

Contraction of Blood Clots Is Impaired in Acute Ischemic Stroke

Valerie Tutwiler, Alina D. Peshkova, Izabella A. Andrianova, Dina R. Khasanova, John W. Weisel, Rustem I. Litvinov

Objective—Obstructive thrombi or thrombotic emboli are the pathogenic basis of ischemic stroke. In vitro blood clots and in vivo thrombi can undergo platelet-driven contraction (retraction), resulting in volume shrinkage. Clot contraction can potentially reduce vessel occlusion and improve blood flow past emboli or thrombi. The aim of this work was to examine a potential pathogenic role of clot contraction in ischemic stroke.

Approach and Results—We used a novel automated method that enabled us to quantify time of initiation and extent and rate of clot contraction in vitro. The main finding is that clot contraction from the blood of stroke patients was reduced compared with healthy subjects. Reduced clot contraction correlated with a lower platelet count and their dysfunction, higher levels of fibrinogen and hematocrit, leukocytosis, and other changes in blood composition that may affect platelet function and properties of blood clots. Platelets from stroke patients were spontaneously activated and displayed reduced responsiveness to additional stimulation. Clinical correlations with respect to severity and stroke pathogenesis suggest that the impaired clot contraction has the potential to be a pathogenic factor in ischemic stroke.

Conclusions—The changeable ability of clots and thrombi to shrink in volume may be a novel unappreciated mechanism that aggravates or alleviates the course and outcomes of ischemic stroke. The clinical importance of clot or thrombus transformations in vivo and the diagnostic and prognostic value of this blood test for clot contraction need further exploration. (*Arterioscler Thromb Vasc Biol.* 2017;37:271-279. DOI: 10.1161/ATVBAHA.116.308622.)

Key Words: blood coagulation ■ clot retraction ■ stroke ■ thrombosis

Stroke is a leading cause of death and disability worldwide^{1,2} and accounts for 1 in every 20 deaths in the United States each year.^{3,4} Despite this clinical importance and numerous studies on circulatory disorders in the brain vessels, the pathogenesis of stroke remains largely unclear, and the results of stroke prophylaxis and treatment are unsatisfactory.⁵ Brain damage caused by abnormalities of cerebral blood flow is often ischemic by nature² and is associated with thrombosis or thromboembolism, indicating involvement of hemostatic reactions.^{6,7} Hypercoagulability and changes in blood rheology in combination with local dysfunction and destruction of endothelium comprise a pathogenic basis for ischemic stroke.^{6,8-10} In addition to disorders of plasma and vascular hemostasis, platelet activation plays a role in ischemic stroke.^{6,10,11} Platelet stimulation and aggregation are followed by release of potent triggers of the coagulation cascade and secondary platelet activation.

See cover image

A main physiological activator of platelets is thrombin, an enzyme that forms in the blood irrespective of the initial cause of intravascular blood clotting. Concomitant with platelet activation, thrombin converts fibrinogen, a soluble plasma protein, into

an insoluble fibrin gel made of a filamentous network.¹²⁻¹⁴ Fibrin sticks to activated platelets via the integrin receptor $\alpha\text{IIb}\beta 3$ to form a platelet-fibrin meshwork comprising the structural basis of a hemostatic clot or an obstructive thrombus.^{12,15,16} When the blood clot is formed in a tube, it undergoes volume shrinkage, and liquid serum is expelled. This process is called clot contraction or retraction.¹⁷ The cellular and molecular composition of the blood clot has been found to influence the rate and extent of clot contraction.¹⁸ Importantly, contraction occurs not only in vitro but also inside a vessel if a clot or thrombus is formed.¹⁹ The importance of clot contraction in vivo is based on at least 2 mechanisms: (1) at the site of tissue injury, the active decrease in clot size reduces blood loss by pulling together wound edges and (3) blood flow is recovered past a contracted formerly obstructive thrombus, the latter being hypothetically an important compensatory or adaptive mechanism in thrombotic states.^{17,19,20}

Despite potential medical significance, clot contraction has not been studied systematically in clinical settings mainly because of no standardized and automated methods for registration and quantification of this process. Previously, a limited number of studies assessed extent of clot contraction in vitro subjectively as a visible ratio of the volume of serum

Received on: June 10, 2016; final version accepted on: November 17, 2016.

From the Department of Cell and Developmental Biology, University of Pennsylvania School of Medicine, Philadelphia (V.T., J.W.W., R.I.L.); School of Biomedical Engineering, Sciences, and Health Systems, Drexel University, Philadelphia, PA (V.T.); Institute of Fundamental Medicine and Biology, Kazan Federal University, Russia (A.D.P., I.A.A., R.I.L.); and Neurology Department, Interregional Clinical Diagnostic Center, Kazan, Russia (D.R.K.).

The online-only Data Supplement is available with this article at <http://atvb.ahajournals.org/lookup/suppl/doi:10.1161/ATVBAHA.116.308622/-/DC1>.

Correspondence to Rustem I. Litvinov, MD, PhD, Department of Cell and Developmental Biology, University of Pennsylvania School of Medicine, 421 Curie Blvd, Biomedical Research Bldg II/III, Room 1109, Philadelphia, PA 19104. E-mail litvinov@mail.med.upenn.edu

© 2016 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol is available at <http://atvb.ahajournals.org>

DOI: 10.1161/ATVBAHA.116.308622

Nonstandard Abbreviations and Acronyms	
NIHSS	National Institutes of Health Stroke Scale
RBC	red blood cell
TF	tissue factor
TIA	transient ischemic stroke
TOAST	Trial of Org 10172 in Acute Stroke Treatment
TRAP	thrombin receptor-activating peptide

and the reduced volume of the clot. It is not surprising that the parameters of clot contraction under various circumstances are hardly comparable and highly scattered with the normal range of the extent of contraction varying from 20% to 90%. In this work, we use a recently described instrumental technique¹⁸ that enabled us to quantify the kinetics of blood clot contraction in normal and disease states.

The goal of this work was to evaluate the potential pathogenic importance of clot contraction in acute ischemic stroke by correlating the parameters of clot contraction with clinical and laboratory characteristics.

Materials and Methods

Materials and Methods are available in the [online-only Data Supplement](#).

Results

Comparison of Hemostatic and Hematologic Profiles in Stroke Patients and Healthy Donors

Hemostatic and hematologic parameters for stroke patients and healthy donors are shown in Table 1. Comparison of these parameters revealed an elevation in fibrinogen and D-dimer levels and a decrease in the rate of ADP-induced platelet aggregation in stroke patients (Table 1). In addition, stroke patients had a reduced platelet count and increased fibrinogen level, hematocrit, erythrocyte sedimentation rate, and leukocyte count. In vitro model experiments with variation of the platelet count and fibrinogen levels suggest that clot contraction can be directly modulated by these blood constituents. In particular, clot contraction was enhanced by the increasing number of platelets (Figure IIA in the [online-only Data Supplement](#)), whereas fibrinogen had a dose-dependent inhibitory effect (Figure IIB in the [online-only Data Supplement](#)), consistent with the changes in blood composition observed in the stroke patients. Because the molecular and cellular composition of the blood impacts the process of clot contraction,¹⁸ these differences collectively suggest that clot contraction has the potential to differ in healthy donors and stroke patients.

Characterization of Clot Contraction in Stroke Patients

Automated optical tracking of clot size (Figure 1A and 1B) revealed a 60% reduction in the average extent of clot contraction at the end point in stroke patients relative to healthy donors (Figure 1C), which resulted in stroke patients having a significantly larger clot size. Stroke patients also had a reduced average velocity and area over the curve while having a prolonged lag time (Figure 1D and 1E). Cross-correlation of

Table 1. Hemostatic and Hematologic Parameters in Stroke Patients and Healthy Subjects

Parameters (Normal Ranges Are Shown in Parentheses)	Healthy Subjects (n=79)	Patients With IS (n=85)
Hemostatic parameters		
aPTT (26–36), s	32.1±0.5	35.4±2.4
Prothrombin ratio (70–130), s	103.6±1.5	105.8±2.8
INR (0.85–1.15)	0.99±0.01	1.00±0.02
Fibrinogen (1.8–4.0), g/L	2.7±0.1	3.3±0.2*
Thrombin time (14–21), s	17.7±0.2	17.4±0.4
D-Dimer (0–0.5), µg/mL	0.23±0.03	1.7±0.4‡
Antithrombin III (80–120), %	90.5±1.0	94.1±4.9
Plasminogen (80–135), %	111.8±2.9	109.8±4.3
Protein C (70–130), %	104.3±2.1	103.7±2.9
Maximal ADP-induced platelet aggregation (60–90), %	64.3±1.6	62.0±4.4
Rate of ADP-induced platelet aggregation (30–45), %/min	36.6±1.0	24.3±2.4§
Hematologic parameters		
Platelet count (180–320), ×10 ⁹ /L	286±10	252±8*
Mean platelet volume (8.6–12.6), fL	9.5±1.4	8.4±0.1†
Red blood cells (3–5), ×10 ¹² /L	4.5±0.1	4.5±0.1
Hematocrit (36–48), %	38.5±0.5	41.6±0.6‡
Hemoglobin (120–160), g/L	134.1±1.8	139.3±2.2
Color index (0.85–1.05)	0.93±0.01	0.92±0.01
Mean cell volume (80–100), fL	85.6±0.8	92.2±0.9§
Mean cell hemoglobin (30–35), pg	32.1±0.4	30.9±0.3†
Leukocytes (4–9), ×10 ⁹ /L	5.8±0.2	7.8±0.2§
Eosinophils (0.5–5), %	3.0±0.3	2.0±0.1*
Monocytes (3–11), %	6.2±0.2	8.1±0.3§
Lymphocytes (19–37), %	34±1	23±1§
Basophiles (0–1), %	0.4±0.1	0.49±0.02
Neutrophils (47–78), %	58.8±1.1	66.3±1.0§
ESR (0–15), mm/h	8.5±1.0	19.4±2.1‡

aPTT indicates activated partial thromboplastin time; ESR, erythrocyte sedimentation rate; INR, international normalized ratio; and IS, ischemic stroke. * $P<0.01$, † $P<0.05$, ‡ $P<0.001$, § $P<0.0001$.

clot contraction parameters revealed that extent of clot contraction, average velocity, and area over the curve were all strongly correlated with each other, suggesting a mechanistic relationship between the parameters. Interestingly, although lag time and extent of clot contraction were not significantly correlated in healthy donors, there was a significant negative correlation between these 2 parameters in stroke patients, suggesting that reduced platelet activity increases the lag time and decreases the extent of clot contraction (Tables I and II in the [online-only Data Supplement](#)).

To see whether patients experiencing a transient ischemic stroke (TIA) have more normalized clot retraction, blood




Figure 1. Optical tracking system used for measurements of clot contraction. **A**, The optical analyzer (side view) used to measure light scattering during the process of clot contraction. **B**, The changes in relative clot size are converted into a kinetic curve that can be analyzed for (a) the extent of clot contraction at 20 min, (b) the lag time or the time to 5% contraction, and (c) the area over the curve (AOC) that characterizes the entire process of clot contraction and integrates the extent and rate of contraction and the lag time. Optical tracking was used to assess differences in between healthy subjects and stroke patients in **(C)** extent of clot contraction, **(D)** average velocity of contraction, **(E)** AOC, and **(F)** lag time. Parameters for healthy subjects and stroke patients were compared using an unpaired, 2-tailed Student *t* test. **** $P < 0.0001$.

samples from patients with TIA were analyzed. It was found that TIA patients also had a significantly reduced extent of contraction, average velocity, lag time, and area over the curve ($P < 0.01$ for average velocity, $P < 0.0001$ for all other parameters), whereas clot contraction in TIA patients did not significantly differ from that of ischemic stroke patients (Table III in the [online-only Data Supplement](#)).

Kinetic Phase Analysis of Clot Contraction in Stroke Patients

It has been shown previously that clot contraction in the blood of healthy subjects occurs in 3 phases: initiation of contraction (phase 1), linear contraction (phase 2), and mechanical stabilization (phase 3).¹⁸ Regression analysis conducted on average kinetics curves (Figure 2) revealed that stroke patients and healthy donors did not have a difference in the duration of the different phases or in the rate constant associated with phase 1 (Table 2). However, the rate constants of the linear phase 2 and

the exponential phase 3 were significantly reduced by 2 times and 10 times, respectively, in stroke patients compared with healthy subjects, indicating impairment of the mechanisms of compaction (phase 2) and stabilization of the clots (phase 3).

Correlation Between Contraction Parameters and Laboratory Tests

Because the observed differences between clot contraction in healthy subjects and stroke patients were likely related to disorders of hemostasis and changed blood composition, we investigated correlations between the extent of clot contraction and hemostatic, hematologic, and biochemical parameters (Table IV in the [online-only Data Supplement](#)). The main finding was that the extent of clot contraction moderately but significantly correlated with the platelet count in stroke patients. We also revealed slight but significant correlations between the extent of clot contraction and prothrombin time (prothrombin index and international normalized ratio) and hematocrit in healthy subjects, whereas these correlations were not present in stroke patients. In stroke patients, only monocyte count weakly but significantly correlated with the extent of clot contraction ($r^2 = 0.2839$; $P < 0.001$).

Analysis of Platelet Function and Morphology in Stroke Patients

Because platelets are critical for the contractile force generated during clot retraction, we studied the functionality of platelets in ischemic stroke using flow cytometry–based evaluation of the baseline activity of quiescent platelets and their responsiveness to chemical activation. We used thrombin receptor–activating peptide (TRAP) to mimic the effect of thrombin on protease-activated receptors without a risk of fibrin formation that would complicate the assay. Platelet reactivity was assessed by surface expression of P-selectin and by the ability to bind fibrinogen as a measure of the integrin $\alpha IIb\beta 3$ activation. In addition, isolated unstimulated platelets from ischemic stroke patients were studied with scanning electron microscopy to assess their functional status reflected by their shape. In stroke, platelets were initially activated as revealed by a higher




Figure 2. Averaged kinetic clot contraction curves obtained in the blood of stroke patients ($n = 85$) and healthy subjects ($n = 79$). Optical tracking was used to measure the relative changes in clot size during 20 min with 15-s intervals. Data are represented as mean \pm SEM.

Table 2. Phase Kinetic Parameters of Clot Contraction in Stroke Patients and Healthy Subjects

Rate Constants	Healthy Subjects	Stroke Patients
Phase 1, 1/s	0.023±0.006	0.014±0.005
Phase 2, %/s	-0.073±0.002	-0.0210±0.0003*
Phase 3, 1/s	0.0021±0.0004	0.00029±0.00006*

* $P<0.0001$.

level of P-selectin expression in unstimulated cells (Table 3; Figure III in the [online-only Data Supplement](#)). This observation was confirmed by the frequently observed shape change and formation of filopodia in platelets from stroke patients (34%) compared with morphologically altered platelets from healthy subjects (9%, $P<0.0001$; Figure IV in the [online-only Data Supplement](#)). In response to TRAP-induced stimulation, platelets from the blood of ischemic stroke patients had a significantly lower expression of P-selectin compared with TRAP-activated normal platelets. TRAP-induced stimulation also resulted in reduced fibrinogen-binding capacity compared with the TRAP-activated normal platelets. Collectively, these results indicate that in ischemic stroke, platelets are initially partially activated and have a reduced responsiveness to a thrombin-like stimulus. It is noteworthy that the ratio of activated versus quiescent cells for both fluorescent markers was ≈ 2 - to 4-fold higher in healthy donors compared with stroke patients, suggesting, in combination with the reduced exposure of P-selectin and fibrinogen-binding activity, that the stroke platelets have a substantially decreased overall activation potential. As a matter of fact, contraction in the group of stroke patients studied for platelet functionality was significantly impaired with an average extent of $28\pm 7\%$ versus $51\pm 1\%$ for control ($P<0.0001$).

Analysis of Clot Contraction Based on the Severity of Stroke

The National Institutes of Health Stroke Scale (NIHSS) is a validated clinical methodology to assess the severity of stroke in scale from 0 to 42, where <15 corresponds to a minor-to-moderate stroke and >15 corresponds to a moderate-to-severe or severe stroke. Modified Rankin scale measures rank the level of disability after stroke with 0 representing no symptoms and 5 representing severe disability.²¹ The Alberta Stroke

Program Early CT score value is inversely related to the area of damage as determined through a topographical evaluation of damage.²² Within the cohort of patients in this study, the NIHSS value correlated strongly with modified Rankin scale score and inversely with Alberta Stroke Program Early CT score (Table V in the [online-only Data Supplement](#)).

The stroke patient population was divided based on NIHSS value to compare those with a value <15 to those with a value >15 . We found that patients with more severe strokes (NIHSS >15) had a moderately but significantly increased extent of clot contraction, average velocity, area over the curve, and a reduced lag time when compared with patients with minor and moderate stroke (NIHSS <15 ; Figure 3). It is noteworthy that regardless of stroke severity the patients had a significantly reduced extent of clot contraction compared with healthy subjects, and neither subgroup of stroke patients differed significantly from the total stroke patient population (Tables VI and VII in the [online-only Data Supplement](#)). Hemostatic, hematologic, and biochemical parameters were compared for the populations of stroke patients divided based on NIHSS score (Table VII in the [online-only Data Supplement](#)). Patients with an NIHSS score >15 had a faster activated partial thromboplastin time and increased fibrinogen level, D-dimer level, platelet count, leukocyte count, and protein S-100 level.

Analysis of Clot Contraction Based on Pathogenesis of Stroke

To reveal whether the notable alterations in clot contraction in stroke patients had a pathogenesis-related significance, we investigated differences between subpopulations of stroke patients as defined by the TOAST classification (Trial of Org 10172 in Acute Stroke Treatment; Table 4), which allows for a distinction to be made between the following pathogeneses of ischemic stroke: (1) large-vessel atherosclerosis, (2) cardioembolism, (3) small-vessel occlusion, (4) stroke of other determined pathogenesis, and (5) stroke of undetermined pathogenesis.²³ Because of the small number of patients with TOAST subtypes 3 to 5, we preceded with analysis of clot contraction in patients with cardioembolic and atherothrombotic strokes. Assessment of extent of clot contraction based on stroke subtype revealed that there was a lower extent of clot contraction in patients with cardioembolic strokes when compared with patients with atherothrombotic strokes (Figure 4).

Table 3. Functional Characterization of Platelets Isolated From the Blood of Ischemic Stroke Patients and Healthy Donors Before and After Stimulation of the Cells With TRAP*

	P-Selectin Expression			Fibrinogen-Binding Capacity		
	Quiescent Platelets	TRAP-Activated	Activated/Quiescent Ratio	Quiescent Platelets	TRAP-Activated	Activated/Quiescent Ratio
Healthy donors (n=11)	3.3±0.8	83.6±3.7	≈ 25	1.0±0.3	58.3±5.6	≈ 58
Ischemic stroke patients (n=11)	9.9±5.1†	71.1±7.3‡	≈ 7	1.1±0.4	42.9±7.4§	≈ 39

TRAP indicates thrombin receptor-activating peptide.

*Numbers (mean±SEM) represent relative flow cytometry counts (%) for the fractions of platelets bearing fluorescently labeled fibrinogen or antibodies to P-selectin.

† $P<0.05$ between the donors and patients.‡ $P<0.01$ between the donors and patients.§ $P<0.001$ between the donors and patients.




Figure 3. Clot contraction parameters in the blood of 2 groups of ischemic stroke patients segregated based on the National Institutes of Health Stroke Scale (NIHSS) values: <15 (minor-to-moderate stroke) and >15 (severe stroke). Optical tracking was used to assess differences in (A) extent of clot contraction, (B) average velocity of contraction, (C) area over the curve, and (D) lag time. Parameters for stroke patients were compared using an unpaired, 2-tailed Student *t* test. **P*<0.05, ***P*<0.01.

Discussion

Stroke is a leading cause of death and disability worldwide.⁴ Although the pathogenesis of ischemic stroke is complex and heterogeneous, a common link between the different pathogeneses is increased activity of the coagulation and fibrinolytic systems¹⁰ in conjunction with macro- and microvessel disruption⁹ manifesting in thrombotic events. A thrombus or

thrombotic embolus can result in blockage of blood flow to a portion of the brain; size and location of the thrombotic event influences stroke severity and functional outcome.²⁴ Brain tissue is a prominent source of tissue factor (TF),²⁵ and locations of brain damage are associated with high levels of TF and other procoagulant compounds,²⁶ which can intensify hypercoagulability and potentially lead to exhaustion of anticoagulant mechanisms and functional impairment of platelets.²⁷

Information gleaned from in vitro and ex vivo studies of clot structure has been widely used to inform clinical findings.²⁸ Thrombi collected from ischemic stroke patients have a platelet–fibrin network interspersed with cells, and those rich in red blood cells (RBCs) are generally more susceptible to thrombolysis.²⁹ In ischemic stroke, the structure of thrombi is nonhomogenous, but, on average, RBCs comprise 34% of the volume,³⁰ which can influence the process of clot contraction.¹⁸ Interestingly, fibrin clots formed in the blood from ischemic stroke patients had abnormal fibrin ultrastructure that could influence contraction of the fibrin–platelet network.²⁸ Collectively, this provides the need for an examination of clot contraction in samples collected from patients with ischemic stroke.

Blood clot contraction remains the least studied stage of blood clotting. This work is the first systematic assessment of clot contraction in ischemic stroke. The main finding of this work is that clots formed from the blood of patients with ischemic stroke have a reduced extent of clot contraction when compared with clots formed from the blood of healthy subjects (Figure 1). This result is somewhat unexpected as ischemic stroke is associated with a procoagulant activity,⁹ but the aggregate of laboratory values collected from this cohort of ischemic stroke patients sheds some light on potential mechanisms for impaired clot contraction.

Platelets contain actomyosin machinery that generates contractile forces^{31–33} that are propagated through the clot via

Table 4. Clinical Classification of Stroke Patients Enrolled in This Study

TOAST Classification				
	Cardioembolic	Large Artery Atherosclerotic	Small-Vessel Occlusion	Total Patients
Number of patients, n (%)	33 (39)	45 (54)	6 (7)	84 (100)
Male sex, n (%)	13 (39)	25 (56)	3 (50)	41 (48)
Age, y	69±2	68±1	60±4	68±1
Ischemic stroke classifications				
NIHSS score at the time of admission (median; 1st, 3rd quartile)	9; 4.5, 17	7; 3, 13	4; 3.5, 6	7; 4; 13.5
mRs score (median; 1st, 3rd quartile)	3; 2, 4.5	3; 2, 4	1; 0.75, 1.75	3; 2, 4
ASPECT score (median; 1st, 3rd quartile)	7; 3, 8	8; 3, 8	9; 9, 9	7; 3.5, 8.5
OCSP classification, n (%)				
TACI	9 (27)	10 (22)	0	19 (23)
PACI	20 (60)	21 (47)	1 (17)	42 (50)
LACI	4 (12)	13 (29)	2 (33)	19 (23)
POCI	0 (0)	1 (2)	3 (50)	4 (5)

ASPECT indicates Alberta Stroke Program Early CT; LACI, lacunar infarct; mRs, modified Rankin scale; NIHSS, National Institutes of Health Stroke Scale; OCSP, Oxfordshire Community Stroke Project; PACI, partial anterior circulation infarct; POCI, posterior circulation infarct; TACI, total anterior circulation infarct; and TOAST, Trial of Org 10172 in Acute Stroke Treatment.




Figure 4. Clot contraction parameters in the blood of patients of clinical subpopulations of ischemic stroke as defined by TOAST classification (Trial of Org 10172 in Acute Stroke Treatment). The extent of clot contraction is reduced in patients with cardioembolic type strokes (n=33) compared with atherothrombotic type strokes (n=45). * $P < 0.05$.

platelet–fibrin interactions.^{34,35} Therefore, reduced contraction in stroke patients is likely because of reduced number and functionality of platelets (Table 1). We confirmed the mechanistic importance of the quantitative and qualitative changes in platelets for clot contraction through in vitro experiments with varying platelet counts (Figure IIA in the [online-only Data Supplement](#)) and through flow cytometry (Table 3; Figure III in the [online-only Data Supplement](#)) and scanning electron microscopy (Figure IV in the [online-only Data Supplement](#)) of platelets from stroke patients. On the basis of the surface expression of P-selectin, fibrinogen-binding capacity, and morphology, it has been found that platelets in the blood of stroke patients are initially activated and, more importantly, have a dramatically reduced capacity to respond to a thrombin-like activating stimulus. This is consistent with an earlier report that states that platelets from ischemic stroke patients are insensitive to thrombin in vitro,²⁷ which may result from cleavage of protease-activated receptors, metabolic exhaustion, or other mechanisms. Although a reduced platelet count in stroke is moderate (Table 1) and can have a relatively minor contribution to impaired clot contraction, it is platelet dysfunction that is likely to be the major pathogenic mechanism for reduced clot contraction observed in the blood of stroke patients. Disruption in both platelet number and function results in partial contractile inefficiency.

The prolonged lag phase observed in ischemic stroke (Figure 1) could be, in part, because of increase of anti-thrombin III activity after activation of the clotting cascade (Table 1), which inhibits thrombin and other clotting factors,³⁶ resulting in the slightly prolonged activated partial thromboplastin time observed in stroke patients (Table 1). Reduced thrombin activity strongly correlates with impaired clot contraction and could be a factor here despite addition of exogenous thrombin to initiate clotting.¹⁸

Increased fibrinogen level that can accompany ischemic stroke³⁷ (Table 1) can suppress clot contraction, which was confirmed in the in vitro experiments (Figure IIB in the [online-only Data Supplement](#)). Higher fibrinogen concentration results in increased fibrin mass and more fibers not associated with activated platelets/platelet aggregates, which

can impede clot contraction.^{12,18} In stroke patients, fibrin and RBCs can undergo oxidative and proteolytic changes that result in the formation of densely matted deposits of fibrin and RBCs, which have been implicated to play a role in persistent thrombi.^{38,39} The abnormal fibrin structure associated with ischemic stroke^{40–42} results in lower clot permeability and a slower rate of lysis.⁴²

Contractile forces that are generated by platelets are able to compress RBCs tightly into the core of clots.¹⁹ Compacted RBCs form a tessellated network and acquire a polyhedral shape, resulting in the terminology polyhedrocytes.¹⁹ Polyhedrocytes have been found inside coronary thrombi extracted from patients with myocardial infarction,⁴³ confirming that intravascular clots undergo contraction. Increased hematocrit results in a reduced extent and rate of clot contraction,¹⁸ and the higher hematocrit in stroke patients (Tables 1 and 2; Figure 2) is consistent with this inhibitory effect and could be additional mechanism of impaired contraction in stroke patients.

Based on this data analysis, we propose that collectively the reduced platelet count, increased hematocrit, and increased fibrinogen contribute to the impaired mechanism of clot contraction, which results in vessel occlusion and brain infarct. The compromised contraction of an intravascular clot or thrombus may play an important role in the pathogenesis of ischemic stroke. Although severity of stroke and infarct size are largely determined by the diameter and location of the occluded vessel, the ability of the thrombi to contract either more or less can aggravate or alleviate the course of ischemic brain damage. For example, if clot or thrombus contraction results in a 50% reduction in the vessel cross-sectional area, this would translate to a reduction of blood flow to 24% of that with no clot, whereas the impaired contraction seen in the stroke patients, resulting in an 80% reduction in the vessel cross-sectional area, would reduce the flow rate to 4% of the original.

Interestingly, clot contraction is also reduced in TIA similar to the reduction seen in ischemic stroke patients (Table III in the [online-only Data Supplement](#)), indicating that clot contraction is impaired in these patients. TIA is characterized as a focal neurological deficit that lasts <24 hours and, although ischemic in nature, does not result in infarction.⁴⁴ Patients with a recent TIA are at a high risk for the occurrence of an ischemic stroke,^{45,46} and 15% of ischemic stroke patients presented with a previous TIA.⁴⁷ Our findings suggest that a premonitory status and predisposition to thrombosis, such as are seen in TIA and ischemic stroke, are critical determinants for the ability of clots to contract rather than a transitory episode of cerebral vascular occlusion and ischemia that may or may not result in persistent thrombosis or thromboembolism. Although rapid treatment of TIA has the potential to help reduce the incidence of ischemic stroke,^{48,49} the TIA samples in our study were examined before the administration of any anticoagulant, thrombolytic, or antiplatelet drugs.

Our results show that despite an overall impairment of contraction in stroke patients, severe strokes had a relative increase in the extent of clot contraction compared with minor/moderate stroke (Figure 4). NIHSS score >15 also had a higher Protein S-100 level, a molecular marker of brain damage,

which is correlated with infarct volume, functional outcome, and ongoing ischemia.⁵⁰ We propose that in some cases of stroke with focal brain damage, the contractile activity of clots and thrombi can be improved as a potential compensatory mechanism to decrease the consequences of thrombosis and restore blood flow. In patients with severe ischemic brain damage, the blood–brain barrier is disrupted,⁵¹ and massive amounts of TF are released into systemic circulation, resulting in activation of the extrinsic pathway of blood coagulation.^{52–54} It has been shown that the level of brain damage is associated with the amount of TF that is released.^{55,56} Also, ischemic stroke patients have increased TF-bearing microparticles⁵⁷ and TF expression on monocytes.⁵⁸

We hypothesize that increased TF could potentially cause a secondary wave of thrombin generation, resulting in a faster activated partial thromboplastin time (Table VI in the [online-only Data Supplement](#)) and higher fibrinogen turnover in severe stroke patients.⁵⁹ Thrombin can induce highly active platelets in severe ischemic stroke patients, which can be determined through expression of P-selectin.^{60,61} Platelet hyperactivity in combination with increased platelet count (Table VI in the [online-only Data Supplement](#)) could explain the increased extent of contraction in patients with more severe stroke compared with less severe stroke (Figure 3).

We propose a dual role of clot contraction in the ischemic stroke (Graphical Abstract in the [online-only Data Supplement](#)). Impaired clot contraction results because of changes in blood composition. This impaired clot contraction has the potential to be a pathogenic factor that aggravates the course of disease by increased blockage of blood flow. Enhanced clot contraction occurs in atherothrombotic patients compared with cardioembolic patients (Figure 4). This supports the idea that TF can potentially enhance contraction in subpopulations of stroke patients because atherothrombotic stroke patients have increased TF⁶² and D-dimer, creating a thrombogenic state,⁶³ and increased P-selectin expression on platelets, indicative of hyperactivity.^{64,65} Cardioembolic strokes are not a result of platelet hyperfunction but rather arise because of blood stasis.⁶⁶ This in conjunction with marginally enhanced clot contraction in severe strokes supports the notion that an increased thrombogenic state may result in the enhanced contraction observed here. This enhanced clot contraction has the potential to act as a compensatory or adaptive factor in ischemic stroke. However, this potential requires further examination.

In summary, the pathophysiological significance of clot contraction in thrombotic states has been explored. Clot contraction is associated with the restoration of blood flow past otherwise obstructive thrombi, potentially influencing the outcome of thrombosis.^{17,19,20} However, compressed, dense thrombi are more impermeable and may be resistant to fibrinolysis. Consequently, clinical consequences of clot contraction in vivo may be ambiguous, resulting in either aggravation or relief of the thrombotic state. Our results reveal that further systematic studies of the clinical importance of clot contraction are worthwhile, as they could inform a novel pathogenic mechanism and result in potential laboratory tests that improve diagnosis of and prognosis in thrombotic conditions such as ischemic stroke.

Acknowledgments

We thank Dr Fazly Ataullakhanov (HemaCore, Ltd) for providing the Thrombodynamics Analyser System used in this study and Drs Timur Demin and Marat Saikhunov for providing clinical data. Scanning electron microscopy was performed at the Interdisciplinary Center for Analytical Microscopy of Kazan Federal University.

Sources of Funding

The work was supported by National Institutes of Health grants HL-090774, UO1HL116330, and T32 H10791, National Science Foundation grant DMR1505662, the Program for Competitive Growth at Kazan Federal University, and American Heart Association grant 16PRE30260002 (V. Tutwiler).

Disclosures

None.

References

- Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet*. 2006;367:1747–1757. doi: 10.1016/S0140-6736(06)68770-9.
- Feigin VL, Lawes CM, Bennett DA, Barker-Collo SL, Parag V. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *Lancet Neurol*. 2009;8:355–369. doi: 10.1016/S1474-4422(09)70025-0.
- Centers for Disease Control and Prevention. Stroke statistics and maps. Centers for Disease Control and Prevention: stroke website. http://www.cdc.gov/stroke/statistics_maps.htm. Updated 2015. Accessed March 15, 2015.
- Mozaffarian D, Benjamin EJ, Go AS, et al. Executive summary: heart disease and stroke statistics - 2016 update. *Circulation*. 2016;133:447–454.
- Wiseman SJ, Doubal FN, Chappell FM, Valdés-Hernández MC, Wang X, Rumley A, Lowe GD, Dennis MS, Wardlaw JM. Plasma biomarkers of inflammation, endothelial function and hemostasis in cerebral small vessel disease. *Cerebrovasc Dis*. 2015;40:157–164. doi: 10.1159/000438494.
- Stoll G, Kleinschnitz C, Nieswandt B. Molecular mechanisms of thrombus formation in ischemic stroke: novel insights and targets for treatment. *Blood*. 2008;112:3555–3562. doi: 10.1182/blood-2008-04-144758.
- Ruggeri ZM. Platelets in atherothrombosis. *Nat Med*. 2002;8:1227–1234. doi: 10.1038/nm1102-1227.
- Levine SR. Hypercoagulable states and stroke: a selective review. *CNS Spectr*. 2005;10:567–578.
- Blum A, Vaispapir V, Keinan-Boker L, Soboh S, Yehuda H, Tamir S. Endothelial dysfunction and procoagulant activity in acute ischemic stroke. *J Vasc Interv Neurol*. 2012;5:33–39.
- Fisher M, Francis R. Altered coagulation in cerebral ischemia. Platelet, thrombin, and plasmin activity. *Arch Neurol*. 1990;47:1075–1079.
- Shen MY, Chen FY, Hsu JF, et al. Plasma L5 levels are elevated in ischemic stroke patients and enhance platelet aggregation. *Blood*. 2016;127:1336–1345. doi: 10.1182/blood-2015-05-646117.
- Weisel JW, Litvinov RI. Mechanisms of fibrin polymerization and clinical implications. *Blood*. 2013;121:1712–1719. doi: 10.1182/blood-2012-09-306639.
- Brown AE, Litvinov RI, Discher DE, Purohit PK, Weisel JW. Multiscale mechanics of fibrin polymer: gel stretching with protein unfolding and loss of water. *Science*. 2009;325:741–744. doi: 10.1126/science.1172484.
- Kim OV, Litvinov RI, Weisel JW, Alber MS. Structural basis for the nonlinear mechanics of fibrin networks under compression. *Biomaterials*. 2014;35:6739–6749. doi: 10.1016/j.biomaterials.2014.04.056.
- Bodin S, Soulet C, Tronchère H, Sié P, Gachet C, Plantavid M, Payrastre B. Integrin-dependent interaction of lipid rafts with the actin cytoskeleton in activated human platelets. *J Cell Sci*. 2005;118(pt 4):759–769. doi: 10.1242/jcs.01648.
- Podolnikova NP, Yakovlev S, Yakubenko VP, Wang X, Gorkun OV, Ugarova TP. The interaction of integrin α IIb β 3 with fibrin occurs through multiple binding sites in the α IIb β -propeller domain. *J Biol Chem*. 2014;289:2371–2383. doi: 10.1074/jbc.M113.518126.
- Carr ME. Development of platelet contractile force as a research and clinical measure of platelet function. *Cell Biochem Biophys*. 2003;38:55–78.
- Tutwiler V, Litvinov RI, Lozhkin AP, Peshkova AD, Lebedeva T, Ataullakhanov FI, Spiller KL, Cines DB, Weisel JW. Kinetics and

- mechanics of clot contraction are governed by the molecular and cellular composition of the blood. *Blood*. 2016;127:149–159. doi: 10.1182/blood-2015-05-647560.
19. Cines DB, Lebedeva T, Nagaswami C, Hayes V, Massefski W, Litvinov RI, Rauova L, Lowery TJ, Weisel JW. Clot contraction: compression of erythrocytes into tightly packed polyhedra and redistribution of platelets and fibrin. *Blood*. 2014;123:1596–1603. doi: 10.1182/blood-2013-08-523860.
 20. Muthard RW, Diamond SL. Blood clots are rapidly assembled hemodynamic sensors: flow arrest triggers intraluminal thrombus contraction. *Arterioscler Thromb Vasc Biol*. 2012;32:2938–2945. doi: 10.1161/ATVBAHA.112.300312.
 21. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke*. 1988;19:604–607.
 22. Pexman JH, Barber PA, Hill MD, Sevick RJ, Demchuk AM, Hudon ME, Hu WY, Buchan AM. Use of the Alberta Stroke Program Early CT Score (ASPECTS) for assessing CT scans in patients with acute stroke. *AJNR Am J Neuroradiol*. 2001;22:1534–1542.
 23. Adams HP, Beneditin BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE 3rd. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke*. 1993;24:35–41.
 24. Puetz V, Dzialowski I, Hill MD, Subramaniam S, Sylaja PN, Krol A, O'Reilly C, Hudon ME, Hu WY, Coutts SB, Barber PA, Watson T, Roy J, Demchuk AM; Calgary CTA Study Group. Intracranial thrombus extent predicts clinical outcome, final infarct size and hemorrhagic transformation in ischemic stroke: the clot burden score. *Int J Stroke*. 2008;3:230–236. doi: 10.1111/j.1747-4949.2008.00221.x.
 25. del Zoppo GJ, Yu JQ, Copeland BR, Thomas WS, Schneiderman J, Morrissey JH. Tissue factor localization in non-human primate cerebral tissue. *Thromb Haemost*. 1992;68:642–647.
 26. Thomas WS, Mori E, Copeland BR, Yu JQ, Morrissey JH, del Zoppo GJ. Tissue factor contributes to microvascular defects after focal cerebral ischemia. *Stroke*. 1993;24:847–853; discussion 847.
 27. Jurk K, Jahn UR, Van Aken H, Schriek C, Droste DW, Ritter MA, Bernd Ringelstein E, Kehrel BE. Platelets in patients with acute ischemic stroke are exhausted and refractory to thrombin, due to cleavage of the seven-transmembrane thrombin receptor (PAR-1). *Thromb Haemost*. 2004;91:334–344. doi: 10.1160/TH03-01-0044.
 28. Undas A, Slowik A, Wolkow P, Szczudlik A, Tracz W. Fibrin clot properties in acute ischemic stroke: relation to neurological deficit. *Thromb Res*. 2010;125:357–361. doi: 10.1016/j.thromres.2009.11.013.
 29. Marder VJ, Chute DJ, Starkman S, et al. Analysis of thrombi retrieved from cerebral arteries of patients with acute ischemic stroke. *Stroke*. 2006;37:2086–2093. doi: 10.1161/01.STR.0000230307.03438.94.
 30. Liebeskind DS, Sanossian N, Yong WH, et al. CT and MRI early vessel signs reflect clot composition in acute stroke. *Stroke*. 2011;42:1237–1243. doi: 10.1161/STROKEAHA.110.605576.
 31. Greulich PE, Brouse CF, Beckham J, Jessen ME, Martin EJ, Carr ME. Reductions in platelet contractile force correlate with duration of cardiopulmonary bypass and blood loss in patients undergoing cardiac surgery. *Thromb Res*. 2002;105:523–529.
 32. Lam WA, Chaudhuri O, Crow A, Webster KD, Li TD, Kita A, Huang J, Fletcher DA. Mechanics and contraction dynamics of single platelets and implications for clot stiffening. *Nat Mater*. 2011;10:61–66. doi: 10.1038/nmat2903.
 33. Niederman R, Pollard TD. Human platelet myosin. II. *In vitro* assembly and structure of myosin filaments. *J Cell Biol*. 1975;67:72–92.
 34. Ehrlicher A, Hartwig JH. Cell mechanics: contracting to stiffness. *Nat Mater*. 2011;10:12–13. doi: 10.1038/nmat2928.
 35. Wufsus AR, Rana K, Brown A, Dorgan JR, Liberatore MW, Neeves KB. Elastic behavior and platelet retraction in low- and high-density fibrin gels. *Biophys J*. 2015;108:173–183. doi: 10.1016/j.bpj.2014.11.007.
 36. Hatton MW, Regoeczi E. The inactivation of thrombin and plasmin by antithrombin III in the presence of sepharose-heparin. *Thromb Res*. 1977;10:645–660.
 37. Kristensen B, Malm J, Nilsson TK, Hultdin J, Carlberg B, Olsson T. Increased fibrinogen levels and acquired hypofibrinolysis in young adults with ischemic stroke. *Stroke*. 1998;29:2261–2267.
 38. Pretorius E, Lipinski B. Thromboembolic ischemic stroke changes red blood cell morphology. *Cardiovasc Pathol*. 2013;22:241–242. doi: 10.1016/j.carpath.2012.11.005.
 39. Lipinski B, Pretorius E, Oberholzer HM, van der Spuy WJ. Interaction of fibrin with red blood cells: the role of iron. *Ultrastruct Pathol*. 2012;36:79–84. doi: 10.3109/01913123.2011.627491.
 40. Pretorius E, Swanepoel AC, Oberholzer HM, van der Spuy WJ, Duim W, Wessels PF. A descriptive investigation of the ultrastructure of fibrin networks in thrombo-embolic ischemic stroke. *J Thromb Thrombolysis*. 2011;31:507–513. doi: 10.1007/s11239-010-0538-5.
 41. Pretorius E, Steyn H, Engelbrecht M, Swanepoel AC, Oberholzer HM. Differences in fibrin fiber diameters in healthy individuals and thromboembolic ischemic stroke patients. *Blood Coagul Fibrinolysis*. 2011;22:696–700. doi: 10.1097/MBC.0b013e32834dbd32.
 42. Pera J, Undas A, Topor-Madry R, Jagiella J, Klimkowicz-Mrowiec A, Slowik A. Fibrin clot properties in acute stroke: what differs cerebral hemorrhage from cerebral ischemia? *Stroke*. 2012;43:1412–1414. doi: 10.1161/STROKEAHA.111.646729.
 43. Ząbczyk M, Sadowski M, Zalewski J, Undas A. Polyhydrocytes in intracoronary thrombi from patients with ST-elevation myocardial infarction. *Int J Cardiol*. 2015;179:186–187. doi: 10.1016/j.ijcard.2014.10.004.
 44. Albers GW, Caplan LR, Easton JD, Fayad PB, Mohr JP, Saver JL, Sherman DG; TIA Working Group. Transient ischemic attack—proposal for a new definition. *N Engl J Med*. 2002;347:1713–1716. doi: 10.1056/NEJMs020987.
 45. Wu CM, McLaughlin K, Lorenzetti DL, Hill MD, Manns BJ, Ghali WA. Early risk of stroke after transient ischemic attack: a systematic review and meta-analysis. *Arch Intern Med*. 2007;167:2417–2422. doi: 10.1001/archinte.167.22.2417.
 46. Coull AJ, Lovett JK, Rothwell PM; Oxford Vascular Study. Population based study of early risk of stroke after transient ischaemic attack or minor stroke: implications for public education and organisation of services. *BMJ*. 2004;328:326. doi: 10.1136/bmj.37991.635266.44.
 47. Brainin M, McShane LM, Steiner M, Dachenhausen A, Seiser A. Silent brain infarcts and transient ischemic attacks. A three-year study of first-ever ischemic stroke patients: the Klosterneuburg Stroke Data Bank. *Stroke*. 1995;26:1348–1352.
 48. Dutta D, Bowen E, Foy C. Four-year follow-up of transient ischemic attacks, strokes, and mimics: a retrospective transient ischemic attack clinic cohort study. *Stroke*. 2015;46:1227–1232. doi: 10.1161/STROKEAHA.114.008632.
 49. Hong KS, Yegiaian S, Lee M, Lee J, Saver JL. Declining stroke and vascular event recurrence rates in secondary prevention trials over the past 50 years and consequences for current trial design. *Circulation*. 2011;123:2111–2119. doi: 10.1161/CIRCULATIONAHA.109.934786.
 50. Nash DL, Bellolio MF, Stead LG. S100 as a marker of acute brain ischemia: a systematic review. *Neurocrit Care*. 2008;8:301–307. doi: 10.1007/s12028-007-9019-x.
 51. Kassner A, Merali Z. Assessment of blood-brain barrier disruption in stroke. *Stroke*. 2015;46:3310–3315. doi: 10.1161/STROKEAHA.115.008861.
 52. Vecht CJ, Sibinga CT, Minderhoud JM. Disseminated intravascular coagulation and head injury. *J Neurol Neurosurg Psychiatry*. 1975;38:567–571.
 53. Pathak A, Dutta S, Marwaha N, Singh D, Varma N, Mathuriya SN. Change in tissue thromboplastin content of brain following trauma. *Neurol India*. 2005;53:178–182.
 54. Undas A, Ariëns RA. Fibrin clot structure and function: a role in the pathophysiology of arterial and venous thromboembolic diseases. *Arterioscler Thromb Vasc Biol*. 2011;31:e88–e99. doi: 10.1161/ATVBAHA.111.230631.
 55. Scherer RU, Spangenberg P. Procoagulant activity in patients with isolated severe head trauma. *Crit Care Med*. 1998;26:149–156.
 56. Maegele M. Coagulopathy after traumatic brain injury: incidence, pathogenesis, and treatment options. *Transfusion*. 2013;53(suppl 1):28S–37S. doi: 10.1111/trf.12033.
 57. Świtońska M, Stomka A, Sinkiewicz W, Żekanowska E. Tissue-factor-bearing microparticles (MPs-TF) in patients with acute ischaemic stroke: the influence of stroke treatment on MPs-TF generation. *Eur J Neurol*. 2015;22:395–401. e28. doi: 10.1111/ene.12591.
 58. Kappelmayer J, Berecki D, Misz M, Oláh L, Fekete I, Csiba L, Blaskó G. Monocytes express tissue factor in young patients with cerebral ischemia. *Cerebrovasc Dis*. 1998;8:235–239.
 59. Barber M, Langhorne P, Rumley A, Lowe GD, Stott DJ. Hemostatic function and progressing ischemic stroke: D-dimer predicts early clinical progression. *Stroke*. 2004;35:1421–1425. doi: 10.1161/01.STR.0000126890.63512.41.
 60. Cha JK, Jeong MH, Kim EK, Lim YJ, Ha BR, Kim SH, Kim JW. Surface expression of P-selectin on platelets is related with clinical worsening in acute ischemic stroke. *J Korean Med Sci*. 2002;17:811–816. doi: 10.3346/jkms.2002.17.6.811.
 61. Vanschoonbeek K, Feijge MA, Keuren JF, Coenraad Hemker H, Lodder JJ, Hamulyák K, van Pampus EC, Heemskerk JW. Thrombin-induced hyperactivity of platelets of young stroke patients: involvement of thrombin

- receptors in the subject-dependent variability in Ca²⁺ signal generation. *Thromb Haemost.* 2002;88:931–937.
62. He M, Wen Z, He X, Xiong S, Liu F, Xu J, Li J, Xie Q, Jian Z, Chen F, Xiao B, Pu X, He S. Observation on tissue factor pathway and some other coagulation parameters during the onset of acute cerebrocardiac thrombotic diseases. *Thromb Res.* 2002;107:223–228.
 63. Reganon E, Vila V, Martínez-Sales V, Vaya A, Lago A, Alonso P, Aznar J. Association between inflammation and hemostatic markers in atherothrombotic stroke. *Thromb Res.* 2003;112:217–221. doi: 10.1016/j.thromres.2003.12.008.
 64. Yamazaki M, Uchiyama S, Iwata M. Measurement of platelet fibrinogen binding and p-selectin expression by flow cytometry in patients with cerebral infarction. *Thromb Res.* 2001;104:197–205.
 65. Zeller JA, Tschoepe D, Kessler C. Circulating platelets show increased activation in patients with acute cerebral ischemia. *Thromb Haemost.* 1999;81:373–377.
 66. Nagao T, Hamamoto M, Kanda A, Tsuganesawa T, Ueda M, Kobayashi K, Miyazaki T, Terashi A. Platelet activation is not involved in acceleration of the coagulation system in acute cardioembolic stroke with nonvalvular atrial fibrillation. *Stroke.* 1995;26:1365–1368.

Highlights

- Blood clot contraction is an understudied aspect of blood clotting and thrombosis, and this work is the systematic study of clot contraction in a thrombotic state.
- Extent of clot contraction is reduced in the blood of ischemic stroke patients compared with healthy subjects.
- The clot contraction in ischemic stroke is likely weakened because of platelet dysfunction combined with altered blood composition.
- Impaired clot contraction may be an important pathogenic factor in thrombosis.