

## Adult Basic Life Support

### 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations

**ABSTRACT:** This 2020 *International Consensus on Cardiopulmonary Resuscitation (CPR) and Emergency Cardiovascular Care Science With Treatment Recommendations* on basic life support summarizes evidence evaluations performed for 22 topics that were prioritized by the Basic Life Support Task Force of the International Liaison Committee on Resuscitation. The evidence reviews include 16 systematic reviews, 5 scoping reviews, and 1 evidence update. Per agreement within the International Liaison Committee on Resuscitation, new or revised treatment recommendations were only made after a systematic review.

Systematic reviews were performed for the following topics: dispatch diagnosis of cardiac arrest, use of a firm surface for CPR, sequence for starting CPR (compressions-airway-breaths versus airway-breaths-compressions), CPR before calling for help, duration of CPR cycles, hand position during compressions, rhythm check timing, feedback for CPR quality, alternative techniques, public access automated external defibrillator programs, analysis of rhythm during chest compressions, CPR before defibrillation, removal of foreign-body airway obstruction, resuscitation care for suspected opioid-associated emergencies, drowning, and harm from CPR to victims not in cardiac arrest.

The topics that resulted in the most extensive task force discussions included CPR during transport, CPR before calling for help, resuscitation care for suspected opioid-associated emergencies, feedback for CPR quality, and analysis of rhythm during chest compressions. After discussion of the scoping reviews and the evidence update, the task force prioritized several topics for new systematic reviews.

Theresa M. Olasveengen,  
MD, PhD

:

Peter T. Morley, MBBS,  
AFRACMA, GCertClinTeac  
On behalf of the Adult  
Basic Life Support  
Collaborators

The full author list is available on page  
S77.

**Key Words:** AHA Scientific Statements  
■ cardiopulmonary resuscitation  
■ defibrillators ■ drowning  
■ emergency medical services  
communication systems ■ heart arrest  
■ heart massage ■ respiration, artificial

© 2020 American Heart Association,  
Inc., European Resuscitation Council,  
and International Liaison Committee on  
Resuscitation

<https://www.ahajournals.org/journal/circ>

## CONTENTS

|   |     |
|---|-----|
| Abstract.....   | S41 |
| Topics Reviewed in This 2020 BLS CoSTR.....   | S43 |
| Early Access and Cardiac Arrest Prevention,<br>Including Emergency Medical Dispatch and DA-CPR...                       | S44 |
| Dispatch Diagnosis of Cardiac Arrest<br>(BLS 740: SysRev).....  | S44 |
| Dispatcher Instructions in CPR<br>(2019 CoSTR BLS 359: SysRev).....   | S46 |
| DA-Assisted Compression-Only<br>CPR Versus Conventional CPR<br>(2017 CoSTR BLS 359: SysRev).....                        | S46 |
| Compression-Only CPR .....  | S47 |
| Lay Rescuer Chest Compression-Only Versus<br>Standard CPR (2017 CoSTR BLS 547 SysRev) ....                              | S47 |
| EMS Chest Compression-Only Compared<br>With Conventional CPR (2017 CoSTR BLS 360:<br>SysRev) .....                      | S47 |
| In-Hospital Chest Compression-Only CPR<br>Versus Conventional CPR (2017 CoSTR BLS 372:<br>SysRev).....                  | S48 |
| Rescuer Fatigue in Chest Compression-Only<br>CPR (BLS 349: ScopRev) .....   | S48 |
| CPR Sequence.....   | S48 |
| Firm Surface for CPR (BLS 370: SysRev) .....  | S48 |
| Starting CPR (C-A-B Compared With A-B-C)<br>(BLS 661: SysRev) .....   | S50 |
| CPR Before Call for Help (BLS 1527: SysRev) .....   | S51 |
| Duration of CPR Cycles (2 Minutes Versus<br>Other) (BLS 346: SysRev).....   | S52 |
| Check for Circulation During BLS<br>(BLS 348: EvUp).....  | S54 |
| Components of High-Quality CPR .....  | S54 |
| Hand Position During Compressions<br>(BLS 357: SysRev).....   | S54 |
| Chest Compression Rate, Chest Compression<br>Depth, and Chest Wall Recoil (BLS 366,<br>BLS 367, BLS 343: ScopRev) ..... | S55 |
| Compression-to-Ventilation Ratio<br>(2017 CoSTR BLS 362: SysRev).....   | S56 |
| Timing of Rhythm Check (BLS 345: SysRev) .....  | S56 |
| Feedback for CPR Quality (BLS 361: SysRev).....   | S57 |
| Alternative Techniques.....   | S61 |
| Alternative Techniques (Cough CPR, Precordial<br>Thump, Fist Pacing) (BLS 374: SysRev).....                             | S61 |
| Defibrillation.....   | S64 |
| Public Access AED Programs (BLS 347:<br>SysRev) .....   | S64 |
| Analysis of Rhythm During Chest<br>Compressions (BLS 373: SysRev) .....   | S65 |
| CPR Before Defibrillation (BLS 363: SysRev).....  | S66 |
| Paddle Size and Placement for Defibrillation<br>(ALS-E-030A: ScopRev) .....   | S67 |
| Special Circumstances.....  | S68 |

|  |     |
|--|-----|
| CPR During Transport (BLS 1509: ScopRev).....  | S68 |
| Removal of Foreign-Body Airway<br>Obstruction (BLS 368: SysRev).....                       | S69 |
| Resuscitation Care for Suspected Opioid-<br>Associated Emergencies (BLS 811: SysRev) ..... | S72 |
| Drowning (BLS 856: SysRev) .....   | S73 |
| Potential Harm From CPR.....   | S76 |
| Harm From CPR to Victims not in Cardiac<br>Arrest (BLS 353: SysRev) .....                  | S76 |
| Harm to Rescuers From CPR (BLS 354:<br>ScopRev) .....                                      | S77 |
| Topics not Reviewed in 2020.....   | S77 |
| Disclosures.....   | S78 |
| References.....  | S80 |

This 2020 document is the fourth in a series of annual International Liaison Committee on Resuscitation (ILCOR) *International Consensus on Cardiopulmonary Resuscitation (CPR) and Emergency Cardiovascular Care (ECC) Science With Treatment Recommendations* (CoSTR) summary publications. This 2020 CoSTR for basic life support (BLS) includes new topics addressed by systematic reviews (SysRevs) performed within the past 12 months and prioritized by the BLS Task Force. It also includes updates of the BLS treatment recommendations published from 2010 through 2019,<sup>1-8</sup> as needed, based on additional evidence evaluations. As a result, this 2020 CoSTR for BLS is the most comprehensive update since 2010.

The 3 major types of evidence evaluation supporting this 2020 document are the SysRev, the scoping review (ScopRev), and the evidence update (EvUp).

The SysRev is a rigorous process, following strict methodology to answer a specific question; each of these ultimately resulted in generation of the task force consensus on science with treatment recommendations included in this document. The SysRevs were performed by a knowledge synthesis unit, an expert systematic reviewer, or the BLS Task Force, and many resulted in separate published SysRevs.

To begin the SysRev, the question to be answered was phrased in terms of the PICOST (population, intervention, comparator, outcome, study design, time frame) format. The methodology used to *identify* the evidence was based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses.<sup>9</sup> The approach used to *evaluate* the evidence was based on that proposed by the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) working group.<sup>10</sup> Using this approach, the task force rated as high, moderate, low, or very low the certainty/confidence in the estimates of effect of an intervention or assessment across a body of evidence (excluding animal studies) for each of the pre-defined outcomes. Randomized controlled trials (RCTs)

generally began the analysis as high-certainty evidence, and observational studies generally began the analysis as low-certainty evidence; examination of the evidence using the GRADE approach could result in downgrading or upgrading of the certainty of evidence. For additional information, refer to this supplement's "Evidence Evaluation Process and Management of Potential Conflicts of Interest."<sup>11</sup> Disclosure information for writing group members is listed in Appendix 1. Disclosure information for peer reviewers is listed in Appendix 2.

When a pre-2015 treatment recommendation was not updated, the language used differs from that used in the GRADE approach because GRADE was not used before 2015.<sup>12,13</sup>

Draft 2020 CoSTRs for BLS were posted on the ILCOR website<sup>14</sup> public comment between December 31, 2019, and February 16, 2020, with comments accepted through February 29, 2020. These new draft 2020 CoSTR statements for BLS received 45 694 views and 27 comments.

This summary statement contains the final wording of the CoSTR statements as approved by the ILCOR task forces and by the ILCOR member councils after review and consideration of comments posted online in response to the draft CoSTRs. Within this publication, each topic includes the PICOST as well as the CoSTR, an expanded "Justification and Evidence-to-Decision Framework Highlights" section, and a list of knowledge gaps requiring future research studies. An evidence-to-decision table is included for each CoSTR in Appendix A in the Supplemental Materials of this document.

The second major type of evidence evaluation performed to support this 2020 CoSTR for BLS is a ScopRev. ScopRevs are designed to identify the extent, range, and nature of evidence on a topic or a question, and they were performed by topic experts in consultation with the BLS Task Force. The task force analyzed the identified evidence and determined its value and implications for resuscitation practice or research. The rationale for the ScopRev, the summary of evidence, and the task force insights are all highlighted in the body of this publication. The most recent treatment recommendation is included. The task force notes whether the ScopRev identified substantive evidence that may result in a change in ILCOR treatment recommendations. If sufficient evidence was identified, the task force suggested consideration of a (future) SysRev to supply sufficient detail to support the development of an updated CoSTR. All ScopRevs are included in their entirety in Appendix B in the Supplemental Materials of this publication.

The third type of evidence evaluation supporting this 2020 CoSTR for BLS is an EvUp. EvUps are generally performed for topics previously reviewed by ILCOR to identify new studies published after the most recent ILCOR evidence evaluation, typically through use of search terms and methodologies from previous reviews. These EvUps were

performed by task force members, collaborating experts, or members of council writing groups. The EvUps are cited in the body of this document with a note about whether the evidence suggested the need to consider a SysRev; the existing ILCOR treatment recommendation was reiterated. In this document, no change in ILCOR treatment recommendations resulted from an EvUp; if substantial new evidence was identified, the task force recommended consideration of a SysRev. All EvUps are included in Appendix C in the Supplemental Materials of this publication.

The BLS Task Force considered the availability of new evidence as well as the evidence needed to create, confirm, or revise treatment recommendations. The chapter topics are organized in sections that approximate the order of the steps of resuscitation. For each reviewed topic, the method of review (SysRev, ScopRev, EvUp) is clearly labeled, with links to the relevant review documents in the Appendixes in the Supplemental Materials.

## TOPICS REVIEWED IN THIS 2020 BLS CoSTR

Note: As indicated above, the new BLS CoSTR evidence reviews were all completed in February 2020. As a result, this document does not address the topic of potential influence of coronavirus disease 2019 (COVID-19) on resuscitation practice. In the spring of 2020, an ILCOR writing group was assembled to identify and evaluate the published evidence regarding risks of aerosol generation and infection transmission during attempted resuscitation of adults, children, and infants. This group developed a consensus on science with treatment recommendations and task force insights. This statement is published as a separate document.<sup>15</sup> As new evidence emerges, the ILCOR task forces will review and update this statement, so the reader is referred to the ILCOR website<sup>14</sup> for the most up-to-date recommendations.

### Early Access and Cardiac Arrest Prevention, Including Emergency Medical Dispatch and Dispatcher-Assisted CPR (DA-CPR)

- Dispatch diagnosis of cardiac arrest (BLS 740: SysRev)
- Dispatcher instructions in CPR (2019 CoSTR BLS 359: SysRev)
- Dispatcher-assisted compression-only CPR versus conventional CPR (2017 CoSTR BLS 359: SysRev)

### Compression-Only CPR

- Lay rescuer chest compression-only versus standard CPR (2017 CoSTR BLS 547: SysRev)
- Emergency medical services (EMS) chest compression-only compared with conventional CPR (2017 CoSTR BLS 360: SysRev)
- In-hospital chest compression-only CPR versus conventional CPR (2017 CoSTR BLS 372: SysRev)

- Rescuer fatigue in chest compression-only CPR (BLS 349: ScopRev)

### CPR Sequence

- Firm surface for CPR (BLS 370: SysRev)
- Starting CPR (compressions-airway-breaths [C-A-B] versus airway-breaths-compressions [A-B-C]) (BLS 661: SysRev)
- CPR before call for help (BLS 1527: SysRev)
- Duration of CPR cycles (2 minutes versus other) (BLS 346: SysRev)
- Check for circulation during BLS (BLS 348: EvUp)

### Components of High-Quality CPR

- Hand position during compressions (BLS 357: SysRev)
- Chest compression rate, chest compression depth, and chest wall recoil (BLS 366, BLS 367, BLS 343: ScopRev)
- Compression-to-ventilation ratio (2017 CoSTR BLS 362: SysRev)
- Timing of rhythm check (BLS 345: SysRev)
- Feedback for CPR quality (BLS 361: SysRev)

### Alternative Techniques

- Alternative techniques (cough CPR, precordial thump, fist pacing) (BLS 374: SysRev)

### Defibrillation

- Public access automated external defibrillator (AED) programs (BLS 347: SysRev)
- Analysis of rhythm during chest compressions (BLS 373: SysRev)
- CPR before defibrillation (BLS 363: SysRev)
- Paddle size and placement for defibrillation (ALS-E-030A: ScopRev)

### Special Circumstances

- CPR during transport (BLS 1509: ScopRev)
- Removal of foreign-body airway obstruction (BLS 368: SysRev)
- Resuscitation care for suspected opioid-associated emergencies (BLS 811: SysRev)
- Drowning (BLS 856: SysRev)

### Potential Harm From CPR

- Harm from CPR to victims not in cardiac arrest (BLS 353: SysRev)
- Harm to rescuers from CPR (BLS 354: ScopRev)

## EARLY ACCESS AND CARDIAC ARREST PREVENTION, INCLUDING EMERGENCY MEDICAL DISPATCH AND DA-CPR

A variety of terms have been used to identify the person(s) at an emergency telephone call center who are charged with answering the call, interacting with the

caller, and assigning the needed care providers to the incident scene (traditionally called dispatchers). Terminology is similarly varied for the process the dispatcher uses to provide real-time CPR instructions to bystanders at the scene of an out-of-hospital cardiac arrest (OHCA). To remain consistent with the ILCOR evidence review, the term DA-CPR will be used to describe such coaching in this update, recognizing that other terms (eg, telecommunicator CPR and telephone CPR) could be substituted.

## Dispatch Diagnosis of Cardiac Arrest (BLS 740: SysRev)

### Rationale for Review

Accurate recognition of cardiac arrest by emergency medical dispatchers at the time of the emergency call is an important early step in cardiac arrest management, enabling initiation of DA-CPR and appropriate and timely emergency response. The overall accuracy of dispatchers in recognizing cardiac arrest is not well known. Furthermore, it is not known if there are specific call characteristics that affect the ability to recognize cardiac arrest.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA
- Intervention: Characteristics of the call process (these might include the specific words by the caller, language or idioms spoken by the caller and understood by the call taker, perceptions of the call receiver, emotional state of the caller, other caller characteristics, type of personnel receiving the call, background noises, etc)
- Comparator: Absence of identified characteristics of the call process
- Outcome: Any diagnostic test outcomes
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded.
- Time frame: All years and all languages were included, provided there was an English abstract. The literature search was updated November 28, 2019.
- International Prospective Register of Systematic Reviews (PROSPERO) registration: CRD42019140265

### Consensus on Science

A variety of algorithms and criteria (both commercial and locally developed) are used by dispatch centers to identify potential life-threatening events, such as cardiac arrest and triage emergency responders, to the scene appropriately. The dispatch centers reported great variability of overall accuracy of these algorithms and criteria for recognizing an OHCA in adults (Table 1).



**Table 1. Overall Diagnostic Performance of Dispatch Centers for Recognizing OHCA**

| Outcome                           | Certainty   | Studies | No. of Patients | Median (IQR)         |
|-----------------------------------|---|---------|-----------------|----------------------|
| Sensitivity                       | Very low (risk of bias, imprecision, inconsistency) | 46*     | 84 534†         | 0.79 (0.69–0.83)     |
| False-negative rate (undertriage) | Very low (risk of bias, imprecision, inconsistency) | 46*     | 84 534†         | 0.21 (0.17–0.32)     |
| Specificity                       | Very low (risk of bias, inconsistency)              | 12‡     | 789 004§        | 0.99 (0.93–1.00)     |
| False-positive rate (overtriage)  | Very low (risk of bias, inconsistency)              | 12‡     | 789 004§        | 0.01 (0.01–0.07)     |
| Negative predictive value         | Low (risk of bias, inconsistency)                   | 12‡     | 789 004§        | 1.00 (0.92–1.00)     |
| Positive predictive value         | Low (risk of bias, inconsistency)                   | 12‡     | 789 004§        | 0.76 (0.50–0.85)     |
| Positive likelihood ratio         | Low (risk of bias, inconsistency)                   | 12‡     | 789 004§        | 54.72 (11.28–152.22) |
| Negative likelihood ratio         | Low (risk of bias, inconsistency)                   | 12‡     | 789 004§        | 0.22 (0.19–0.24)     |

IQR indicates interquartile range; and OHCA, out-of-hospital cardiac arrest.

Sensitivity = proportion of confirmed cardiac arrest patients labeled as cardiac arrest by the dispatcher. False-negative rate = proportion of confirmed cardiac arrest patients who are not labeled as cardiac arrest by the dispatcher. Specificity = proportion of patients without confirmed cardiac arrest identified who are not labeled as cardiac arrest by dispatchers. False-positive rate = proportion of patients without cardiac arrest who are incorrectly labeled as cardiac arrest by the dispatcher. Negative predictive value = the proportion of patients labeled as not cardiac arrest by the dispatcher who are found not to have confirmed cardiac arrest. Positive predictive value = the proportion of patients labeled as cardiac arrest by dispatchers who are found to have confirmed cardiac arrest. Positive likelihood ratio = the likelihood of a patient with confirmed cardiac arrest to be labeled positive compared with a person without cardiac arrest (the higher the likelihood ratio, the better the test to rule in cardiac arrest). Negative likelihood ratio = the likelihood of a patient with confirmed cardiac arrest to be labeled negative compared with a person without cardiac arrest (the smaller the likelihood ratio, the better the test to rule out cardiac arrest).

\*References 16–61.

†Patients strictly with confirmed OHCA.

‡References 16,21,22,27,34,39,41,42,47,48,60,61.

§All patients inclusive of those without and with confirmed OHCA.

We compared subgroups of studies that used pre-determined or proprietary dispatching algorithms with those that used less structured criteria for diagnosis of cardiac arrest (dispatch algorithms versus criteria-based dispatch) and studies that reported different credential or training requirements for emergency dispatchers. No identifiable differences were noted in these subgroup analyses. Heterogeneity in studies and lack of adjusted analyses precluded meta-analysis for any subgroup.

### Treatment Recommendations

We recommend that dispatch centers implement a standardized algorithm and/or standardized criteria to immediately determine if a patient is in cardiac arrest at the time of emergency call (strong recommendation, very-low-certainty evidence).

We suggest that dispatch centers monitor and track diagnostic capability.

We suggest that dispatch centers look for ways to optimize sensitivity (minimize false negatives).

We recommend high-quality research that examines gaps in this area.

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-1](#). In making these new recommendations, we prioritized the desirable benefits (increase in potential lifesaving treatment) that would result from the immediate accurate identification of cardiac arrest by dispatchers. These benefits include the provision of DA-CPR and dispatching of appropriate EMS resources compared with the undesirable consequences of lack of early recognition of the event, such as delays to CPR and AED use.

We realize that efforts to minimize the frequency of undertriage (false-negative) may increase the frequency of overtriage (false-positive cases). Importantly, whether in cardiac arrest or not, the potential acuity of such patients still demands the need for immediate EMS assistance at the scene. In tiered response systems, if first-arriving EMS responders find a less emergent situation on arrival, the secondary advanced life support (ALS) response could be cancelled. In either event, the consequences of failing to recognize a genuine cardiac arrest in a timely manner is significant enough to justify some false-positive events. By comparison, the default position of most trauma systems is to have a high overtriage rate and a low undertriage rate because of similar concerns.

We were unable to make any recommendations on specific algorithms or criteria for identification of cardiac arrest because the variability across studies did not allow for direct comparisons or pooling of data. Furthermore, as the result of unexplained variability across studies, even among those using similar dispatch criteria, there was considerable variation in their diagnostic accuracy, which prevented pooling of data to find overall diagnostic accuracy measures for each of the algorithms. One factor that significantly influences the diagnostic accuracy is the prevalence of cardiac arrest in the reported population. In multiple studies, the denominator of total evaluated calls was different—some studies reporting cardiac arrests as a proportion of all emergency calls, others reporting cardiac arrests as a proportion of calls strictly among patients who were described as being unresponsive, and still other studies that (retrospectively) only included patients who were actually in cardiac arrest at the time of the call. Reporting the accuracy of identifying

a cardiac arrest as a proportion of all emergency calls can produce misleadingly favorable diagnostic statistics because, for the majority of such calls, it is obvious at the time that the patient is not in cardiac arrest.

Last, although studies that examined barriers to cardiac arrest identification were identified, these studies were not done in a manner that enabled calculation of the effect of these characteristics on OHCA diagnosis or on dispatcher performance.

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- Are there other potentially important criteria or ancillary tools in addition to standard dispatch algorithms that might improve dispatcher recognition of cardiac arrest? These might include use of a remote video link or pulse detection technologies via a caller's mobile telephone.
- What are the potential obstacles to dispatcher recognition of cardiac arrest (eg, language barriers, caller characteristics, patient characteristics)?
- Could the use of artificial intelligence improve recognition of cardiac arrest compared with emergency medical dispatcher recognition?
- What are the operational costs required for implementing and monitoring dispatcher recognition programs?
- What is the most accurate dispatch algorithm, and what are the optimal criteria for rapidly recognizing cardiac arrest?
- What is the relationship between dispatch algorithms and time to cardiac arrest recognition and time to initiation of DA-CPR?

### Dispatcher Instructions in CPR (2019 CoSTR BLS 359: SysRev)

DA-CPR has been reported in individual studies to significantly increase the rate of bystander CPR and survival from cardiac arrest. In 2019, we undertook a SysRev and meta-analysis to evaluate the impact of DA-CPR programs on key clinical outcomes after OHCA.<sup>62</sup> Consensus on science, values, preferences, and task force insights and knowledge gaps can be found in the *2019 International Consensus on CPR and ECC Science With Treatment Recommendations*.<sup>7,8</sup>

#### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with presumed OHCA
- Intervention: Patients/cases or EMS systems for which DA-CPR is offered
- Comparator: Studies with comparators in which either systems or specific cardiac arrest patients/cases were not offered DA-CPR were included

- Outcome: Critical—survival with favorable neurological function (at hospital discharge, 1 month, or 6 months), survival (to hospital discharge, 1 month, or 1 year), short-term survival (return of spontaneous circulation [ROSC], hospital admission), provision of bystander CPR; important—initial shockable rhythm, time to CPR
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) eligible for inclusion
- Time frame: All years and all languages included with the last search, performed July 1, 2018; ongoing or unpublished studies identified through a search of ClinicalTrials.gov online registry
- PROSPERO registration: CRD42018091427

### Treatment Recommendations

We recommend that emergency medical dispatch centers have systems in place to enable call handlers to provide CPR instructions for adult patients in cardiac arrest (strong recommendation, very-low-certainty evidence).

We recommend that emergency medical dispatchers provide CPR instructions (when deemed necessary) for adult patients in cardiac arrest (strong recommendation, very-low-certainty evidence).<sup>7,8</sup>

### DA-Assisted Compression-Only CPR Versus Conventional CPR (2017 CoSTR BLS 359: SysRev)

Emergency medical dispatchers typically are trained to provide telephone instructions for both compression-only CPR and conventional CPR with mouth-to-mouth ventilation. There is still some degree of controversy about whether it is sufficient for dispatchers to instruct callers to do only compression-only CPR for adult cardiac arrests or whether it is feasible to teach untrained lay rescuers over the phone how to perform mouth-to-mouth ventilation. This topic has been included in a SysRev and meta-analysis.<sup>63</sup> The task force CoSTR as well as values and preferences can be found in the *2017 International Consensus on CPR and ECC Science With Treatment Recommendations Summary*.<sup>5,6</sup> These note that the treatment recommendations prioritized the effective treatment for the most common causes of cardiac arrest (ie, cardiac causes). There remains uncertainty about the optimal approach when the cardiac arrest is caused by noncardiac causes, especially hypoxia.

#### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA
- Intervention: Dispatcher-assisted compression-only CPR
- Comparator: Dispatcher-assisted standard CPR

- Outcome: The primary outcome was favorable neurological outcomes, measured by cerebral performance or a modified Rankin scale. Secondary outcomes were survival, ROSC, and quality of life.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Study designs without a comparator group (ie, case series, cross-sectional studies), reviews, and pooled analyses were excluded.
- Time frame: Published studies in English were searched on January 15, 2016.
- PROSPERO registration: CRD42016047811

### **Treatment Recommendation**

We recommend that dispatchers provide chest compression-only CPR instructions to callers for adults with suspected OHCA (strong recommendation, low-certainty evidence).<sup>5,6</sup>

## **COMPRESSION-ONLY CPR**

One of the primary measures taken to improve survival after cardiac arrest is a focused effort to improve the quality of CPR. Although the impact of high-quality chest compressions has been studied extensively,<sup>64–69</sup> the role of ventilation and oxygenation in the initial management of cardiac arrest is less clear. Shortly after the publication of the *2015 International Consensus on CPR and ECC Science With Treatment Recommendations*,<sup>3,4</sup> a 23 711-patient RCT was published evaluating the effectiveness of continuous chest compressions (during which breaths were given without pausing chest compressions) in the EMS setting.<sup>70</sup> In parallel, developments of large national and regional registries are continually providing new insights into the epidemiology of cardiac arrest and effects of bystander CPR on outcomes.<sup>71</sup> These emerging publications generated an urgent need to review all available evidence on continuous compression strategies to provide an updated evidence evaluation that includes the latest science available. This topic has been included in a 2017 SysRev and meta-analysis.<sup>63</sup> The BLS Task Force CoSTR and its values and preferences can be found in the 2017 CoSTR summary.<sup>5,6</sup>

### **Lay Rescuer Chest Compression-Only Versus Standard CPR (2017 CoSTR BLS 547: SysRev)**

#### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with OHCA
- Intervention: Lay rescuer compression-only CPR
- Comparator: Lay rescuer standard CPR
- Outcome: The primary outcome was favorable neurological outcomes, measured by cerebral

performance or a modified Rankin scale. Secondary outcomes were survival, ROSC, and quality of life.

- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Study designs without a comparator group (ie, case series, cross-sectional studies), reviews, and pooled analyses were excluded.
- Time frame: Published studies in English were searched on January 15, 2016.
- PROSPERO registration: CRD42016047811

### **Treatment Recommendations**

We continue to recommend that bystanders perform chest compressions for all patients in cardiac arrest (good practice statement).

We suggest that bystanders who are trained, able, and willing to give rescue breaths and chest compressions do so for all adult patients in cardiac arrest (weak recommendation, very-low-certainty evidence).<sup>5,6</sup>

## **EMS Chest Compression-Only Compared With Conventional CPR (2017 CoSTR BLS 360: SysRev)**

#### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with OHCA treated by EMS
- Intervention: Compression-only CPR or minimally interrupted CPR (protocol for resuscitation based on commencing an initial 200 uninterrupted chest compressions and passive oxygen insufflation).
- Comparator: Standard CPR
- Outcome: The primary outcome was favorable neurological outcomes, measured by cerebral performance or a modified Rankin scale. Secondary outcomes were survival, ROSC, and quality of life.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Study designs without a comparator group (ie, case series, cross-sectional studies), reviews, and pooled analyses were excluded.
- Time frame: Published studies in English were searched on January 15, 2016.
- PROSPERO registration: CRD42016047811

### **Treatment Recommendations**

We recommend that EMS providers perform CPR with 30 compressions to 2 breaths (30:2 ratio) or continuous chest compressions with positive pressure ventilation delivered without pausing chest compressions until a tracheal tube or supraglottic device has been placed (strong recommendation, high-certainty evidence).

We suggest that, when EMS systems have adopted minimally interrupted cardiac resuscitation, this strategy

is a reasonable alternative to conventional CPR for witnessed shockable OHCA (weak recommendation, very-low-certainty evidence).<sup>5,6</sup>

## **In-Hospital Chest Compression–Only CPR Versus Conventional CPR (2017 CoSTR BLS 372: SysRev)**

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with in-hospital cardiac arrest (IHCA)
- Intervention: Compression-only CPR
- Comparator: Standard CPR
- Outcome: The primary outcome was favorable neurological outcomes, measured by cerebral performance or a modified Rankin scale. Secondary outcomes were survival, ROSC, and quality of life.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Study designs without a comparator group (ie, case series, cross-sectional studies), reviews, and pooled analyses were excluded.
- Time frame: Published studies in English were searched on January 15, 2016.
- PROSPERO registration: CRD42016047811

### **Treatment Recommendation**

Whenever tracheal intubation or a supraglottic airway is achieved during in-hospital CPR, we suggest that providers perform continuous compressions with positive pressure ventilation delivered without pausing chest compressions (weak recommendation, very-low-certainty evidence).<sup>5,6</sup>

## **Rescuer Fatigue in Chest Compression–Only CPR (BLS 349: ScopRev)**

### **Rationale for Review**

This topic was not a part of the 2017 SysRev<sup>63</sup> and CoSTR summary on continuous compressions versus standard CPR.<sup>5,6</sup> It was prioritized by the BLS Task Force for an updated evidence review, because this topic had not been reviewed by ILCOR since 2005.

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Rescuers performing CPR
- Intervention: Compression-only CPR
- Comparator: Standard CPR
- Outcome: Rescuer fatigue, CPR quality parameters (compression rate, compression depth, compression pauses, leaning or incomplete release, etc)

- Study design: RCTs, interrupted time series, controlled before-and-after studies, cohort studies, and manikin studies were eligible for inclusion.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 29, 2019.

### **Summary of Evidence**

This ScopRev is included in [Supplement Appendix B-1](#). Fifteen manikin studies evaluating fatigue at various compression-to-ventilation ratios were identified. These studies compared fatigue and its effects on CPR quality in volunteers performing continuous compressions and 30:2 or 15:2 CPR.<sup>72–86</sup> Evidence from these manikin studies comparing fatigue and effects on CPR quality suggest that continuous compressions are effective in the first 2 minutes with regard to depth and frequency, and there are indications that short periods of rest (pauses in compression) reduce rescuer fatigue and increase CPR quality.

### **Task Force Insights**

Continuous compression strategies increasingly have been advocated in an effort to increase overall bystander CPR rates. Evidence reviews evaluating the effect of continuous chest compressions versus standard CPR on critical outcomes, such as long-term survival, have been performed by the BLS Task Force in a separate published CoSTR.<sup>5,6</sup>

Although the BLS Task Force regards rescuer fatigue as an important barrier to high-quality bystander CPR, a higher value is placed on patient-centered outcomes.

### **Treatment Recommendation**

This treatment recommendation (below) is unchanged from 2015.<sup>3,4</sup>

We suggest pausing chest compressions every 2 minutes to assess the cardiac rhythm (weak recommendation, low-certainty evidence).

In making this recommendation, we placed a high priority on consistency with previous recommendations and the absence of contradictory evidence to prompt a change. We placed value on simplifying resuscitation logistics by coordinating rhythm and pulse checks with standard recommendations for rotating the provider performing chest compressions every 2 minutes.

## **CPR SEQUENCE**

### **Firm Surface for CPR (BLS 370: SysRev)**

#### **Rationale for Review**

This topic was prioritized for review by the BLS Task Force because it had not been updated since 2010.<sup>1,2</sup> Members of the task force reported variation in backboard use and the practice of moving a patient from



**Table 2. Firm Surface for CPR**

| Group                   | Certainty                  | Studies  | No. of Participants | Results   |
|-------------------------|----------------------------|--|---------------------|---|
| Mattress type           | Low (serious indirectness) | Four manikin RCTs* <sup>87–90</sup>                  | 33                  | No study identified a difference in chest compression depth between mattress types    |
| Floor compared with bed | Low (serious indirectness) | Two manikin RCTs (meta-analyzed) <sup>88,91</sup>    | 64                  | No effect on chest compression depth: mean difference 4.29 mm (95% CI, –0.70 to 9.27) |
|                         |                            | Two manikin RCTs* <sup>89,92</sup>                   | 34                  | Neither study identified a difference in chest compression depth between groups       |
| Backboard use           | Low (serious indirectness) | Six manikin RCTs (meta-analyzed) <sup>90,93–97</sup> | 221                 | Improved chest compression depth: mean difference 2.74 mm (95% CI, 1.19 to 4.28)      |
|                         |                            | One manikin RCT* <sup>98</sup>                       | 24                  | No difference in chest compression depth between groups                               |

\*Heterogeneity precluded meta-analysis

CPR indicates cardiopulmonary resuscitation; and RCT, randomized controlled trial.

the bed to the floor to improve the quality of CPR, so it was considered timely to review the published evidence.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults or children in cardiac arrest (OHCA and IHCA) on a bed
- Intervention: CPR on a hard surface (eg, backboard, floor, deflatable or specialist mattress)
- Comparator: CPR on a regular mattress
- Outcome: Survival, survival with a favorable neurologic outcome, ROSC, CPR quality
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Randomized manikin/simulation/cadaver studies were only included if insufficient human studies were identified. Unpublished studies (eg, conference abstracts, trial protocols), nonrandomized manikin/simulation/cadaver studies, animal studies, experimental/laboratory models, mathematical models, narrative reviews, and editorials and opinions with no primary data were excluded.
- Time frame: January 1, 2009, to September 16, 2019
- PROSPERO registration: CRD42019154791

### Consensus on Science

The identified science has been grouped under the following subheadings: mattress type, floor compared with bed, and backboard in Table 2.

### Treatment Recommendations

We suggest performing manual chest compressions on a firm surface when possible (weak recommendation, very-low-certainty evidence).

During IHCA, we suggest that, when a bed has a CPR mode that increases mattress stiffness, it should be activated (weak recommendation, very-low-certainty evidence).

During IHCA, we suggest against moving a patient from a bed to the floor to improve chest compression depth (weak recommendation, very-low-certainty evidence).

The confidence in effect estimates is so low that the task force was unable to make a recommendation about the use of a backboard strategy.

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-2](#).

The context for this question was that, when manual chest compressions are performed on a mattress, the compression force is dissipated through both chest compression and compression of the mattress under the patient. Manikin models indicate that the amount of mattress compression ranges from 12% to 57% of total compression depth, with softer mattresses compressed the most.<sup>87,90,99,100</sup> This mattress compression can lead to reduced spinal-sternal displacement and a reduction in effective chest compression depth.

Effective compression depths can be achieved even on a soft surface, providing the CPR provider increases overall compression depth to compensate for mattress compression.<sup>90,97,101–105</sup> CPR feedback devices that account for mattress compression (eg, the use of dual accelerometers or increasing compression depth targets) can help CPR providers ensure adequate compression depth when CPR is performed on a mattress.<sup>95,99,101,103,105,106</sup>

In making these recommendations, the task force highlights the importance of high-quality chest compressions for optimizing outcomes from cardiac arrest.

The task force noted that there were no clinical studies reporting on the critical outcomes of survival and favorable neurological outcome or important outcome of chest compression quality.

The weak recommendations are based on extrapolation from manikin studies, typically undertaken on a mattress placed on a hospital bed, for which manual CPR was performed by a trained healthcare professional. The hospital beds involved in the studies typically had rigid bases. The task force noted that, although this configuration is common in hospitals in many developed countries, it may not be applicable to all hospitals

or the out-of-hospital setting. The absence of studies simulating out-of-hospital settings (where beds may be softer) and in which the CPR provider may be a single untrained rescuer led the task force to focus recommendations on the in-hospital setting.

The task force supported performing manual chest compressions on a firm surface when possible because this reduces the risks of shallow compressions attributable to performing CPR on a soft surface. On the other hand, moving a patient onto a hard surface can be a major barrier to CPR, and the importance of performing CPR on a firm surface needs to be weighed against the likelihood of significant delay in providing CPR. In the setting of DA-CPR, in particular, logistical aspects of moving patients from bed to floor can impede if not thwart the performance of CPR.

The task force considered that, when a mattress with CPR function was available, activating a CPR function on a mattress, although unlikely to substantially improve compression depth, posed a low risk of harm to rescuers and patients, leading to a weak recommendation of support.

In considering whether to transfer a patient from a hospital bed to the floor to improve compression depth, the task force considered that the risks of harm (eg, interruption in CPR, risk of losing vascular access for intravenous drug delivery, and more confined space) to the patient and resuscitation team outweighed any small improvement in chest compression depth, leading to a weak recommendation against routine use of this practice.

The task force was unable to make a recommendation for the use of a CPR backboard during IHCA. Within the limitations of manikin studies, the available evidence indicates a marginal benefit to chest compression depth from use of a backboard. For example, placing a firm surface (eg, a backboard) between the patient and a soft surface may merely transfer the same force from CPR to the underlying softness and not obviate potential concern over chest compression depth. No studies specifically evaluated backboard deployment or any impact this has on interruptions to chest compressions and/or displacement of tubes and catheters during insertion. For healthcare systems that have already incorporated backboards into routine use during IHCA, the evidence was considered insufficient to suggest against their continued use. For healthcare systems that have not introduced backboards, the limited improvement in compression depth and uncertainty about harms seemed insufficient to justify the costs of purchasing backboards and training staff in their use. When backboards are deployed, users should be aware that mattress stiffness, backboard size (larger is better), and orientation (longitudinal is better) influence their effectiveness.<sup>107–111</sup>

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- Studies reporting clinical outcomes
- Studies examining the logistical aspects of backboard deployment or moving a patient from a bed to the floor
- Studies relevant to OHCA
- Studies in both high- and low-resource settings, in which hospital bed or prehospital stretcher configurations may vary

### Starting CPR (C-A-B Compared With A-B-C) (BLS 661: SysRev)

Although, internationally, most adult BLS guidelines recommend commencing chest compressions before rescue breaths, debate about this sequence continues. In addition, there is variability in the sequences used for pediatric resuscitation and for aquatic rescue, with different approaches in various jurisdictions.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA
- Intervention: Commencing CPR beginning with compressions first (30:2)
- Comparator: CPR beginning with ventilation first (2:30)
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; and ROSC
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion.
- Exclusion criteria: Unpublished studies (eg, conference abstracts, trial protocols) and animal studies were excluded. Studies of dispatcher- or telephone-assisted CPR were excluded.
- Time frame: All languages were included as long as there was an English abstract. The literature search was updated in September 2019.

### Consensus on Science

This current SysRev did not identify any additional human or manikin studies published since the 2015 CoSTR SysRev.<sup>3,4</sup> The published evidence remains limited to 4 manikin studies: 1 randomized study<sup>112</sup> focused on adult resuscitation, 1 randomized study focused on pediatric resuscitation,<sup>113</sup> and 2 observational studies focused on adult resuscitation.<sup>114,115</sup> The results from these studies are summarized in Table 3.

The overall certainty of evidence was rated as very low for all outcomes primarily because of a very serious risk of bias and indirectness. The individual observational studies were all at a critical risk of bias because of confounding, and the RCTs were all at critical risk of bias

**Table 3. Starting CPR**

| Outcome  | Certainty | Studies   | No. of Patients  | Results  |
|--|-----------|---|--|--|
| Time to commencement of chest compressions   | Very low  | 1 RCT (manikin): Lubrano 2012 <sup>113</sup>  | 155 two-person teams   | Statistically significant 24-s difference ( $P<0.05$ ) in favor of C-A-B   |
|  |           | 2 observational (manikin): Kobayashi 2008, <sup>114</sup> Sekiguchi 2013 <sup>115</sup> | 40 individual rescuers <sup>115</sup> and 33 six-person teams <sup>114</sup> | The observational studies found statistically significant decreases of 20 s ( $P<0.001$ ) <sup>115</sup> and 26 s ( $P<0.001$ ) <sup>114</sup> in favor of C-A-B.  |
| Time to commencement of rescue breaths   | Very low  | 2 RCTs (manikin): Marsch 2013, <sup>112</sup> Lubrano 2012 <sup>113</sup>               | 210 two-person teams   | In a respiratory arrest scenario, there was a 4-second difference ( $P<0.05$ ) in favor of C-A-B <sup>113</sup> ; in a cardiac arrest scenario, A-B-C decreased the time to commencement of rescue breaths by 6 s ( $P<0.05$ ), and C-A-B decreased time to commencement of rescue breaths by 5 s ( $P<0.05$ ). <sup>112</sup> |
| Time to completion of first CPR cycle (30 chest compressions and 2 rescue breaths) | Very low  | 1 RCT (manikin): Marsch 2013 <sup>112</sup>   | 55 two-person teams  | C-A-B decreased time to completion of first CPR cycle by 15 s ( $P<0.001$ ).   |

A-B-C indicates airway-breathing-compression; C-A-B, compression-airway-breathing; CPR, cardiopulmonary resuscitation; and RCT, randomized controlled trial.

because of lack of blinding. Because of this and a high degree of heterogeneity, no meta-analyses could be performed. Individual studies are difficult to interpret.

### Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.<sup>3,4</sup>

We suggest commencing CPR with compressions rather than ventilation in adults with cardiac arrest (weak recommendation, very-low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-3](#). No change was made to this adult treatment recommendation. For all outcomes, starting CPR with compressions resulted in faster times to key elements of resuscitation (rescue breaths, chest compressions, completion of first CPR cycle) across the 4 papers reviewed, with the exception of simulated pediatric resuscitation, for which starting with compressions delayed time to commencement of rescue breaths in cardiac arrest by 6 seconds. This difference was statistically significant but reflects a delay that is not considered clinically significant.<sup>113</sup> This delay in commencing rescue breaths may be acceptable given the decreased time to other elements of resuscitation; however, the certainty of the evidence is very low, and all studies reviewed were manikin studies. There is no clinical evidence to guide whether to initiate compressions before ventilation in adult cardiac arrest. There should also be consideration given to the impact of simplification of training requirements by using a single approach compared with separate approaches for adults and children.

### Knowledge Gaps

- No human studies evaluating this question in any setting were identified.
- Important uncertainties regarding timing and delays in initiation of the CPR components (chest compressions, opening airway, and rescue breaths)

remain and may not be readily extrapolated from manikin studies.

### CPR Before Call for Help (BLS 1527: SysRev)

This question was suggested by the resuscitation community during the public commentary process. The question of the optimal sequence for calling for help and starting CPR frequently arises during CPR training courses, and a SysRev of the literature to guide recommendations was therefore prioritized by the BLS Task Force. Searching for new science from the era of increased availability of communication devices and hands-free alternatives for lone rescuers was also considered important in this evidence review.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA
- Intervention: CPR before call for help; immediate CPR by a lone bystander performed for a short time interval (ie, 1 minute) before alerting EMS dispatch center with a mobile phone
- Comparator: An immediate call for help to the EMS dispatch center by a lone bystander with a mobile phone
- Outcome: Survival with favorable neurological outcome until and beyond hospital discharge or 30 days; survival until and beyond hospital discharge or 30 days; ROSC
- Study design: We included RCTs, nonrandomized studies, and case series with at least 5 cases. We considered papers in all languages provided there was an English language abstract available for review. We excluded unpublished studies, conference abstracts, manikin or simulation studies, narrative reviews, editorials or opinions with no primary data, animal studies and experimental/laboratory models.
- Time frame: All years and all languages were included as long as there was an English abstract;

unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 2019.

### Consensus on Science

For the critical outcome of survival with favorable neurological outcome, we identified only a single observational study.<sup>116</sup> The overall certainty of evidence was rated as very low because of a very serious risk of bias. With the identification of only 1 study, no meta-analyses were performed.

For the critical outcome of survival with favorable neurological outcome, we identified very-low-certainty evidence (downgraded for very serious risk of bias) from 1 cohort study including 17 461 OHCA occurrences from Japan (2005–2012), which showed no benefit from a “CPR-first” strategy (cohort of 5 446 OHCA patients) compared with a “call-first” strategy (cohort of 1 820 OHCA patients).<sup>116</sup>

Adjusted analyses were performed on various subgroups and suggested significant improvements in survival with a favorable neurological outcome with a “CPR-first” strategy compared with a “call-first” strategy for noncardiac etiology OHCA (adjusted odds ratio [AOR], 2.01; 95% CI, 1.39–2.98); under 65 years of age (AOR, 1.38; 95% CI, 1.09–1.76); under 20 years of age (AOR, 3.74; 95% CI, 1.46–9.61); and both under 65 years of age and noncardiac etiology together (AOR, 4.31; 95% CI, 2.38–8.48).<sup>116</sup>

### Treatment Recommendation

We recommend that a lone bystander with a mobile phone should dial EMS, activate the speaker or other hands-free option on the mobile phone, and immediately begin CPR with dispatcher assistance, if required (strong recommendation, very-low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-4](#). This SysRev was based on a new PICOST question suggested during public commenting and, therefore, includes a new treatment recommendation. The included paper analyzed only 17 461 OHCA occurrences from 925 288 recorded in the Japan national registry in the period from 2005 to 2012. Analysis was limited to cases in which lay rescuers witnessed the OHCA and spontaneously performed CPR (without the need for dispatcher assistance), and the groups compared were different with respect to age, gender, initial rhythm, bystander CPR characteristics, and EMS intervals. Although some factors were adjusted for in subgroup analyses, there is significant risk of confounding. Despite very-low-certainty evidence, there was consensus among the BLS Task Force to make a strong recommendation.

There were many exclusion criteria: unwitnessed, prehospital involvement of physician or unknown,

EMS-witnessed OHCA, bystander-witnessed cases with missing data on time to intervention, no bystander CPR, DA-CPR, no intervention in 0 to 1 minutes, no CPR at all within 4 minutes, and etiology (cardiac or noncardiac) unknown.

There were some benefits noted in subgroup analyses, but these groups were not specified a priori. We cannot expect a bystander to reliably determine whether a cardiac arrest is of cardiac or noncardiac etiology. The results are not generalizable to all OHCA cases because they refer specifically to bystander-witnessed cases in which the bystander spontaneously initiates CPR after only a short delay.

The timings of interventions were determined after the event by EMS personnel who interviewed the bystanders. These timings may be imprecise or inaccurate in an undetermined number of cases.

The wide availability of mobile phones may reduce the likelihood that a lone bystander would have to leave a victim to phone EMS. Pragmatically, it is now often possible to perform both actions simultaneously, and the focus should be on empowering people to recognize OHCA and initiate both an EMS call and CPR as soon as possible. In the absence of any evidence to the contrary, this would apply to both witnessed and unwitnessed OHCA, except in circumstances when there are appropriate reasons not to start CPR. When more than 1 bystander is at the scene, calling EMS and initiating CPR can be performed simultaneously. For the single rescuer, a call-first strategy ensures that EMS providers are dispatched as soon as possible, bringing additional assets (including a defibrillator) that might otherwise be delayed by a later call. Telecommunicator prompting may promote the initiation of bystander CPR that might not otherwise occur or may support better quality CPR (eg, instructing the caller to press hard and count aloud, helping to pace the compression rate).

In the situation when a lone rescuer would have to leave a victim alone to dial EMS, the priority is prompt activation of EMS before subsequently returning to the victim to initiate CPR as soon as possible.

### Knowledge Gaps

There is no evidence comparing an immediate call to EMS for help with a call after 1 minute of CPR in the specific circumstance of a lone bystander with a mobile phone. There is also no evidence about how long it takes to call EMS after a witnessed cardiac arrest. The delay between a witnessed arrest and a call to EMS may be substantial.

### Duration of CPR Cycles (2 Minutes Versus Other) (BLS 346: SysRev)

#### Rationale for Review

The recommendations for CPR cycle duration have changed with time, but these changes have never



**Table 4. 1-Minute CPR Duration Compared With 3-Minute Duration for Postshock VF/pVT**

| Outcome  | Certainty                       | Studies                      | No. of Patients | Results   |
|--|---------------------------------|------------------------------|-----------------|---|
| Hospital discharge with favorable neurological outcome | Low (risk of bias, imprecision) | RCT: Wik 2003 <sup>117</sup> | 200             | No difference:<br>Relative risk 1.68 (95% CI, 0.85–3.32), 78 more patients/1000 (–17 to 266)  |
| Survival to hospital discharge                         | Low (risk of bias, imprecision) | RCT: Wik 2003 <sup>117</sup> | 200             | No difference:<br>Relative risk 1.52 (95% CI, 0.83–2.77), 76 more patients/1000 (–25 to 258)  |
| ROSC   | Low (risk of bias, imprecision) | RCT: Wik 2003 <sup>117</sup> | 200             | No difference:<br>Relative risk 1.22 (95% CI, 0.92–1.50), 101 more patients/1000 (–37 to 229) |

CPR indicates cardiopulmonary resuscitation; RCT, randomized controlled trial; ROSC, return of spontaneous circulation; pVT, pulseless ventricular tachycardia; and VF, ventricular fibrillation.

Both relative and absolute risks are written as mean values (95% CIs).

been based on high-certainty evidence that any specific interval or CPR cycle duration was superior in terms of patient survival. Because the topic has not been reviewed since 2015,<sup>3,4</sup> when no direct evidence was identified, the following PICOST question was prioritized for evidence review.

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with cardiac arrest
- Intervention: Pausing chest compressions at another interval
- Comparator: Pausing chest compressions every 2 minutes to assess the cardiac rhythm
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to September 2019.

### **Consensus on Science**

Data were derived from 2 RCTs<sup>117,118</sup> for which the principal focus was on the period of time allotted for CPR *before* the first rhythm analysis. Assessment of the duration (in minutes) of uninterrupted CPR

between subsequent rhythm checks and outcome were not formally reported analyses in either study. The published data in these 2 studies enabled an ad hoc analysis by ILCOR evidence evaluation experts that indirectly addressed this question. Outcomes were not adjusted for possible confounders.

### **1-Minute CPR Duration Compared With 3-Minute Duration for Postshock Ventricular Fibrillation (VF)/Pulseless Ventricular Tachycardia (pVT)**

In the 2003 study including 1-minute and 3-minute durations of uninterrupted CPR between rhythm checks,<sup>117</sup> the control group included patients who received immediate defibrillation (up to 3 stacked shocks) for VF/VT followed by 1 minute of CPR for patients in refractory VF/VT at the next rhythm check and 3 minutes of CPR for those patients who exhibited nonshockable rhythms after 1 to 3 shocks. The intervention group included patients who received 3 minutes of CPR before the first defibrillation attempt (up to 3 stacked shocks) for VF/VT followed by CPR for 3 minutes regardless of postshock rhythm. Of note, none of the patients received 2-minute periods of CPR. This RCT showed no benefit from the intervention compared with the control CPR duration between rhythms checks for all of the outcomes listed (Table 4).

### **1-Minute CPR Duration Compared With 2-Minute CPR Duration**

In the 1 study that included 1-minute and 2-minute durations of uninterrupted CPR between rhythm checks,<sup>118</sup> the 2-minute group included patients who were enrolled in the RCT after implementation of new guidelines introducing single shocks, 30:2 CPR, and

**Table 5. 1-Minute CPR Duration Compared With 2-Minute CPR Duration**

| Outcome                        | Certainty  | Studies                        | No. of Patients | Results  |
|--------------------------------|--|--------------------------------|-----------------|--|
| Survival to hospital discharge | Very low (serious risk of bias, indirectness, imprecision) | RCT: Baker 2008 <sup>118</sup> | 202             | No difference:<br>Relative risk 0.49 (95% CI, 0.23–1.06), 92 fewer patients/1000 (–139 to 11)  |
| ROSC                           | Very low (serious risk of bias, indirectness, imprecision) | RCT: Baker 2008 <sup>118</sup> | 202             | No difference:<br>Relative risk 0.95 (95% CI, 0.73–1.24), 27 fewer patients/1000 (–144 to 128) |

CPR indicates cardiopulmonary resuscitation; RCT, randomized controlled trial; and ROSC, return of spontaneous circulation.

Both relative and absolute risks are written as mean values (95% CIs).

2-minute CPR cycles between rhythm checks. The 1-minute group included patients who were enrolled in the RCT before implementation of new guidelines and were therefore treated with stacked shocks (up to 3 in refractory VF/VT), 15:2 CPR, and 1-minute CPR cycles between rhythm checks. No clear benefit from either the 1- or 2-minute duration between rhythm checks was observed (Table 5).

### **Treatment Recommendation**

This treatment recommendation (below) is unchanged from 2015.<sup>3,4</sup>

We suggest pausing chest compressions every 2 minutes to assess the cardiac rhythm (weak recommendation, low-certainty evidence).

### **Justification and Evidence-to-Decision Framework Highlights**

The evidence-to-decision table is included in [Supplement Appendix A-5](#). No change was made to this treatment recommendation. This topic was prioritized for review by the BLS Task Force because it had not been updated since the 2015 CoSTR. Although the current review identified 2 older studies that included comparisons of groups with different CPR durations between rhythm checks, each had significant limitations. Both studies were designed to address the question of CPR first compared with defibrillation first. As a result, the certainty of evidence derived from these studies is low, and recommendations regarding optimal duration of CPR before a scheduled rhythm analysis are seriously confounded.

In making the suggestion to pause chest compressions every 2 minutes to assess cardiac rhythm, we placed a high value on being consistent with previous recommendations, and noting the only limited indirect evidence identified in this review. The BLS Task Force acknowledges that every change in guidelines comes with a significant risk and cost as CPR educators and providers are asked to change current practice and implement new treatment strategies for complex and high-stress medical emergencies.

### **Knowledge Gaps**

- Does the optimal CPR duration (ie, interval between rhythm analyses) differ for patients with different initial or postshock cardiac rhythms?
- Does the duration between collapse and EMS arrival affect the optimal CPR duration/interval between rhythm checks?
- Do different intervals between rhythm checks interfere with the overriding goal of minimizing interruptions in chest compressions?
- What is the relationship between rescuer fatigue, chest compression quality, and the optimal CPR duration/interval between rhythm checks?

## **Check for Circulation During BLS (BLS 348: EvUp)**

An EvUp (see [Supplement Appendix C-1](#)) identified no evidence to justify a SysRev or a change in the 2015 treatment recommendation.<sup>3,4</sup>

Future reviews could focus on combination/alternative techniques used to confirm presence of circulation: plethysmography, arterial pressure monitoring, end-tidal carbon dioxide (ETCO<sub>2</sub>), near infrared spectroscopy, ultrasound, and more.

### **Treatment Recommendation**

Outside of the ALS environment, where invasive monitoring is available, there are insufficient data about the value of a pulse check while performing CPR. We therefore do not make a treatment recommendation regarding the value of a pulse check.<sup>3,4</sup>

## **COMPONENTS OF HIGH-QUALITY CPR**

### **Hand Position During Compressions (BLS 357: SysRev)**

#### **Rationale for Review**

The recommendations for hand position during compressions have changed with time, but these changes have been based on only low- or very-low-certainty evidence, with no data demonstrating that a specific hand position was optimal in terms of patient survival. The topic has not been reviewed since 2015,<sup>3,4</sup> when no direct evidence was identified, so the following PICOST question was prioritized for evidence review.

#### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with cardiac arrest
- Intervention: Delivery of chest compressions on the lower half of the sternum
- Comparator: Any other location for chest compressions
- Outcome: Any clinical outcome. Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome. Physiological outcomes, such as blood pressure, coronary perfusion pressure, or ETCO<sub>2</sub>, also were considered important.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded.
- Time frame: SysRev search strategy: All years and all languages were included as long as there was an English abstract.

## Consensus on Science

There were no studies reporting the critical outcomes of favorable neurological outcome, survival, or the important outcome of ROSC. For the important outcome of physiological end points, we identified 3 very-low-certainty studies (downgraded for bias, indirectness, and imprecision).<sup>119–121</sup> One crossover study in 17 adults with prolonged resuscitation from nontraumatic cardiac arrest observed improved peak arterial pressure during compression systole ( $114 \pm 51$  mmHg compared with  $95 \pm 42$  mmHg) and  $\text{ETCO}_2$  ( $11.0 \pm 6.7$  mmHg compared with  $9.6 \pm 6.9$  mmHg) when compressions were performed over the lower third of the sternum compared with the center of the chest, but arterial pressure during compression recoil, peak right atrial pressure, and coronary perfusion pressure did not differ.<sup>120</sup> A second crossover study in 30 adults with cardiac arrest observed no difference in  $\text{ETCO}_2$  values resulting from changes in hand placement.<sup>121</sup> A third crossover study in 10 children observed higher peak systolic pressure and higher mean arterial pressure when compressions were performed on the lower third of the sternum compared with the middle of the sternum.<sup>119</sup>

## Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.<sup>3,4</sup>

We suggest performing chest compressions on the lower half of the sternum on adults in cardiac arrest (weak recommendation, very-low-certainty evidence).

## Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-6](#). In making this recommendation, we placed high value on consistency with current treatment recommendations in the absence of compelling clinical data suggesting the need to change the recommended hand placement for performing chest compressions.

## Knowledge Gaps

- We did not identify any studies that evaluated the effect of any specific hand position on short- or long-term survival after cardiac arrest; only physiological surrogate outcomes have been reported.
- Imaging studies suggest that there might be important differences in anatomy depending on age, gender, body mass index, presence or absence of chronic heart conditions, and more.
- Important gaps remain in evaluating how to identify optimal hand placement and/or compression point when using physiological feedback during CPR.

## Chest Compression Rate, Chest Compression Depth, and Chest Wall Recoil (BLS 366, BLS 367, BLS 343: ScopRev)

### Rationale for Review

The BLS Task Force requested a ScopRev related to chest compression rate, chest compression depth, and chest wall recoil to identify any recent published evidence that provided more information on these chest compression components as discrete entities and to assess whether studies have reported interactions among these chest compression components. Therefore, a ScopRev was undertaken to understand whether the science to date has focused on single chest compression components or interactions among chest compression components and identify the evidence related to the chest compression components to determine whether the body of evidence published since the 2015 CoSTR for BLS<sup>3,4</sup> indicates the need for a full SysRev of the evidence related to chest compression components.<sup>122</sup>

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with cardiac arrest
- Intervention/Comparator: (1)  $\geq 2$  chest compression depths measured in millimeters, centimeters, or inches or (2)  $\geq 2$  chest compression rates measured in compressions per minute or (3)  $\geq 2$  measures of chest wall recoil or (4)  $\geq 2$  measures of leaning or leaning compared with no leaning
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC or survival to a defined time point and physiological measures (eg, blood pressure and  $\text{ETCO}_2$ ) were ranked as important outcomes.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to June 2019.

### Summary of Evidence

In addition to the 14 studies identified in the 2015 CoSTR for BLS,<sup>3,4</sup> an additional 8 studies<sup>123–129a</sup> were identified, so a total of 22 studies were included in this ScopRev, which has been published in full.<sup>122</sup> Five observational studies examined both chest compression rate and chest compression depth.<sup>127,128,129a,130,131</sup> One RCT,<sup>124</sup> 1 crossover trial,<sup>132</sup> and 6 observational studies<sup>125,129,133–136</sup> examined chest compression rate only. One RCT<sup>137</sup> and 6 observational studies examined chest compression depth only,<sup>67,138–142</sup> and 2 observational studies examined

chest wall recoil.<sup>123,126</sup> No studies were identified that examined different measures of leaning. This ScopRev (see [Supplement Appendix B-2](#)) does highlight significant gaps in the research evidence related to chest compression components, namely a lack of high-level evidence, a paucity of studies of IHCA, and a failure to account for the possibility of interactions between chest compression components.

### **Task Force Insights**

In the evidence identified in this ScopRev, most studies focused on a single chest compression component, whereas several studies suggested the presence of confounding interactions that prompt caution when evaluating any chest compression component in isolation. Most studies identified in this review focused on OHCA, highlighting a major gap in research involving IHCA.

This ScopRev did not identify sufficient new evidence that would justify conducting new SysRevs or reconsideration of current resuscitation guidelines.

### **Treatment Recommendation**

These treatment recommendations (below) are unchanged from 2015.<sup>3,4</sup>

We recommend a manual chest compression rate of 100 to 120/min (strong recommendation, very-low-certainty evidence).

We recommend a chest compression depth of approximately 5 cm (2 in) (strong recommendation, low-certainty evidence) while avoiding excessive chest compression depths (greater than 6 cm [greater than 2.4 in] in an average adult) during manual CPR (weak recommendation, low-certainty evidence).

We suggest that rescuers performing manual CPR avoid leaning on the chest between compressions to allow full chest wall recoil (weak recommendation, very-low-certainty evidence).

## **Compression-to-Ventilation Ratio (2017 CoSTR BLS 362: SysRev)**

### **Rationale for Review**

The first ILCOR review to be performed after the 2015 CoSTR was a large SysRev<sup>63</sup> of compression strategies across different settings and populations. One of these comparisons addressed the optimal compression-to-ventilation ratio. Task force values and preferences can be found in the 2017 CoSTR summary.<sup>5,6</sup>

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with OHCA
- Intervention: Any compression-to-ventilation ratio other than 30:2
- Comparator: Compression-to-ventilation ratio of 30:2

- Outcome: The primary outcome was favorable neurological outcomes, measured by cerebral performance or a modified Rankin scale. Secondary outcomes were survival, ROSC, and quality of life.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Study designs without a comparator group (ie, case series, cross-sectional studies), reviews, and pooled analyses were excluded.
- Time frame: Published studies in English were searched on January 15, 2016.
- PROSPERO registration: CRD42016047811

### **Treatment Recommendation**

We suggest a compression-to-ventilation ratio of 30:2 compared with any other ratio in patients with cardiac arrest (weak recommendation, very-low-quality evidence).<sup>5,6</sup>

## **Timing of Rhythm Check (BLS 345: SysRev)**

### **Rationale for Review**

Adverse outcomes after cardiac arrest have been associated with frequent or prolonged interruptions in chest compressions. Because rhythm checks during resuscitation are frequent causes of pauses in compressions, this SysRev was undertaken to assess the evidence available to identify the optimal timing for rhythm checks.

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults with presumed cardiac arrest in in-hospital or out-of-hospital settings receiving a defibrillation attempt during CPR
- Intervention: Checking the cardiac rhythm immediately after defibrillation
- Comparator: Immediate resumption of chest compressions with delayed check of the cardiac rhythm
- Outcome: Critical—survival with good neurological function (ie, at hospital discharge, 1 month, 6 months, 1 year), survival (ie, hospital discharge, 1 month, 6 months, 1 year); important—short-term survival (ROSC, hospital admission), rates of recurrence of fibrillation/refibrillation, CPR quality parameters (ie, compression fraction).
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Animal/laboratory studies, mathematical models, simulation and manikin studies, algorithm studies for rhythm analysis recognition with no outcome data, unpublished studies (eg, conference abstracts, trial protocols), and reviews were excluded.



**Table 6.** Timing of Rhythm Check

| Outcome  | Certainty   | Studies   | No. of Patients | Results   |
|--|---|---|-----------------|---|
| Hospital discharge with favorable neurological outcome | Low (risk of bias, indirectness)<br>Very low (risk of bias, indirectness, imprecision)    | 1 RCT <sup>145</sup><br>3 observational <sup>146–148</sup>      | 415<br>763      | No difference:<br>Relative risk 0.90 (95% CI, 0.70–1.15), 40 fewer patients/1000 (–119 to 60)<br>Lower survival in immediate rhythm check:<br>Relative risk 0.62 (95% CI, 0.51–0.75), 174 fewer patients/1000 (–224 to –13) |
| Survival to hospital discharge                         | Low (serious risk of bias, indirectness)<br>Very low (serious risk of bias, indirectness) | 2 RCTs <sup>143,145</sup><br>3 observational <sup>146–148</sup> | 1 260<br>3 094  | No difference:<br>Relative risk 0.89 (95% CI, 0.72–1.10), 24 fewer patients/1000 (–63 to 23)<br>Lower survival in immediate rhythm check:<br>Relative risk 0.55 (95% CI, 0.45–0.67), 76 fewer patients/1000 (–93 to –56)    |
| Survival to hospital admission                         | Low (serious risk of bias, indirectness)  | 2 RCTs <sup>143,145</sup>                                       | 1 260           | No difference:<br>Relative risk 1.02 (95% CI, 0.91–1.14), 9 more patients/1000 (–43 to 69)  |
| ROSC   | Very low (serious risk of bias, indirectness)   | 2 observational <sup>147,148</sup>                              | 2 969           | Lower survival in immediate rhythm check:<br>Relative risk 0.69 (95% CI, 0.61–0.78), 111 fewer patients/1000 (–139 to –80)  |
| VF recurrence  | Very low (serious risk of bias, indirectness, imprecision)                                | 2 RCTs <sup>144,145</sup>                                       | 551             | No difference:<br>Relative risk 1.08 (95% CI, 0.95–1.22), 47 more patients/1000 (–13 to 5)  |

RCT indicates randomized controlled trial; ROSC, return of spontaneous circulation; and VF, ventricular fibrillation.  
Both relative and absolute risks are written as mean values (95% CIs).

- Time frame: All years and all languages were included provided there was an English abstract. The literature search was updated to November 2, 2019.

### Consensus on Science

Three RCTs<sup>143–145</sup> and 3 observational studies<sup>146–148</sup> were identified comparing immediate rhythm checks to immediate resumption of chest compressions. Outcomes assessed varied from hospital discharge with favorable neurological outcome to recurrence of VF. The meta-analysis of the RCTs did not demonstrate any differences between immediate rhythm analysis and immediate compressions, but unadjusted analysis of observational data suggested that immediate compressions were associated with better outcomes (Table 6).

### Treatment Recommendation

We suggest immediate resumption of chest compressions after shock delivery for adults in cardiac arrest in any setting (weak recommendation, very-low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-7](#). No change was made to this treatment recommendation. Although there is only very-low-certainty evidence addressing this question, worse short- and long-term outcomes have been reported with immediate rhythm checks after shock delivery. The effect of an immediate rhythm check on the incidence of VF recurrence is unclear. An observational study exploring this specific issue did not find that VF recurrence within 30 seconds of defibrillation (ie, successful shock)

was linked to the timing of resumption of chest compressions,<sup>149</sup> and this may not be a major factor affecting outcomes. Protocols including immediate cardiac rhythm check after shock delivery are reported to have reduced chest compression fractions; these increased pauses could be a potential cause of worse outcomes.

### Knowledge Gaps

- There were no studies that evaluated this question in the pediatric/in-hospital setting.
- No RCTs compared the specific intervention with standard care in any patient population, although 1 RCT assessed a CPR protocol characterized by different timing of rhythm checks, different compression-to-ventilation ratios, different duration of uninterrupted CPR between shocks, and different ventilation strategies.
- Currently available studies comparing different CPR protocols are characterized not only by different timing of rhythm checks but also by compression-to-ventilation ratios, compression intervals between shocks, and ventilation strategies that differ from standard care. More data are needed comparing groups receiving standard care with differences between control and intervention groups in only the timing of rhythm checks.

### Feedback for CPR Quality (BLS 361: SysRev)

#### Rationale for Review

CPR feedback or prompt devices are intended to improve CPR quality, probability of ROSC, and survival from cardiac arrest. Feedback devices involve technology that can measure various aspects of CPR mechanics, including ventilation rate, chest compression

mechanics (eg, depth, rate, recoil), and measures of flow time (CPR fraction, pre- and postshock pauses). These data can be presented to the provider in real time and/or provided in a summary report at the end of a resuscitation. Real-time displays can involve voice prompts, visual dials, numeric displays, wave forms, verbal prompts, and visual alarms. Visual displays enable the rescuer to see compression-to-compression quality parameters, including compression depth and rate in real time. Audio prompts may guide CPR rate (eg, metronome) and may offer verbal prompts to rescuers (eg, “push harder,” “good compressions”). Prompt devices that do not include the measurement and feedback of CPR quality metrics can include audible or visual metronomes set at the recommended rate for compressions or ventilation.

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with cardiac arrest
- Intervention: Real-time feedback and prompt devices regarding the mechanics of CPR quality (eg, rate and depth of compressions and/or ventilations)
- Comparator: No feedback
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC, bystander CPR rates, time to first compressions, time to first shock, and CPR quality were ranked as important outcomes.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Studies involving manikins only or the use of CPR quality data for delayed feedback (eg, debriefing or quality assurance programs) were excluded from this review.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to September 2019.

### **Consensus on Science**

Three discrete forms of real-time CPR guidance devices were identified: (1) digital audiovisual feedback, including corrective audio prompts; (2) analogue audio and tactile “clicker” feedback for chest compression depth and release; and (3) metronome guidance for chest compression rate. The analogue “clicker” device, designed to be placed on the patient’s chest under the hands of a CPR provider, involves a mechanism that produces a “click” noise and sensation when sufficient pressure is applied. Because there was considerable clinical heterogeneity across studies with respect to the type of devices used, the

mechanism of CPR quality measurement, the mode of feedback, patient types, locations (eg, in-hospital and out-of-hospital), and baseline (control group) CPR quality, we did not conduct any meta-analyses (Tables 7, 8, and 9).

### **Treatment Recommendations**

We suggest the use of real-time audiovisual feedback and prompt devices during CPR in clinical practice as part of a comprehensive quality improvement program for cardiac arrest designed to ensure high-quality CPR delivery and resuscitation care across resuscitation systems (weak recommendation, very-low-certainty evidence).

We suggest against the use of real-time audiovisual feedback and prompt devices in isolation (ie, not part of a comprehensive quality improvement program) (weak recommendation, very-low-certainty evidence).

### **Justification and Evidence-to-Decision Framework Highlights**

The evidence-to-decision table is included in [Supplement Appendix A-8](#). There was significant debate among task force members on whether to recommend for or against the use of these devices for real-time feedback on the basis of available data. On one side of the debate, the task force acknowledged that the bulk of higher-certainty data from key studies did not demonstrate a clinically or statistically significant association between real-time feedback and improved patient outcomes and that these devices require additional resources to purchase and implement. On the other side of the debate, we acknowledged several studies that demonstrated clinically important improvements in outcomes associated with the use of feedback devices. Most notable was the study by Goharani et al,<sup>159</sup> newly added to the evidence base considered in 2020, which was an RCT of 900 IHCA patients from Iran. This study demonstrated a +25.6% absolute increase in survival to hospital discharge with the use of an analogue “clicker” device that provided real-time feedback on compression depth and recoil (54% versus 28.4%;  $P<0.001$ ). Task force members did interpret this study to be supportive of the use of feedback devices; however, they also felt that this study represented an outlier. Members felt that replication of this result would be necessary before the task force could make any supportive recommendation for the specific type of device used in the study by Goharani et al.<sup>159</sup>

The task force also considered data from several observational studies demonstrating improvements in favorable neurological outcome that were not statistically significant. In addition, the task force considered statistically significant improvements in various aspects of CPR quality, including CPR rate

**Table 7. Real-Time Digital Audiovisual Feedback**

| Outcome                                      | Studies   | No. of Patients  | Results  |
|--|---|------------------|--|
| Survival with favorable neurological outcome | 1 cluster RCT, <sup>150</sup><br>low-certainty evidence (downgraded for very serious risk of bias)<br>4 observational, <sup>151–154</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)   | 1586<br><br>1100 | No difference:<br>Relative risk 1.02; 95% CI, 0.76–1.36; $P=0.9$<br>Absolute risk 0.19% (95% CI, –3.18% to 2.82%), or 2 more patients/1000 survived with the intervention (95% CI, 24 fewer patients/1000 to 36 more patients/1000 survived with the intervention)<br>Better outcome with feedback:<br>Adjusted odds ratio 2.69 (95% CI, 1.04–6.94) <sup>153</sup><br>No difference:<br>Adjusted relative risk 5.75%; 95% CI, –18.51% to 3.85% <sup>152</sup><br>Adjusted odds ratio 0.92; 95% CI, 0.37–2.30 <sup>151</sup><br>2/16 versus 0/16; $P=0.14$ <sup>154</sup>   |
| Survival to hospital discharge               | 1 cluster RCT, <sup>150</sup><br>low-certainty evidence (downgraded for very serious risk of bias)<br>6 observational: 5 in adults <sup>131,151–153,155</sup> and 1 in children, <sup>156</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)                     | 1586<br><br>1592 | No difference:<br>Relative risk: 0.91 (95% CI, 0.69–1.19); $P=0.5$<br>Absolute risk: –1.16% (95% CI, –4.37% to 2.02%), or 9 fewer patients/1000 survived with the intervention (95% CI, 31 fewer patients/1000 to 19 more patients/1000 survived with the intervention)<br>No difference:<br>Adjusted odds ratio 0.90; 95% CI, 0.39–2.06; $P=0.80$ ; <sup>151</sup><br>Adjusted relative risk –0.91; 95% CI, –11.18 to 12.33; <sup>152</sup><br>Adjusted relative risk 5.23; 95% CI, –0.49 to 10.89; <sup>153</sup><br>Adjusted relative risk –0.18; 95% CI, –11.46 to 8.64; <sup>155</sup><br>Adjusted relative risk 1.37; 95% CI, –2.47 to 6.91; <sup>131</sup><br>8 children (ages 1–7 y) with IHCA (1/4 versus 1/4) <sup>156</sup>   |
| Survival to 30 days                          | 1 observational, <sup>157</sup><br>very-low-certainty evidence (downgraded for serious risk of bias)  | 196              | No difference:<br>Adjusted relative risk –0.84; 95% CI, –13.88 to 14.82; $P=0.9$ <sup>157</sup>  |
| Survival to 24 h                             | 1 cluster RCT, <sup>150</sup><br>low-certainty evidence (downgraded for very serious risk of bias)<br>2 observational, <sup>152,154</sup> very-low-certainty evidence (downgraded for very serious risk of bias)  | 1586<br><br>219  | No difference:<br>Relative risk 0.96 (95% CI, 0.82–1.13; $P=0.6$ ); ARR, –1.09% (95% CI, –3.35% to 5.50%), or 4 fewer patients/1000 survived with the intervention (95% CI, 18 fewer patients/1000 to 13 more patients/1000 survived with the intervention)<br>No difference:<br>2/16 versus 0/16 <sup>154</sup><br>Adjusted relative risk 13.13; 95% CI, –0.66 to 28.02 <sup>152</sup>  |
| ROSC   | 1 cluster RCT, <sup>150</sup><br>low-certainty evidence (downgraded for very serious risk of bias)<br>8 observational <sup>152,154</sup> : 7 in adults <sup>131,151–155,158</sup> and 1 in children, <sup>156</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias) | 1586<br><br>2263 | No difference:<br>Relative risk 1.01 (95% CI, 0.91–1.13; $P=0.9$ ); Adjusted relative risk –0.45% (95% CI, –5.33% to 4.43%), or 1 more patient/1000 survived with the intervention (95% CI, 9 fewer patients/1000 to 13 more patients/1000 survived with the intervention)<br>No benefit<br>9/16 versus 10/16 <sup>154</sup><br>Adjusted odds ratio 0.62; 95% CI, 0.31–1.22; $P=0.17$ ; <sup>151</sup><br>Adjusted relative risk –3.17; 95% CI, –10.73 to 4.35; <sup>153</sup><br>Adjusted relative risk –4.39; 95% CI, –3.35 to 12.06; <sup>158</sup><br>Adjusted relative risk 4.55; 95% CI, –11.59 to 19.90; <sup>155</sup><br>Adjusted relative risk 5.65; 95% CI, –2.89 to 15.09; <sup>131</sup><br>Adjusted relative risk 1.11; 95% CI, –15.56 to 13.69; $P=0.9$ ; <sup>131,151–155,158</sup><br>8 children (ages 1–7 y): 3/4 versus 1/4 <sup>156</sup><br>Better outcome with feedback:<br>Adjusted relative risk 17.55; 95% CI, 1.79–32.46) <sup>152</sup> |
| Chest compression rate                       | 1 cluster RCT, <sup>150</sup><br>moderate-certainty evidence<br>6 observational: 5 in adults <sup>131,151,153–155</sup> and 1 in children, <sup>156</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)   | 1586<br><br>1441 | Better CPR quality with feedback:<br>Difference of –4.7/min (95% CI, –6.4 to –3.0/min) when feedback was used<br>No difference:<br>One observational study <sup>155</sup><br>Better CPR quality with feedback:<br>4 observational studies <sup>131,151,153,154</sup> showed lower compression rates in the group with CPR feedback<br>The pediatric study <sup>156</sup> found a median difference of –10/min with feedback.   |
| Compression depth                            | 1 cluster RCT, <sup>150</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)<br>6 observational: 5 in adults <sup>131,151,153–155</sup> and 1 in children, <sup>156</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)                | 1586<br><br>1441 | Better CPR quality with feedback:<br>Significant +1.6 mm (95% CI, 0.5–2.7 mm) (cluster-adjusted) difference in chest compression depth with feedback.<br>Better CPR quality with feedback:<br>Three observational studies <sup>131,153,154</sup> showed deeper chest compressions in the groups with CPR feedback <sup>131,153,154</sup><br>No difference:<br>One observational study <sup>155</sup> ;<br>the pediatric study <sup>156</sup> found no difference in median compression depth.  |

(Continued)

**Table 7. Continued**

| Outcome                    | Studies  | No. of Patients  | Results  |
|----------------------------|--|------------------|--|
| Chest compression fraction | 1 cluster RCT, <sup>150</sup><br>moderate-certainty evidence<br>6 observational: 5 in adults <sup>131,151,153–155</sup><br>and 1 in children, <sup>156</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias) | 1586<br><br>1441 | Better CPR quality with feedback:<br>Difference of +2% (66% compared with 64%; $P=0.016$ )<br>Better CPR quality with feedback:<br><br>2 studies reported statistically significant increases in CPR fraction associated with feedback <sup>151,155</sup><br>No difference:<br>3 studies did not observe a statistically or clinically important difference. <sup>131,153,154</sup><br>The sample size of the pediatric study <sup>156</sup> was too small to enable inferential statistical analysis. |
| Ventilation rate           | 1 cluster RCT, <sup>150</sup><br>moderate-certainty evidence<br>3 observational, <sup>131,153,155</sup><br>very-low-certainty evidence (downgraded for very serious risk of bias)  | 1586<br><br>1001 | No difference<br><br>No difference   |

ARR indicates adjusted relative risk; CPR, cardiopulmonary resuscitation; RCT, randomized controlled trial; and ROSC, return of spontaneous circulation.

and CPR fraction, associated with the use of feedback devices.

The task force also felt that a permissive recommendation was appropriate because of the role that these devices play in CPR quality monitoring, benchmarking, and quality improvement programs by collecting data across patients treated by a system. These roles were not included in the scope of this PICOST; however, the task force was concerned that a recommendation against the use of these devices for real-time feedback would discourage use for other important activities. The task force also recognized that implementing and maintaining high-quality CPR in hospital and EMS systems would be difficult without the use of these devices to provide an objective method of CPR quality measurement in those systems.

In summary, the task force agreed that CPR feedback devices that measure aspects of CPR quality were reasonable to consider for healthcare systems, given the importance of high-quality CPR. Without any signal of patient harm in the data reviewed, we agreed that a weak recommendation in favor of their use in this manner was appropriate.

We also agreed that there was no consistent signal from the data reviewed indicating that the real-time feedback function of these devices has a significant effect on individual cardiac arrest patient outcomes, suggesting that the devices should not be implemented for this reason alone outside of a comprehensive quality assurance program.

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- What is the effect of feedback devices on patient outcomes when used by lay people with AEDs?
- Is there an interaction between the effect of real-time feedback devices and the skill set of the provider (eg, in low-performing services with baseline CPR metrics) that are below recommended values?
- What are the most effective parameters to feedback to users (ie, measures of brain or other tissue perfusion, electrocardiographic characteristics, other physiological measurements)?
- What are the most effective modalities for feedback to be provided to users?

**Table 8. Analogue Audio and Tactile “Clicker” Feedback**

| Outcome                        | Studies   | No. of Patients | Results   |
|--------------------------------|---|-----------------|---|
| Survival to hospital discharge | 1 RCT, <sup>159</sup><br>very-low-certainty evidence (downgraded for serious risk of bias)      | 900             | Better outcome with feedback:<br>Relative risk 1.90 (95% CI, 1.60–2.25; $P<0.001$ );<br>Adjusted relative risk 25.56% (95% CI, 19.22%–31.60%), or 91 more patients/1000 survived with the intervention (95% CI, 61 more patients/1000 to 126 more patients/1000 survived with the intervention)   |
| ROSC                           | 2 RCTs, <sup>159,160</sup><br>very-low-certainty evidence (downgraded for serious risk of bias) | 980             | Better outcome with feedback:<br>Relative risk 1.57 (95% CI, 1.38–1.78; $P<0.001$ );<br>Adjusted relative risk 24.22% (95% CI, 17.79%–30.36%), or 58 more patients/1000 survived with the intervention (95% CI, 38 more patients/1000 to 79 more patients/1000) <sup>159</sup><br>Relative risk 2.07 (95% CI, 1.20–3.29; $P<0.001$ );<br>Adjusted relative risk 37.50% (95% CI, 15.70%–54.68%), or 108 more patients/1000 survived with the intervention (95% CI, 20 more patients/1000 to 232 more patients/1000) <sup>160</sup> |

RCT indicates randomized controlled trial; and ROSC, return of spontaneous circulation.



**Table 9. Metronome Rate Guidance**

| Outcome             | Studies   | No. of Patients | Results   |
|---------------------|---|-----------------|---|
| Survival to 30 days | 1 observational, <sup>157</sup> very-low-certainty evidence (downgraded for serious risk of bias)     | 196             | No difference: Relative risk 1.66; 95% CI, -17.71 to 14.86; $P=0.8^{157}$   |
| Survival to 7 days  | 1 observational, <sup>161</sup> very-low-certainty evidence (downgraded for serious risk of bias)     | 30              | No difference: 3/17 versus 2/13; $P=ns^{161}$   |
| ROSC                | 2 observational, <sup>157,161</sup> very-low-certainty evidence (downgraded for serious risk of bias) | 226             | No difference: Adjusted relative risk 4.97; 95% CI, -21.11 to 11.76; $P=0.6^{157}$ 7/13 versus 8/17; $P=ns^{161}$ |

ROSC indicates return of spontaneous circulation.

## ALTERNATIVE TECHNIQUES

### Alternative Techniques (Cough CPR, Precordial Thump, Fist Pacing) (BLS 374: SysRev)

#### Rationale for Review

Reports of "cough CPR" circulate on social media, and this technique may be perceived by the public as an effective way of preventing cardiac arrest. Precordial thumping and fist pacing are techniques previously recommended to healthcare professionals. In this review, we update the available evidence for these alternative techniques.

#### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with cardiac arrest
- Intervention: Cough CPR; precordial thump; fist pacing
- Comparator: Standard CPR
- Outcome: Survival with favorable neurological outcome until and beyond hospital discharge or 30 days; survival until and beyond hospital discharge or 30 days; ROSC
- Study design: We included RCTs, nonrandomized studies, and case series with at least 5 cases. We considered papers in all languages provided there was an English language abstract available for review. We excluded unpublished studies, conference abstracts, manikin or simulation studies, narrative reviews, editorials or opinions with no

primary data, animal studies, and experimental/laboratory models.

- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 2019.
- PROSPERO registration: CRD42019152925

#### Consensus on Science

##### Cough CPR

For the critical outcome of survival to hospital discharge<sup>162</sup> and important outcome of restoration of cardiac output/circulation (at or shortly after the onset of a potentially nonperfusing rhythm in which the patient has not yet lost consciousness or cardiac output),<sup>163–165</sup> we identified only 4 observational studies. All studies were in adult patients only. The overall certainty of evidence was rated as very low for all outcomes as a result of very serious risk of bias. For this reason and because of a high degree of heterogeneity across studies, no meta-analyses could be performed, and individual studies were difficult to interpret. Additional information may be found in Table 10.

##### Precordial Thump

For the critical outcomes of survival to hospital discharge, we identified 5 observational studies.<sup>162,166–169</sup> Two of these studies, both out-of-hospital, directly compared precordial thump with standard CPR.<sup>166,167</sup> For the important outcome of ROSC, we identified 1 observational study.<sup>170</sup> For the important outcome of restoration of cardiac output/circulation, we identified 10 observational studies.<sup>171–180</sup> All studies were in adult

**Table 10. Observational Studies of Cough CPR for Conscious Patients With No Comparator Group**

| Outcome                        | Certainty                            | Studies  | No. of Patients   | Results   |
|--------------------------------|--------------------------------------|--|---|---|
| Survival to hospital discharge | Very low (very serious risk of bias) | Caldwell 1985 <sup>162</sup>   | 6 (in-hospital VT)  | 6/6 (100%), selective reporting of cases achieving outcome  |
| ROSC                           | Very low (very serious risk of bias) | Marozsan 1990, Nieman 1980 <sup>164,165</sup> , Petelenz 1998 <sup>163</sup> | 20 (in-hospital, 2 studies): n=6 VF, n=13 asystole, n=1 bradycardia; 66 (out-of-hospital, 1 study): rhythms unknown | In-hospital: 18/20 (90%), selective reporting of cases achieving outcome in 1 study (n=7) <sup>165</sup> ; out-of-hospital: 66/66 (100%), selective reporting of cases achieving outcome <sup>163</sup> |

CPR indicates cardiopulmonary resuscitation; ROSC, return of spontaneous circulation; VF, ventricular fibrillation; and VT, ventricular tachycardia.

patients only. The overall certainty of evidence was rated as very low for all outcomes primarily because of very serious risk of bias. Because of this and a high degree of heterogeneity across the studies, no meta-analyses could be performed, and individual studies were difficult to interpret. Additional information may be found in Tables 11 and 12.

### Fist Pacing

For the critical outcome of survival to hospital discharge,<sup>181,182</sup> the important outcome of ROSC,<sup>183</sup> and the important outcome of restoration of cardiac output/circulation,<sup>184</sup> we identified only 4 observational studies. One study included children (age range, 11–84 years).<sup>181</sup> The overall certainty of evidence was rated as very low for all outcomes, mainly because of very serious risk of bias. Because of this and a high degree of heterogeneity, no meta-analyses could be performed, and individual studies were difficult to interpret. Additional information may be found in Table 13.

### Treatment Recommendations

We recommend against the routine use of cough CPR for cardiac arrest (strong recommendation, very-low-certainty evidence).

We suggest that cough CPR may be considered only as a temporizing measure in the exceptional circumstance of a witnessed, monitored IHCA (eg, in a cardiac catheterization laboratory) if a nonperfusing rhythm is recognized promptly before loss of consciousness (weak recommendation, very-low-certainty evidence).

We recommend against the use of a precordial thump for cardiac arrest (strong recommendation, very-low-certainty evidence).

We recommend against fist pacing for cardiac arrest (strong recommendation, very-low-certainty evidence).

We suggest that fist pacing may be considered only as a temporizing measure in the exceptional circumstance of a witnessed, monitored, IHCA (eg, in a cardiac catheterization laboratory) due to bradycardia if such a nonperfusing rhythm is recognized promptly before loss of consciousness (weak recommendation, very-low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-9](#). This topic was last reviewed in the 2010

*International Consensus on CPR and ECC Science With Treatment Recommendations*.<sup>1,2</sup> Although treatment recommendations remain essentially unchanged, the BLS Task Force has tried to update the recommendations with the intention of clarifying the special circumstances when these alternative techniques might be appropriate.

The very-low-quality evidence identified precludes meaningful meta-analysis. Two studies (both on precordial thump) had a direct comparator group (standard CPR), and both had a very serious risk of bias. The others were limited case series or cohorts without comparator groups.

Cough CPR is described as a repeated deep breath followed by a cough every few seconds. There is no evidence for the effectiveness of cough CPR in established cardiac arrest (ie, in an unconscious, pulseless patient), nor is its initiation even feasible under such circumstances. Very-low-quality evidence from 1 study<sup>163</sup> addresses the use of cough CPR for prodromal symptoms of collapse in high-risk patients in whom the cardiac rhythm was not known and the likelihood of progressing to cardiac arrest was uncertain. Suggesting a benefit of cough CPR for the general population would require us to accept that an untrained patient could reliably identify a cardiac arrest rhythm in time to initiate coughing to maintain a cardiac output. This seems highly unlikely.

There are periodic stories (on social media, for example) instructing members of the public to perform cough CPR in case of imminent collapse, so it is important that we address this topic. We should be clear that we do not recommend cough CPR for OHCA. The risks are (1) that it delays effective treatment (early call for help, early CPR and defibrillation if the patient loses consciousness and stops breathing normally) and (2) that members of the public confusing “cardiac arrest” with “heart attack” delay seeking help when suffering chest pain or other symptoms indicating a possible ischemic cardiac event.

There is no evidence to contradict the 2010 CoSTR treatment recommendation<sup>1,2</sup> that providers can consider cough CPR in the exceptional circumstance of monitored, witnessed in IHCAs. The victim must remain conscious and be able to follow instructions for coughing. There is limited very-low-certainty evidence that this may be effective in all arrhythmias that can cause cardiac arrest, not limited to just VF and VT. This evidence is reported for adult patients only. There is some evidence that cough CPR increases aortic, left atrial, and

**Table 11. Observational Studies of Precordial Thump With Comparator Group**

| Outcome                        | Certainty   | Studies  | No. of Patients                              | Results  |
|--------------------------------|---|--|--|--|
| Survival to hospital discharge | Very low (downgraded for very serious risk of bias) | Nehme 2013, <sup>166</sup><br>Pellis 2009 <sup>167</sup> | 797 (n=500 VF/VT, n=101 PEA, n=196 asystole) | No difference: 71% versus 70% (P=ns) <sup>166</sup> and 5.6% versus 6.4% (P=ns) <sup>167</sup> |
| ROSC                           | Very low (downgraded for very serious risk of bias) | Nehme 2013, <sup>166</sup><br>Pellis 2009 <sup>167</sup> | 797 (n=500 VF/VT, n=101 PEA, n=196 asystole) | No difference: 93% versus 90% (P=ns) <sup>166</sup> and 22% versus 20% (P=ns) <sup>167</sup>   |

ns indicates nonsignificant; PEA, pulseless electric activity; ROSC, return of spontaneous circulation; VT, ventricular tachycardia; and VF, ventricular fibrillation.

**Table 12. Observational Studies of Precordial Thump With No Comparator Group**

| Outcome                        | Certainty                            | Studies  | No. of Patients   | Results  |
|--------------------------------|--------------------------------------|--|---|--|
| Survival to hospital discharge | Very low (very serious risk of bias) | Caldwell 1985, <sup>162</sup> Gertsch 1992, <sup>168</sup> Rajagopalan 1971 <sup>169</sup> ; Caldwell 1985 <sup>162</sup>  | 35 (in-hospital, 3 studies): n=29 VT, n=2 VF, n=2 asystole, n=2 unknown; 3 (out-of-hospital, 1 study): n=1 VT, n=2 VF | In-hospital: 20/35 (57%); 2/2 (100%) VF, 14/29 (48%) VT, 2/2 (100%) asystole, 2/2 (100%) unknown; out-of-hospital: 2/3 (67%)   |
| ROSC                           | Very low (very serious risk of bias) | Miller 1984, <sup>170</sup> Rahner 1978, <sup>171</sup> Cotoi 1980, <sup>172</sup> Pennington 1970, <sup>173</sup> Morgera 1979, <sup>174</sup> Haman 2009, <sup>175</sup> Amir 2007, <sup>176</sup> Befeler 1978, <sup>177</sup> Miller 1985, <sup>178</sup> Nejima 1991, <sup>179</sup> Volkmann 1990 <sup>180</sup> | 50 (out-of-hospital): n=27 VT, n=23 VF; 366 (in-hospital): n=320 VT, n=38 VF, n=8 Morgagni-Adams-Stokes attack        | Out-of-hospital: 23/50 (46%); 11/27 (41%) VT, 12/23 (52%) VF; 88/366 (24%); in-hospital: 80/320 (25%) VT, 8/8 (100%) Morgagni-Adams-Stokes, 0/38 (0%) VF; selective reporting of cases achieving outcome in 3 studies (n=39: n=31 VT, n=8 Morgagni-Adams-Stokes <sup>171–173</sup> ) |

ROSC, return of spontaneous circulation; VF, ventricular fibrillation; and VT, ventricular tachycardia.

left ventricular pressures, but a causative link between cough CPR and termination of malignant arrhythmias is lacking. It would not be appropriate to prioritize cough CPR instead of other measures with proven efficacy, but clinicians may consider it as a temporary measure if there is a delay to defibrillation.

A precordial thump is described as a sharp, high-velocity blow to the middle of the sternum with immediate retraction by the ulnar aspect of the fist. We weighed the potential benefit of precordial thumps against the potential for harm. A precordial thump can potentially interrupt life-threatening VT by generating an electric impulse, resulting in a premature ventricular depolarization. However, there is a risk of deterioration of cardiac rhythm (from VT to VF, akin to an “R on T” phenomenon), reported in some studies,<sup>170,171</sup> and a risk of delaying CPR or defibrillation. Delay to definitive treatment is of particular concern in situations when lay rescuers are providing cardiac arrest interventions.

A causal link between precordial thump and the critical outcomes of survival to hospital discharge and ROSC is lacking. Defibrillation is a more effective treatment for the termination of VF and VT and should be prioritized. There is concern from 1 study (very-low-certainty evidence) that use of precordial thump could compromise first shock success.<sup>166</sup>

In many of the included studies, it is unclear whether the tachyarrhythmia (VT) represents cardiac arrest or impending loss of cardiac output. It is very likely that this is not so for many of the cases included in the studies reviewed.

Across studies, there is a lack of standardization in the technique of precordial thump, the number of

times it was used, pharmacological therapy delivered before or after its delivery, and—in some cases—its timing related to the onset of the tachyarrhythmia.

Fist (or percussion) pacing is described as the delivery of serial, rhythmic, relatively low-velocity blows to the sternum by a closed fist. The evidence for the effectiveness of fist pacing is limited to a few small case series (totaling 147 patients among them) suggesting that cardiac output can be maintained if fist pacing is initiated very quickly after onset of asystole or severe bradycardia—and strictly for such rhythms. An electric impulse is generated sufficient to cause myocardial depolarization and contraction. Fist pacing is not used for tachyarrhythmias.

There is no evidence comparing fist pacing with standard CPR (chest compressions) in established bradysystolic cardiac arrest. We again highlight the importance of prompt, high-quality chest compressions for the treatment of cardiac arrest.

There is no evidence to contradict the 2010 CoSTR treatment recommendation<sup>1,2</sup> that providers can consider fist pacing in the exceptional circumstance of monitored, witnessed IHCA due to bradysystole. It would not be appropriate to prioritize fist pacing instead of other measures with proven efficacy, but clinicians may consider it as a temporary measure if there is a delay to electric pacing or pharmacological therapies.

### Knowledge Gaps

- There are no data directly comparing cough CPR or fist pacing with standard CPR.
- There are no data for any alternative CPR technique assessing survival with a favorable neurological outcome.

**Table 13. Observational Studies of Fist Pacing With No Comparator Group**

| Outcome                        | Certainty                            | Studies  | No. of Patients   | Results   |
|--------------------------------|--------------------------------------|--|---|---|
| Survival to hospital discharge | Very low (very serious risk of bias) | Klumbies 1988, <sup>181</sup> Scherf 1960 <sup>182</sup> | 111 (in-hospital): n=51 asystole, n=20 “life-threatening bradycardia,” n=29 unclear/delayed monitoring, n=11 “ventricular standstill” | 63/111 (57%)  |
| ROSC                           | Very low (very serious risk of bias) | Iseri 1987 <sup>183</sup> ; Paliege 1982 <sup>184</sup>  | 5 (in-hospital): all asystole; 42 (in-hospital): n=35 asystole, n=7 “extreme bradycardia”   | 5/5 (100%); selective reporting of cases achieving outcome; 41/42 (98%) |

ROSC indicates return of spontaneous circulation.

- There is limited, very-low quality evidence assessing the critical outcome of survival to hospital discharge.
- There are no data on any outcome after alternative CPR techniques performed in children.

## DEFIBRILLATION

### Public Access AED Programs (BLS 347: SysRev)

#### Rationale for Review

This topic was prioritized for review by the BLS Task Force because it had not been updated since 2015.<sup>3,4</sup> Public access AED programs were recommended by ILCOR after review of the evidence before 2015, and since then several additional studies have been published.

#### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA
- Intervention: Implementation of a public access AED program
- Comparator: Traditional EMS response
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC, bystander CPR rates, time to first compressions, time to first shock, and CPR quality were ranked as important outcomes.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded.
- Time frame: All years and all languages were included as long as there was an English abstract. The literature search was updated to October 2019.

#### Consensus on Science

SysRevs on the effects of public access defibrillation (PAD) on OHCA survival have been published previously.<sup>185,186</sup> This review is focused on comparing outcomes in systems with public access AED programs versus systems with traditional EMS response and included 1 RCT and 30 observational studies. PAD is defined as defibrillation with an onsite AED by a layperson in the OHCA setting. The PAD group included only patients defibrillated by a lay person using an onsite AED. The control group included all patients not receiving PAD—meaning not treated with an onsite AED by a lay person—and included patients defibrillated by professional first responders, such as police or firefighters.

For the critical outcome of survival to 1 year with favorable neurological outcome, we identified low-certainty evidence (downgraded for risk of bias) from

1 observational trial<sup>187</sup> enrolling 62 patients showing improvement (43% versus 0%;  $P=0.02$ ) after a PAD program in a subway system.

For the critical outcome of survival to 30 days with favorable neurological outcome, we identified low-certainty evidence (downgraded for risk of bias and inconsistency) from 7 observational studies<sup>188–194</sup> enrolling 43 116 patients demonstrating improved survival with a PAD program (OR, 6.60; 95% CI, 3.54–12.28).

For the critical outcome of survival to hospital discharge with favorable neurological outcome, we identified low-certainty evidence (downgraded for risk of bias) from 8 observational studies. The studies<sup>187,195–201</sup> included 11 837 patients demonstrating improved survival with PAD program (OR, 2.89; 95% CI, 1.79–4.66).

For the critical outcome of survival to 30 days, we identified low-certainty evidence (downgraded for risk of bias) from 8 observational studies<sup>189,190,192,193,202–205</sup> enrolling 85 589 patients demonstrating improved outcome with a PAD program (OR, 3.66; 95% CI, 2.63–5.11).

For the critical outcome of survival to hospital discharge, we identified moderate-certainty evidence (downgraded for risk of bias) from 1 RCT<sup>206</sup> enrolling 235 OHCA patients showing improved survival with PAD compared with no PAD (RR, 2.0; 95% CI, 1.07–3.77) and low-certainty evidence (downgraded for risk of bias) from 16 observational studies enrolling 40 243 patients showing improved survival associated with PAD programs (OR, 3.24; 95% CI, 2.13–4.92).<sup>195–199,201,207–217</sup>

#### Treatment Recommendation

We recommend the implementation of PAD programs for patients with OHCA (strong recommendation, low certainty evidence).

#### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-10](#). PAD programs are implemented at the community level to improve outcomes for patients with OHCA. In making this recommendation, we placed a high value on the potential life-saving capability of an AED for a shockable rhythm and on keeping with the previous treatment recommendation when there were no compelling data suggesting the need to change. We recognize that there are barriers to the implementation of PAD programs. The ILCOR scientific statement on public access defibrillation addresses key interventions (early detection, optimizing availability, signage, novel delivery methods, public awareness, device registration, mobile apps for AED retrieval and personal access defibrillation) that should be considered as part of all PAD programs. Cost-effectiveness of PAD programs may vary according to country. A recent review found cost-effectiveness ratios between 37 200 and 1 152 400 US dollars/quality-adjusted life-years.<sup>185</sup> Another recent cost-effectiveness analysis study<sup>218</sup> from



the United States concluded that public access AEDs are a cost-effective public health intervention.

Among 31 included studies, there was only 1 RCT, which showed improved survival to discharge in the CPR-plus-AED group compared with the CPR-only group. Observational studies were mostly retrospective analyses of data from large registries and generally showed improved survival outcomes associated with PAD. However, there were some inconsistencies among the observational studies, as some were unable to show any significant differences in outcomes.<sup>187,193,196,215</sup> There was also important heterogeneity among studies in the meta-analysis. The location of cardiac arrest was various and included airports,<sup>212</sup> subways,<sup>187</sup> and sports facilities.<sup>200</sup> The population varied, with 2 studies including only children.<sup>190,194</sup> The control group also varied among studies because some patients in control groups received first responder defibrillation, whereas others did not. Some studies were before-and-after studies in which historic controls included periods before PAD implementation<sup>193,215,217</sup> or the initial period of implementation.<sup>187</sup> Despite such heterogeneity, all patients in those studies had OHCA, and most studies showed that implementation of PAD improved survival.

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- Optimal placement/location of AEDs
- Optimal role of emergency medical dispatchers in identifying nearest AED and alerting callers to their location
- How AEDs could be most effectively integrated into citizen responder programs

## Analysis of Rhythm During Chest Compressions (BLS 373: SysRev)

### Rationale for Review

High-quality CPR with few pauses in chest compressions is emphasized in current guidelines and CPR training. Rhythm analysis and pulse checks require pauses in chest compressions, and artifact-filtering algorithms for analysis of electrocardiographic rhythm during CPR have been proposed as a method to reduce pauses in chest compressions.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with cardiac arrest
- Intervention: Analysis of cardiac rhythm during chest compressions
- Comparator: Standard care (analysis of cardiac rhythm during pauses in chest compressions)
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital

discharge were ranked as critical outcomes. ROSC was ranked as an important outcome. CPR quality metrics, such as time of chest compression fraction, pauses in compressions, compressions per minute, time to commencing CPR, time to first shock, etc, were included as important outcomes.

- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to September 23, 2019.

### Consensus on Science

Fourteen full-text papers were identified and reviewed,<sup>219–232</sup> but none assessed any critical or important patient-related outcomes. Most of these studies use previously collected electrocardiographs, electric impedance, and/or accelerometer signals recorded during CPR for cardiac arrest to evaluate the ability of various algorithms<sup>220–229</sup> or machine learning<sup>230</sup> to detect shockable rhythms during chest compressions. Although these studies did not evaluate the effect of the artifact-filtering algorithms on any critical or important outcomes, they provided insights into the feasibility and potential benefits of this technology. We also identified studies evaluating artifact-filtering algorithms in animal models of cardiac arrest<sup>219,231</sup> and simulation studies.<sup>232</sup> Sensitivities and specificities are generally reported in the 90% to 99% range, but none of these studies evaluated the use of this technology during actual cardiac arrest and resuscitation.

### Treatment Recommendations

We suggest against the routine use of artifact-filtering algorithms for analysis of electrocardiographic rhythm during CPR (weak recommendation, very-low-certainty evidence).

We suggest that the usefulness of artifact-filtering algorithms for analysis of electrocardiographic rhythm during CPR be assessed in clinical trials or research initiatives (weak recommendation, very-low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-11](#). In making a recommendation against routine use, we placed priority on avoiding the costs of introducing a new technology when its effects on patient outcomes and risk of harm remain to be determined.

In making a recommendation for further research; the task force is acknowledging that (1) there is thus far

insufficient evidence to support a decision for or against routine use, (2) further research has potential for reducing uncertainty about the effects, and (3) further research is thought to be of good value for the anticipated costs. This treatment recommendation was changed from a previous weak suggestion that, for EMS systems that had already integrated artifact-filtering algorithms into clinical practice, it would be reasonable to continue with their use.<sup>3,4</sup> The task force acknowledges that some EMS systems may have implemented artifact-filtering algorithms for analysis of electrocardiographic rhythm during CPR and strongly encourages such systems to report their experiences to build the evidence base about the use of these technologies in clinical practice.

### Knowledge Gaps

There were no studies identified that evaluated feasibility, efficacy, or effectiveness of artifact-filtering algorithms for analysis of electrocardiographic rhythm during CPR in any setting for any patient population.

## CPR Before Defibrillation (BLS 363: SysRev)

### Rationale for Review

Previous treatment recommendations for CPR before defibrillation have been based on RCTs, but the results from these trials have been inconsistent, and important uncertainty about the optimal timing of defibrillation remains. This topic has not been reviewed by ILCOR since the 2015 CoSTR<sup>3,4</sup> and therefore was prioritized by the BLS Task Force.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with cardiac arrest and a shockable rhythm at initiation of CPR

- Intervention: A prolonged period of chest compressions before defibrillation (90–180 seconds)
- Comparator: A short period of chest compressions until the defibrillator is ready
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 27, 2019.

### Consensus on Science

Five RCTs were identified comparing a shorter with a longer interval of chest compressions before defibrillation.<sup>117,118,233–235</sup> Outcomes assessed varied from 1-year survival with favorable neurological outcome to ROSC. No clear benefit from CPR before defibrillation was found in meta-analysis of any of the critical or important outcomes (Table 14).

### Treatment Recommendation

This treatment recommendation (below) is modified slightly from the 2015 CoSTR.<sup>3,4</sup> We suggest a short period of CPR until the defibrillator is ready for analysis and/or defibrillation in unmonitored cardiac arrest. (weak recommendation, low-certainty evidence).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-12](#). This topic was prioritized by the

**Table 14. CPR Before Defibrillation**

| Outcome  | Certainty                        | Studies  | No. of Patients | Results  |
|--|----------------------------------|--|-----------------|--|
| 1 y with favorable neurological outcome                | Low (risk of bias, imprecision)  | Wik 2003 <sup>117</sup>  | 200             | No difference:<br>Relative risk 1.15 (95% CI, –0.57 to 2.34), 19 more patients/1000 (–54 to 167) |
| Hospital discharge with favorable neurological outcome | Low (inconsistency, imprecision) | Wik 2003, <sup>117</sup> Baker 2008, <sup>118</sup> Ma 2012, <sup>234</sup> Stiell 2011 <sup>235</sup>                             | 10 424          | No difference:<br>Relative risk 1.02 (95% CI, –0.01 to 0.01), 1 more patient/1000 (–7 to 11)     |
| Survival to 1 y  | Low (risk of bias, imprecision)  | Wik 2003, <sup>117</sup> Jacobs 2005 <sup>233</sup>  | 456             | No difference:<br>Relative risk 1.19 (95% CI, 0.69–2.04), 18 more patients/1000 (–29 to 98)      |
| Survival to hospital discharge                         | Low (risk of bias, imprecision)  | Wik 2003, <sup>117</sup> Jacobs 2005, <sup>233</sup> Baker 2008, <sup>118</sup> Ma 2012, <sup>234</sup> Stiell 2011 <sup>235</sup> | 10 680          | No difference:<br>Relative risk 1.01 (95% CI, 0.90–1.15), 1 more patient/1000 (–8 to 13)         |
| ROSC   | Low (risk of bias, imprecision)  | Wik 2003, <sup>117</sup> Jacobs 2005, <sup>233</sup> Baker 2008, <sup>118</sup> Ma 2012, <sup>234</sup> Stiell 2011 <sup>235</sup> | 10 680          | No difference:<br>Relative risk 1.03 (95% CI, 0.97–1.10), 8 more patients/1000 (–9 to 27)        |

CPR indicates cardiopulmonary resuscitation; and ROSC, return of spontaneous circulation. Both relative and absolute risks are written as mean values (95% CIs).

BLS Task Force, as it had not been reviewed since the 2015 CoSTR.<sup>3,4</sup> Given the availability of comparative data from several RCTs, we did not include non-RCTs. No new RCTs were identified, and no changes were made to the treatment recommendation; however, because the outcome templates have been altered for the 2020 ILCOR review process, the review has been updated.

In continuing to make the recommendation to provide CPR until the defibrillator is ready for analysis and/or defibrillation in unmonitored cardiac arrest, we placed a high value on being consistent with previous recommendations. The BLS Task Force acknowledges that every change in guidelines comes with a significant risk and cost as CPR educators and providers are asked to change current practice and implement new treatment strategies for complex and high-stress medical emergencies.

Important issues remained in the evaluation of the 5 included RCTs and led the BLS Task Force to downgrade the certainty of the treatment recommendation. The trial by Jacobs et al<sup>233</sup> did not use a random sequence generation and did not conceal randomization before rhythm analysis, leading to potential bias. In all RCTs, the treating EMS personnel could not be blinded to the interventional strategy after randomization. There was also significant heterogeneity in these trials with regard to the duration of CPR provided before defibrillation, with a range of 90 to 180 seconds. For the purposes of this review, the 90 to 180 seconds of CPR was considered a combined group. It is also important to note that the trials were conducted in different countries (Australia, Canada, Norway, Taiwan, United States) with varying EMS system structural configurations (BLS, ALS, physician on scene) as well as response times and treatment protocols. Only 1 of the included trials attempted to document and adjust for the quality of the intervention (or chest compressions) before defibrillation,<sup>235</sup> leaving the possibility that the intervention in the other trials was of varying quality. The studies also included only adult (age  $\geq 18$  years) OHCA patients and cannot be generalized to the IHCA or pediatric populations.

Two subgroup analyses were considered in the 2015 CoSTR. One subgroup analysis looked at enrollments based on EMS response interval, comparing those with intervals of less than 4 to 5 minutes versus those with intervals of 4 to  $\geq 5$  minutes. Within this subgroup, 1 study<sup>117</sup> found a favorable relationship with CPR for 180 seconds before defibrillation when the response interval was  $\geq 5$  minutes, but this relationship was not confirmed in 3 other RCTs.<sup>118,233,235</sup> The second subgroup analysis<sup>236</sup> examined outcomes from early compared with late analysis on the basis of baseline EMS agency VF/pVT survival rates. Among EMS agencies with low baseline survival to hospital discharge (defined as less than 20% for an initial rhythm of VF/pVT), higher neurologically favorable survival was associated with early analysis and

shock delivery as opposed to CPR and delayed analysis and shock delivery. Yet, for EMS agencies with higher baseline survival to hospital discharge (greater than 20%), 3 minutes of CPR followed by analysis and defibrillation resulted in higher neurologically favorable survival. These subgroup analyses underscore the difficulty in making "one size fits all" recommendations for resuscitation systems, which may vary considerably in both populations served and treatments offered.

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- What effect does the quality of bystander CPR have?
- Can electrocardiographic waveform characteristics be used to determine optimal strategy?
- If a CPR-first strategy is adopted, what is the optimal duration of CPR (90 seconds, 120 seconds, or 180 seconds)?
- What system-level characteristics might influence adopted strategy?

## Paddle Size and Placement for Defibrillation (ALS-E-030A: ScopRev)

### Rationale for Review

This topic was suggested by the Australian Resuscitation Council. The BLS Task Force was supportive of an updated evidence review because this topic had not been reviewed by ILCOR since 2010.<sup>237,238</sup>

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest
- Intervention: The use of any specific pad size/orientation and position
- Comparator: Standard resuscitation or other specific paddle/pad size/orientation and position
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome. Termination of VF and rates of recurrence of fibrillation/refibrillation were included as important outcomes.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded. It was anticipated that there would be insufficient studies from which to draw a conclusion; case series were included in the initial search and included as long as they contained at least 5 cases.
- Time frame: Since January 1, 2009: All languages were included as long as there was an English

abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to November 11, 2019.

### Summary of Evidence

We did not identify any new evidence that directly addressed this question. See [Appendix B-3](#) for full ScopRev.

### Task Force Insights

Key issues from BLS Task Force discussions were as follows:

Although some studies have shown that anteroposterior electrode placement is more effective than the traditional anterolateral position in elective cardioversion of atrial fibrillation, the majority have failed to demonstrate any clear advantage of any specific electrode position. Transmyocardial current during defibrillation is likely to be maximal when the electrodes are placed so that the area of the heart that is fibrillating lies directly between them (ie, ventricles in VF/pVT, atria in atrial fibrillation). Therefore, the optimal electrode position may not be the same for ventricular and atrial arrhythmias.

Recent approaches including double sequential defibrillation, in which differently oriented sequential defibrillations are delivered, have been evaluated by the Advanced Life Support Task Force in a separate evidence review.

This ScopRev was unable to identify any new studies that needed to be added to the previous SysRev. In light of this, we believe that the existing CoSTR does not need to be modified (with the exception of removing reference to “paddles,” because modern equipment using self-adhesive pads have replaced paddles).

### Treatment Recommendation

These treatment recommendations (below) are unchanged from 2010.<sup>237,238</sup> It is reasonable to place pads on the exposed chest in an anterior-lateral position. An acceptable alternative position is anterior posterior. In large-breasted individuals, it is reasonable to place the left electrode pad lateral to or underneath the left breast, avoiding breast tissue. Consideration should be given to the rapid removal of excessive chest hair before the application of pads, but emphasis must be on minimizing delay in shock delivery.

There is insufficient evidence to recommend a specific electrode size for optimal external defibrillation in adults. However, it is reasonable to use a pad size greater than 8 cm.

## SPECIAL CIRCUMSTANCES

### CPR During Transport (BLS 1509: ScopRev)

#### Rationale for Review

This topic has not been reviewed since before 2005.

#### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with OHCA

- Intervention: Transport to hospital
- Comparator: Completing CPR on scene
- Outcome: Critical: survival with good neurological function (ie, at hospital discharge, 1 month, 6 months, 1 year) and survival (ie, hospital discharge, 1 month, 6 months, 1 year); important: short-term survival (ROSC, hospital admission) and CPR quality parameters (ie, compression fraction rate, depth, leaning, etc)
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded.
- Time frame: All years and all languages were included as long as there was an English abstract.

### Summary of Evidence

This ScopRev is included in [Supplement Appendix B-2](#).

#### Studies Reporting Survival Among OHCA Patients Transported With CPR in Progress (Arriving at Hospital Without a Pulse)

There were 8 nonrandomized studies<sup>239–246</sup> reporting that ROSC was achieved in the emergency department in approximately 9.5% of cases, with 2.9% surviving to hospital discharge.

#### Studies Reporting Quality of Manual CPR on Scene Compared With During Transport

There were 5 nonrandomized studies<sup>247–251</sup> comparing the quality of CPR on scene with the quality of CPR during transport to hospital. Two studies<sup>247,250</sup> concluded that the quality of CPR during transport is no worse than the quality of CPR on scene, whereas 2 studies<sup>249,251</sup> concluded that the quality of CPR was poorer during transport than on scene.

There were 4 RCTs<sup>252–255</sup> and 4 nonrandomized studies<sup>256–259</sup> comparing the quality of CPR on scene with the quality of CPR during transport, using manikins. Manikin studies suggest that CPR quality is poorer during transport than when on scene.

#### Studies Comparing Manual Versus Mechanical CPR During Transport

There were 3 RCTs<sup>260–262</sup> and 3 nonrandomized studies<sup>263–265</sup> reporting survival outcomes for OHCA patients transported with manual CPR compared with mechanical CPR. RCTs showed no benefit from mechanical CPR with respect to ROSC or survival to discharge. The nonrandomized studies reported conflicting results. Two RCTs<sup>260,261</sup> and 3 nonrandomized studies<sup>266–268</sup> suggested variable improvements in physiological parameters with mechanical CPR. Four manikin RCTs<sup>254,255,269,270</sup> and 3 nonrandomized manikin studies<sup>257,271,272</sup> suggested that mechanical CPR provided consistent CPR, whereas the quality of manual CPR declined during transport.



### *Studies Addressing Duration and/or Distance of Transport on Outcomes*

Five nonrandomized studies<sup>246,273–276</sup> suggested that the duration of transport with CPR and the distance transported with CPR does not adversely impact patient outcomes.

There was significant heterogeneity among study populations, study methodologies, outcome measures utilized, and outcomes reported. Findings are grouped into themes, and a narrative analysis is provided.

### **Task Force Insights**

There was considerable task force debate concerning the appropriate outcome for this PICOST:

- Is the quality of CPR during transport better/no different/worse than the quality of CPR on scene?
- Are clinical outcomes affected by the decision to transport with CPR?
- When should the decision to transport with ongoing CPR be made?
- Does the distance of transport affect outcomes of CPR during transport?
- Can we identify which patient groups will/will not benefit from transport with ongoing CPR?
- Should we recommend the use of mechanical CPR during transport?
- What are the risks associated with CPR during transport?

The task force acknowledges several confounding factors when interpreting evidence, such as the use of feedback devices to improve CPR quality during transport and the implementation of high-performance CPR within EMS systems. It was noted that studies of CPR quality reported mean outcome measures and acknowledged that the quality of CPR may fluctuate considerably during transport. Although there is little evidence about risk to providers when performing CPR during transport, there are several reports highlighting the risk of injury when unrestrained in the back of an ambulance. The task force recognizes that performing CPR in the back of a moving ambulance does increase the risk to providers. The decision to transport to hospital or cease in the field might also be dependent on available resources at receiving hospitals—if no additional treatment can be added in the hospital, providers and patients are subjected to additional risk with little potential benefit.

This topic has not been addressed by ILCOR for many years. This ScopRev has identified new evidence addressing this topic. The BLS Task Force recognizes that it may be appropriate to undertake more than 1 SysRev on the basis of these findings. The BLS Task Force will seek public feedback to prioritize the questions to explore in the near future. The BLS Task Force will request as a first priority a SysRev comparing the quality of CPR metrics on scene compared with during transport.

## **Removal of Foreign-Body Airway Obstruction (BLS 368: SysRev)**

### **Rationale for Review**

Foreign-body airway obstruction is a common problem. Many cases are likely resolved easily without the need to involve healthcare providers. Foreign-body airway obstruction is, however, an important cause of early death that typically affects the very young and the elderly or individuals with impaired neurological function/swallowing. Current strategies to relieve foreign-body airway obstruction are well known to many people; delays in treatment increase the risk of death, but interventions themselves can cause harm and death. The topic of relief of foreign-body airway obstruction has not been reviewed since 2010.<sup>1,2</sup> In recent years, manual suction devices (airway clearance devices) that use a vacuum to remove foreign bodies have become commercially available. These devices have not previously been reviewed by ILCOR and are included in this SysRev.

### **Population, Intervention, Comparator, Outcome, Study Design, and Time Frame**

- Population: Adults and children with foreign-body airway obstruction
- Intervention: Interventions to remove foreign-body airway obstruction, such as finger sweep, back slaps or blows, abdominal thrusts, chest thrusts, and suction-based airway clearance devices
- Comparator: No action
- Outcome: Survival with good neurological outcome, survival, ROSC, relief of airway obstruction, harms/complications
- Study design: RCTs, nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies), and case series ( $\geq 5$  cases) were eligible for inclusion. Case reports of injuries/complications were eligible.
- Time frame: All years and all languages were included as long as there was an English abstract. Unpublished studies (eg, conference abstracts, trial protocols), animal studies, manikin studies, and cadaver studies were excluded. The literature was searched to September 2019.
- PROSPERO registration: CRD42019154784

### **Consensus on Science**

The review focused on studies published in the peer-reviewed literature. All studies identified were observational, consisting mostly of case series. The overall certainty of evidence was very low for all outcomes primarily because of very serious risk of bias and imprecision. Key limitations with interpretation of the case series identified include publication bias (reports of successful use or harm are more likely to be published); lack of information about the denominator (ie, the number of times an intervention was used compared with the number of successes

or harms reported); and, in many reports, more than 1 intervention attempted. For these reasons and because of the high degree of heterogeneity across the case reports, no meta-analyses were performed, and individual studies were difficult to interpret. Evidence relating to the use of back blows, abdominal thrusts, chest compressions, and finger sweeps is presented in Table 15.

### Magill Forceps

For the critical outcome of survival with favorable neurological outcome, we identified very-low-certainty evidence from 1 observational study<sup>343</sup> enrolling 240 adults and children with OHCA with foreign-body airway obstruction, which showed benefit associated with the use of Magill forceps by EMS personnel compared with no use (OR, 3.96 [95% CI, 1.21–13.00]; 107 more patients/1000 survived with the intervention [95% CI, 8 more patients/1000 to 324 more patients/1000 survived with the intervention]). This outcome was achieved despite the much lower incidence of bystander CPR provided to the Magill forceps group.

For the critical outcome of survival, we identified very-low-certainty evidence from 1 observational study<sup>343</sup> enrolling 240 patients with OHCA associated with foreign-body airway obstruction. The rate of survival with EMS use of Magill forceps was 27% versus 17% in the control group ( $P=0.086$ ) despite a lower rate of bystander CPR before EMS arrival (57% versus 80%;  $P<0.001$ ).

For the important outcome of relief of foreign-body airway obstruction, we identified very-low-certainty

evidence from 4 case series studies<sup>278,285,343,344</sup> reporting successful relief of foreign-body airway obstruction in 417 patients treated with Magill forceps.

### Airway Clearance Devices

For the critical outcome of survival and the important outcome of relief of foreign-body airway obstruction, we identified a single observational study with very-low-certainty evidence reporting about 9 adult patients with foreign-body airway obstruction who survived after treatment with a suction-based airway clearance device.<sup>345</sup>

### Foreign-Body Airway Obstruction Removal by Bystanders

For the critical outcome of survival with good neurological outcome, we identified very-low-certainty evidence downgraded for very serious risk of bias from 1 observational study<sup>278</sup> enrolling 41 patients with foreign-body airway obstruction, which showed benefit from bystander attempts to remove the foreign-body airway obstruction compared with no bystander attempts (intervention versus control, 74% versus 32%;  $P=0.0075$ ).

### Treatment Recommendations

We suggest that back slaps are used initially in adults and children with a foreign-body airway obstruction and an ineffective cough (weak recommendation, very-low-certainty evidence).

We suggest that abdominal thrusts are used in adults and children (older than 1 year) with a foreign-body airway obstruction and an ineffective cough when back

**Table 15. Removal of Foreign Body Airway Obstruction**

| Intervention               | Outcome               | Studies                                      | No. of Patients | Results  |
|----------------------------|-----------------------|--|-----------------|--|
| Back blows                 | Survival              | 1 observational <sup>277</sup>               | 13              | All 13 patients survived   |
|                            | Relief of obstruction | 3 observational <sup>277–279</sup>           | 75              | All 75 patients had relief of obstruction  |
|                            | Injury/harm           | 4 observational <sup>280–282a</sup>          | 4               | 3 vascular injuries, 1 thoracic injury   |
| Abdominal thrusts          | Survival              | 2 observational <sup>283,284</sup>           | 189             | All 189 patients survived  |
|                            | Relief of obstruction | 6 observational <sup>277–279,283–285</sup>   | 417             | All 417 patients had relief of obstruction   |
|                            | Injury/harm           | 49 observational <sup>281,282a,286–333</sup> | 52              | 17 gastric/esophageal injuries, 15 vascular injuries, 12 thoracic injuries, 8 abdominal injuries |
| Chest thrusts/compressions | Survival              | 1 observational <sup>334</sup>               | 138             | All 138 patients survived  |
|                            | Relief of obstruction | 1 observational <sup>279</sup>               | 28              | All 28 patients had relief of obstruction  |
|                            | Injury/harm           | 4 observational <sup>280,312,323,326</sup>   | 5               | 3 gastric/esophageal injuries, 2 vascular injuries.  |
| Finger sweep               | Survival              | 1 observational <sup>277</sup>               | 6               | All 6 patients survived  |
|                            | Relief of obstruction | 2 observational <sup>277,279</sup>           | 36              | All 36 patients had relief of obstruction  |
|                            | Injury/harm           | 8 observational <sup>335–342</sup>           | 10              | 5 dislodgement of object, 5 injury to nasopharynx  |

slaps are ineffective (weak recommendation, very-low-certainty evidence).

We suggest that rescuers consider the manual extraction of visible items in the mouth (weak recommendation, very-low-certainty evidence).

We suggest against the use of blind finger sweeps in patients with a foreign-body airway obstruction (weak recommendation, very-low-certainty evidence).

We suggest that appropriately skilled healthcare providers use Magill forceps to remove a foreign-body airway obstruction in patients with OHCA from foreign-body airway obstruction (weak recommendation, very-low-certainty evidence).

We suggest that chest thrusts be used in unconscious adults and children with a foreign-body airway obstruction (weak recommendation, very-low-certainty evidence).

We suggest that bystanders undertake interventions to support foreign-body airway obstruction removal as soon as possible after recognition (weak recommendation, very-low-certainty evidence).

We suggest against the routine use of suction-based airway clearance devices (weak recommendation, very-low-certainty evidence).

### **Justification and Evidence-to-Decision Framework Highlights**

The evidence-to-decision table is included in [Supplement Appendix A-13](#). The current treatment recommendations are similar to previous recommendations, but the BLS Task Force has provided some additional guidance about the recommended sequence of steps to relieve airway obstruction. The task force recognizes the importance of early removal of a foreign-body airway obstruction to prevent cardiac arrest. Bystanders should be encouraged to assist victims by rapidly attempting to remove the obstruction. The initial response to foreign-body airway obstruction in a conscious individual should be to encourage coughing because this is a normal physiological response that may be effective and is unlikely to cause harm. The sequence of interventions in individuals without an effective cough suggested in treatment recommendations seeks to balance the benefits of early removal of the foreign-body airway obstruction with the potential harms of interventions, such as abdominal thrusts.

We prioritized consistency with current treatment recommendations. We note the difference in methodologic approaches used in this review compared with previous reviews. In particular, previous reviews included cadaver, animal, and manikin studies.

We note that evidence for all outcomes is assessed as very low certainty. Research on foreign-body airway obstruction is challenging because many with a foreign-body airway obstruction are treated immediately and effectively by bystanders or by coughing. It would be difficult if not impossible to perform an RCT of treatments for foreign-body airway obstruction.

The task force distinguished between a situation in which a foreign-body airway obstruction can be seen in the mouth and a situation in which no object can be seen. When an object can be seen in the mouth, the manual removal of the item was considered appropriate. When an object cannot be seen in the mouth, the potential harm associated with the rescuer placing and moving their fingers in the victim's mouth (a blind finger sweep) and the lack of clear benefit to this approach led to a suggestion against the use of blind finger sweeps.

The task force treatment recommendation limits use of abdominal thrusts to adults and children beyond infancy. This was driven by concerns that, in infants, the limited protection of the upper abdominal organs by the lower ribs may mean that the potential harm of abdominal thrusts outweighs any potential benefit. This is consistent with previous treatment recommendations.

The task force treatment recommendation supporting the use of chest thrusts/compressions is based on case series reports of successful relief of foreign-body airway obstruction (unknown whether patients were in cardiac arrest) and an observational study that found that chest compressions improved neurologically intact survival in unresponsive patients with foreign-body airway obstruction. Our current recommendation is consistent with previous treatment recommendations.

The introduction of a treatment recommendation supporting the use of Magill forceps by suitably trained healthcare providers reflects the potential benefit of the intervention and the availability of relevant equipment to trained individuals. The task force expects that these trained healthcare providers will already be skilled in advanced airway management. The treatment recommendation is based on evidence from case series of successful relief in victims with foreign-body airway obstruction (unknown whether patients were in cardiac arrest) and an observational study that found that EMS use of Magill forceps was associated with improved neurologically intact survival in those with OHCA from foreign-body airway obstruction.

The task force acknowledges that there are very limited data in the peer reviewed literature assessing the efficacy of suction-based airway clearance devices (a case series of 9 adults). The task force agreed that the peer-reviewed published data were insufficient to support the implementation of a new technology with an associated financial and training cost. The task force has outlined recommendations for further research in relation to these devices.

We identified no evidence that specifically examined foreign-body airway obstruction removal in pregnant individuals. The task force suggests that abdominal thrusts are avoided in this group due to risk of injury to the fetus.

### **Knowledge Gaps**

- There is a need for high-quality observational studies that accurately describe the incidence of foreign-body

airway obstruction, patient demographics (age, setting, comorbidities, food type, level of consciousness), full range of interventions delivered, who delivered interventions (health professional/lay responder), success rates of interventions, harm of interventions, and outcomes. It is unlikely that such a study can be conducted using only health service data.

- There is a need for further evidence on the benefits and harms of suction-based airway clearance devices. The task force encourages the registration of all device uses. Reports should detail key demographics (eg, age, setting, comorbidities, food type, level of consciousness), full range of interventions provided, who provided the intervention (lay compared with healthcare professional), and outcomes. This evidence initially may come in the form of published case series.

## Resuscitation Care for Suspected Opioid-Associated Emergencies (BLS 811: SysRev)

### Rationale for Review

Deaths from drug overdose are an increasing public health burden in many countries. In the United States alone, more than 70 000 deaths were attributed to drug overdose in 2017.<sup>345a</sup> Overdose deaths have been increasing since 2013; although there is increasing research into overdose prevention and response education, there is a need for a SysRev to guide development of best practice guidelines for bystander resuscitation in suspected opioid-induced emergencies.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children with suspected opioid-associated cardiorespiratory arrest in the prehospital setting
- Intervention: Bystander naloxone administration (intramuscular or intranasal) in addition to standard CPR
- Comparator: Conventional CPR only
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, and controlled before-and-after studies, cohort studies) were eligible for inclusion.
- Time frame: All years and all languages were included as long as there was an English abstract. Unpublished studies (eg, conference abstracts, trial protocols), animal studies, manikin studies, and cadaver studies were excluded. The literature was searched to October 2019.

### Consensus on Science

We did not identify any studies reporting any critical or important outcomes for adults or children with suspected opioid-associated cardiorespiratory arrest in any setting, comparing bystander-administered naloxone

(intramuscular or intranasal) plus conventional CPR with conventional CPR only.

### Treatment Recommendation

We suggest that CPR be started without delay in any unconscious person not breathing normally and that naloxone be used by lay rescuers in suspected opioid-related respiratory or circulatory arrest (weak recommendation based on expert consensus).

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-14](#). There is no direct evidence comparing outcomes for patients with opioid-induced respiratory or cardiac arrest treated with naloxone in addition to standard CPR compared with those treated with CPR alone. Despite this, the BLS Task Force decided to make a suggestion for the use of naloxone on the basis of expert opinion alone, wanting to underline the importance and challenge of the opioid epidemic. Although administering naloxone is unlikely to directly harm the patient, rescuers should be prepared for behavioral changes that may occur after drug administration. Patients who are resuscitated from a narcotic overdose may become agitated and sometimes violent.

Although no evidence directly evaluating the clinical question was identified, we did identify a summary of 4 case series including 66 patients, in which 39 of 39 patients who received naloxone after opioid overdose recovered compared with 24 of 27 who did not receive naloxone after opioid overdose.<sup>346</sup> At the population level, there is evidence to demonstrate improved outcomes in communities after implementation of various naloxone distribution schemes. A recent SysRev identified 22 observational studies evaluating the effect of overdose education and naloxone distribution using Bradford Hill criteria and found a link between implementation of these programs and decreased mortality rates.<sup>347</sup>

Diagnosis of respiratory or cardiac arrest is not always straightforward, and lay rescuers would be expected to have a high suspicion of cardiac or respiratory arrest in any unresponsive person with suspected drug overdose. Administration of naloxone is likely to have preventive effects if given after a drug overdose that has not yet caused respiratory or cardiac arrest, and the potential for desirable effects in a broader population strengthens the suggestion to administer naloxone in this setting. Furthermore, there are very few reports of side effects from naloxone.<sup>348</sup> Although it is possible that bystanders might spend valuable time finding and administering naloxone instead of starting CPR during respiratory or cardiac arrest, lack of reports of harm from large-scale implementation of naloxone distribution schemes indicate that this is unlikely a big problem.

Because there is no formal evaluation of naloxone with CPR compared with CPR alone in opioid overdose, it



is not possible to formally balance desirable and undesirable effects of naloxone administration by laypeople. As a response to the growing epidemic, naloxone has been widely distributed by healthcare authorities to laypeople in various opioid overdose prevention schemes. Overall, these programs report beneficial outcomes at the population level. The BLS Task Force therefore considers it very likely that the desirable effects outweigh undesirable effects and that use of naloxone is acceptable by key stakeholders as well as the general population.

### Knowledge Gaps

Current knowledge gaps include but are not limited to the following:

- There is currently no evidence evaluating the role of naloxone use among bystanders attempting CPR in suspected opioid-related respiratory or circulatory arrest.
- Further research is needed to determine the optimal components of resuscitation and the role of naloxone during bystander CPR.

## Drowning (BLS 856: SysRev)

### Rationale for Review

This question was initiated in response to a request that ILCOR review the evidence for prognostic factors that predict outcome in relation to a drowning incident. Drowning was last reviewed in 2015.<sup>3,4</sup> Drowning is the third leading cause of unintentional injury death worldwide, accounting for over 360 000 deaths annually.<sup>349</sup> Care of a submersion victim in high-resource countries often involves a multiagency approach, with several different organizations independently responsible for different phases of the victim's care, beginning with initial aquatic rescue, through on-scene resuscitation and transfer to hospital, and with in-hospital and rehabilitative care. Attempting to rescue a submerged victim has substantial resource implications and may place rescuers at risk.

### Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children who are submerged in water
- Intervention: Any particular factor in search-and-rescue operations (eg, duration of submersion, salinity of water, water temperature, age of victim)
- Comparator: Compared with no factors
- Outcome: Survival to hospital discharge with good neurological outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. It was anticipated that there would be insufficient studies from which to draw a

conclusion; case series were included in the initial search as long as they contained at least 5 cases.

- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 1, 2019.

### Consensus on Science

#### Age

For the critical outcome of survival with favorable neurological outcome, we identified very-low-certainty evidence from 11 observational studies (downgraded for bias inconsistency, indirectness, and imprecision) comprising 4 105 patients.<sup>350–359,359a</sup> Of the 8 pediatric studies, 6 found that young age, variably defined as less than 3, 4, 5, or 6 years, was not associated with favorable neurological outcome.<sup>350–354,356</sup> A single pediatric study including 166 children less than 15 years of age reported better outcomes in children age less than 5 years (RR, 0.12; 95% CI, 0.03–0.44).<sup>355</sup> Four studies considered drowning victims of all ages; 2 found no association between age and outcome.<sup>357,358</sup> One reported worse outcomes associated with children aged greater than 5 years (RR, 0.66; 95% CI, 0.51–0.85).<sup>359</sup>

For the critical outcome of survival, we identified very-low-certainty evidence (downgraded for risk of bias, inconsistency, indirectness, and imprecision) from 6 observational studies including 1313 patients.<sup>360–365</sup> Three studies found that age was not associated with outcome.<sup>361,363,365</sup> Two reported better outcomes associated with younger ages (less than 58 years: RR, 0.27; 95% CI, 0.08–0.96<sup>362</sup>; less than 46 years: RR, 0.98; 95% CI, 0.99–0.99),<sup>364</sup> and 1 favored older age ( $\geq 3$  years: RR, 1.51; 95% CI, 1.19–1.9).<sup>360</sup>

#### EMS Response Interval

For the critical outcome of survival, we identified very-low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision) from 2 observational studies including 746 patients in the Swedish EMS OHCA registry.<sup>362,366</sup> EMS response intervals of less than 10 minutes were associated with better survival (RR, 0.29; 95% CI, 0.13–0.66)<sup>366</sup> and a reported OR of 0.44 (95% CI, 0.06–0.83).<sup>362</sup>

#### Salinity

For the critical outcome of survival with favorable neurological outcome, we identified very-low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision) from 6 observational studies<sup>354,357,359,359a,367,368</sup> 1799 including 3 584 drowning victims, of which 980 occurred in salt water and 2 604 in fresh water. Two reported that drowning in salt water was associated with better outcomes (RRs, 1.3 [95% CI, 1.12–1.5]<sup>357</sup> and 1.2 [95% CI, 1.1–1.4],<sup>354</sup> and 4 found no association between water salinity

and outcome (RRs, 1.1 [95% CI, 0.95–1.2],<sup>367</sup> 1.14 [95% CI, 0.9–1.4],<sup>359</sup> 1.1 [95% CI, 0.70–1.72],<sup>368</sup> and 1.15 [95% CI, 0.91–1.45].<sup>359a</sup>

For the critical outcome of survival, we identified very-low-certainty evidence (downgraded for risk of bias imprecision, inconsistency, indirectness, and imprecision) from 5 observational studies.<sup>360,363,368–370</sup> One reported better outcomes associated with salt water submersion (RR, 1.34; 95% CI, 1.19–1.52),<sup>369</sup> 3 showed no association between water salinity and survival (RRs, 1.22 [95% CI, 0.95–1.56],<sup>360</sup> 0.88 [95% CI, 0.40–1.92],<sup>368</sup> and 0.94 [95% CI, 0.62–1.4],<sup>370</sup> and 1 reported worse survival associated with salt water drowning (RR, 0.18; 95% CI, 0.03–1.43).<sup>363</sup>

### **Submersion Duration**

For the purposes of this review, we considered studies in 3 groups. We defined those with short submersion duration (less than 5–6 minutes), those with intermediate duration (less than 10 minutes), and those with prolonged submersion duration (less than 15–25 minutes).

#### **Short Submersion Intervals (Less Than 5–6 Minutes)**

For the critical outcome of survival with favorable neurological outcome, we identified moderate-certainty evidence from 15 observational studies (downgraded for bias and indirectness, upgraded for dose response) including 2 746 drowning victims.<sup>350,352–356,359,359a,371–377</sup> All studies noted worse outcomes associated with submersion durations exceeding 5 minutes (RRs between 0.05<sup>359</sup> and 0.61.<sup>355</sup> The 943/1 075 patients (87.7%) who had outcome information available and were submerged for short durations had good outcomes compared with the 139/1 238 (11.2%) who had longer submersion durations.

For the critical outcome of survival, we identified low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision; upgraded for dose response) from 6 observational studies comprising 392 cases.<sup>360,361,369,375,378,379</sup> All studies noted an association between worse outcomes with prolonged compared with short submersion durations (RRs between 0.27<sup>378</sup> and 0.83.<sup>379</sup> The 204/217 patients (94.0%) submerged for short durations had good outcomes compared with the 54/98 (55.1%) who had longer submersion durations.

#### **Intermediate Submersion Intervals (Less Than 10 Minutes)**

For the critical outcome of survival with favorable neurological outcome, we identified moderate-certainty evidence (downgraded for bias, indirectness, and imprecision; upgraded for dose response) from 9 observational studies including 2 453 victims of drowning.<sup>352,354,355,359,371,372,374,380,381</sup> All studies noted an association between worse outcomes and prolonged submersion durations compared with intermediate submersion durations (RRs between 0.02.<sup>359</sup> and 0.45.<sup>355,372</sup> The 787/1 019 patients (77.2%) submerged

for intermediate durations had good outcomes compared with the 36/962 (3.7%) who had longer submersion durations.

For the critical outcome of survival, we identified low-certainty evidence (downgraded for bias, indirectness and imprecision; upgraded for dose response) from 2 observational studies<sup>369,382</sup> reporting about 121 victims of drowning. In the first study,<sup>369</sup> 56/73 (77%) submerged for less than 10 minutes survived compared with none of the 7 patients who were submerged for more prolonged periods survived (RR, not estimable; absolute difference, 76.7%; 95% CI, 39.7%–94.9%). The second study<sup>382</sup> also noted better survival rates associated with a submersion duration of less than 10 minutes (46/50 [96%] survived) compared with submersion duration of more than 10 minutes (2/5 [40%] survived).<sup>382</sup>

#### **Prolonged Submersion Intervals (Less Than 15–25 Minutes)**

For the critical outcome of survival with favorable neurological outcome, we identified low-certainty evidence (downgraded for bias and imprecision, upgraded for dose response) from 3 observational studies including reports of 739 victims of drowning.<sup>352,354,374</sup> In 1 study (n=398),<sup>354</sup> submersion for less than 20 minutes was associated with better outcomes (289/370 [78%] compared with 1/27 [4%] survived; RR, 0.05; 95% CI, 0.01–0.31). The second series<sup>352</sup> reported better outcomes associated with a submersion duration of less than 25 minutes (68/101 survivors, or a 67% survival rate) compared with a submersion duration longer than 25 minutes (0/4 survivors, or a 0% survival rate).<sup>352</sup> In the third study, which included hypothermic children in cardiac arrest, 12/66 (18%) submerged for less than 25 minutes survived compared with 0/39 who were submerged for more than 25 minutes.<sup>374</sup>

For the critical outcome of survival, we identified very-low-certainty evidence (downgraded for bias, indirectness, and imprecision) from a single study<sup>378</sup> comprising 49 patients. Those with a submersion duration of less than 15 minutes had an overall survival rate of 82% (33/39) compared with none of the 2 victims whose submersion duration exceeded 15 minutes (RR, not estimable; absolute difference, 84.6%; 95% CI, 17.3%–92.8%).

### **Water Temperature**

For the critical outcome of survival with favorable neurological outcome, we identified very-low-certainty evidence (downgraded for bias, inconsistency, indirectness, and imprecision) from 2 studies<sup>359,374</sup> including 1254 victims of drowning. The largest study (n=1094) included all unintentional drownings in open waters (lakes, ponds, rivers, ocean) in a single large region, collected from medical examiners, EMS systems, and all regional hospitals.<sup>359</sup> Water temperatures were measured within a month of the drowning incident. Univariable analysis according to temperatures less than or greater than 6°C or less than or

greater than 16°C did not find any association between water temperature and neurological survival. Multivariable analysis also showed no association between water temperature and outcome. The second study included 160 children who required resuscitation and were hypothermic after submersion.<sup>374</sup> Water temperatures were estimated on the basis of the season. Submersion in the winter, with water temperature estimated as 0°C to 8°C, was associated with better outcomes than submersion in spring or summer, with water temperature estimated at 6°C to 28°C (univariable OR, 4.55; 95% CI, 1.37–15.09).

For the critical outcome of survival, we identified very-low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision) from a single study<sup>362</sup> including 250 victims of drowning. This study included only those who had OHCA and received EMS care, and it included those with intentional (suicide and homicide) drowning. This study found no relationship between water temperature less than or greater than 15°C and outcome (RR, 0.94; 95% CI, 0.34–2.62; absolute difference, 0.36%; 95% CI, –6.4% to 6.5%).

### Witnessed Status

The definition of witnessed compared with unwitnessed drowning was inconsistently defined in the studies reviewed. It was often unclear if the term “witnessed” related to the submersion or the cardiac arrest.

For the critical outcome of survival with favorable neurological outcome, we found very-low-certainty evidence (downgraded for indirectness and imprecision) from 3 observational studies<sup>358,359a,383</sup> involving 2 707 patients. Two studies reported better neurological outcomes associated with a witness to the event (unadjusted odds ratio, 16.33 [95% CI, 5.58–47.77]; AOR, 11.8 [95% CI, 2.84–49.08]<sup>358</sup>; and unadjusted odds ratio, 2.6 [95% CI, 1.69–4.01]; AOR, 3.27 [95% CI, 2.0–5.36]<sup>383</sup>). Neither of the analyses included submersion duration, which several studies have reported as an independent predictor.

For the critical outcome of survival, we found low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision) from 4 studies<sup>358,363,364,366</sup> involving 2 857 victims. Two studies<sup>362,364</sup> were from the same EMS system, and both used multivariable analysis. The smaller study (n=255) showed that witnessed status was not associated with improved survival (RR, 0.55; 95% CI, 0.17–1.75; absolute difference, 3%; 95% CI, –3.1% to 11.2%).<sup>362</sup> However, in the larger subsequent study from that same EMS system, witnessed status predicted better outcome (reported univariable analysis:  $P=0.05$ ; AOR, 2.5; 95% CI, 1.38–4.52).<sup>364</sup> Another study<sup>363</sup> found no association between witnessed status and improved survival (RR, 0.82; 95% CI, 0.26–2.59). A large observational study from Japan<sup>358</sup> reported an UAOR of 7.38 (95% CI, 3.81–14.3) and an AOR of 6.5 (95% CI, 2.81–15.02) with witnessed compared with unwitnessed drowning, although the unusual population of much older victims, most drowning

in bathtubs, and a very low rate of favorable outcomes limited the generalizability of these findings.

### Treatment Recommendations

These treatment recommendations are unchanged from 2015.<sup>3,4</sup> We recommend that submersion duration be used as a prognostic indicator when making decisions surrounding search and rescue resource management/operations (strong recommendation, moderate-certainty evidence).

We suggest against the use of age, EMS response time, water type (fresh or salt), water temperature, and witness status when making prognostic decisions (weak recommendation, very-low-certainty evidence).

We acknowledge that this review excluded exceptional and rare case reports that identify good outcomes after prolonged submersion in icy water.

### Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-15](#). The 2015 CoSTR benefited from significant feedback from ILCOR task forces as well as through public consultation and input from the drowning research and clinical communities.<sup>3,4</sup> In making the original recommendations, the task force placed priority on producing simple guidance that may assist those responsible for managing search and rescue operations. The public comments highlighted the difficult moral dilemmas facing the rescuer in an emotionally charged and fast-moving environment requiring dynamic risk assessments that consider the likelihood of a favorable outcome with the risks posed to those undertaking the rescue. It must also be noted that there is substantial difficulty inherent in determining the submersion duration and the bias of studies using it as a predictive variable. The key finding of the 2015 review was that submersion durations of less than 10 minutes are associated with a very high chance of favorable outcome, and submersion durations more than 25 minutes are associated with a low chance of favorable outcomes.<sup>3,4</sup>

The findings from the 6 new papers identified in this update<sup>359a,368,370,375,376,383</sup> are consistent with the 2015 treatment recommendation.<sup>3,4</sup> The previously identified limitations of this review (exclusion of factors after the victim is rescued, for example, bystander CPR<sup>383–385</sup>; specialist interventions, such as the use of extracorporeal membrane oxygenation<sup>386–393</sup>; and the lack of prospective validation of submersion duration as a clinical decision rule) persist. Similarly, continued reports of rare survival after prolonged (more than 30 minutes) submersion<sup>387,392,394</sup> highlight the need for case-by-case decisions that balance risk and potential for benefit.

### Knowledge Gaps

Submersion duration should be assessed in all future drowning studies and be part of multivariable analyses.



To better clarify the value of this predictor, studies should include all victims rescued from the water and not only subcategories.

## POTENTIAL HARM FROM CPR

### Harm From CPR to Victims Not in Cardiac Arrest (BLS 353: SysRev)

#### *Rationale for Review*

Many lay rescuers are reluctant to begin CPR even when a victim is in cardiac arrest because of concern that delivering chest compressions to a person who is not in cardiac arrest could cause serious harm. Case reports and case series of serious harm to persons receiving CPR who are not in cardiac arrest are likely to be published because they are of general interest to a broad group of healthcare providers. A lack of reported cases demonstrating serious harm could strengthen arguments that desirable effects will far outweigh undesirable effects. This topic was last reviewed in 2015.<sup>3,4</sup>

#### *Population, Intervention, Comparator, Outcome, Study Design, and Time Frame*

- Population: Adults and children without OHCA
- Intervention: Provision of chest compressions from lay rescuers
- Comparator: No use of chest compressions
- Outcome: Change in survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; harm (eg, rib fracture); complications; major bleeding; risk of complications (eg, aspiration); survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival to admission
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded. It was anticipated that there would be insufficient studies from which to draw a conclusion; case series and case reports were included in the initial search.
- Time frame: All years and all languages were included as long as there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 13, 2019.

#### *Consensus on Science*

For the important outcome of harm, we identified very-low-certainty evidence (downgraded for risk of bias and imprecision) from 4 observational studies enrolling 762 patients who were not in cardiac arrest but received CPR by lay rescuers out-of-hospital.

Three of the studies<sup>395–397</sup> reviewed the medical records to identify harm, and 1 included follow-up telephone interviews.<sup>395</sup> Pooled data from the first 3 studies, encompassing 345 patients, found an incidence of rhabdomyolysis of 0.3% (n=1), bone fracture (ribs and clavicle) of 1.7% (95% CI, 0.4%–3.1%), pain in the area of chest compression of 8.7% (95% CI, 5.7%–11.7%), and no clinically relevant visceral injury. The fourth study<sup>34</sup> relied on fire department observations at the scene; there were no reported injuries in 417 patients.

#### *Treatment Recommendation*

This treatment recommendation is unchanged from 2015.<sup>3,4</sup> We recommend that lay people initiate CPR for presumed cardiac arrest without concerns of harm to patients not in cardiac arrest (strong recommendation, very-low-certainty evidence).

#### *Justification and Evidence-to-Decision Framework Highlights*

The evidence-to-decision table is included in [Supplement Appendix A-16](#). No change was made to this treatment recommendation. In continuing to make this discordant recommendation (strong recommendation based on very-low-certainty evidence), the BLS Task Force placed a much higher value on the potential survival benefits of CPR initiated by lay persons for patients in cardiac arrest and a lower value on the low risk of injury to patients not in cardiac arrest. The intention of this recommendation is to strongly encourage and support lay rescuers who are willing to initiate CPR in any setting when they believe someone is in cardiac arrest. The intention is also to support emergency medical dispatchers in their efforts to provide DA-CPR instructions in suspected cardiac arrest calls.

#### *Knowledge Gaps*

- Studies are needed to identify harm and provide follow-up after hospital discharge. Many of the conditions prompting initiation of CPR for persons not in cardiac arrest are associated with reduced responsiveness and have poor prognoses. Whether chest compressions and rescue breaths could accentuate these conditions independent of physical injury is not known at the present time.
- The incidence of chest wall fractures was substantially lower than the incidence reported after CPR in patients who were in cardiac arrest. This is likely the result of a shorter duration of CPR (approximately 6 minutes) initiated by lay persons but stopped by professional rescuers and the younger patient age in the studies reviewed. However, it is possible that the lack of systematic follow-up leads to under-reporting of these injuries, and additional research is warranted.



- Could the accuracy of DA protocol be enhanced to reduce the frequency of CPR performed on patients not in cardiac arrest without compromising the initiation of CPR on patients in cardiac arrest?

## Harm to Rescuers From CPR (BLS 354: ScopRev)

### Rationale for Review

The BLS Task Force prioritized an updated evidence review because this topic had not been reviewed by IL-COR since 2010, and that review addressed only injury from CPR to victims who are not in cardiac arrest.<sup>1,2</sup> This 2020 review focused on any potential harm to the rescuers during CPR, including harm during chest compressions, during mouth-to-mouth ventilation, and with the use of defibrillators.

### Summary of Evidence

The complete ScopRev is included in [Supplement Appendix B-5](#). The review identified 5 experimental studies and 1 case report published since 2008. The 5 experimental studies reported the perception of rescuers in an experimental setting during shock administration for elective cardioversion. In these studies, the authors also measured current flow and the average leakage current in different experiments.

### Task Force Insights

We identified many gaps in the published literature. No RCTs were identified that met our inclusion criteria. Most identified studies addressed safety of shock delivery during chest compressions when rescuers wore gloves.

Despite limited evidence evaluating rescuer safety, there was broad agreement within the BLS Task Force that the lack of published evidence supports the interpretation that CPR is generally safe for rescuers. A few reports demonstrate the possibility of disease transmission in the course of performing mouth-to-mouth ventilation. The isolated reports of adverse effects resulting from the widespread and frequent use of CPR suggest that performing CPR is relatively safe. Delivery of a defibrillator shock with an AED during BLS is also safe. The incidence and morbidity of defibrillator-related injuries in the rescuers are low.

The BLS Task Force considers the overall body of new evidence identified by this ScopRev insufficient to warrant a full SysRev. The few reports of harm to rescuers from performing CPR and defibrillation are supportive of general recommendations that lay rescuers may safely perform CPR and use an AED.

### Treatment Recommendation

Evidence supporting rescuer safety during CPR is limited. The few isolated reports of adverse effects resulting from the widespread and frequent use of CPR suggest that performing CPR is relatively safe. Delivery of a defibrillator shock with an AED during BLS is also safe. The incidence and morbidity of defibrillator-related injuries in the rescuers are low.

## TOPICS NOT REVIEWED IN 2020

Topics not reviewed or updated are the following:

- BLS 352: Passive ventilation technique
- BLS 358: Minimizing pauses in chest compressions

## ARTICLE INFORMATION

The American Heart Association requests that this document be cited as follows: Olasveengen TM, Mancini ME, Perkins GD, Avis S, Brooks S, Castrén M, Chung SP, Considine J, Couper K, Escalante R, Hatanaka T, Hung KKC, Kudenchuk P, Lim SH, Nishiyama C, Ristagno G, Semeraro F, Smith CM, Smyth MA, Vaillancourt C, Nolan JP, Hazinski MF, Morley PT; on behalf of the Adult Basic Life Support Collaborators. Adult basic life support: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(suppl 1):S41–S91. doi: 10.1161/CIR.0000000000000892

Supplemental materials are available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/CIR.0000000000000892>

This article has been copublished in *Resuscitation*. Published by Elsevier Ireland Ltd. All rights reserved.

### Authors

Theresa M. Olasveengen, MD, PhD; Mary E. Mancini, RN, PhD; Gavin D. Perkins, MB, ChB, MMed, MD; Suzanne Avis, B App Sc, MPH; Steven Brooks, MD, MHSc; Maaret Castrén, MD, PhD; Sung Phil Chung, MD, PhD; Julie Considine, RN, PhD; Keith Couper, RN, PhD; Raffo Escalante, MD; Tetsuo Hatanaka, MD, PhD; Kevin K.C. Hung, MBChB, MPH; Peter Kudenchuk, MD; Swee Han Lim, MBBS; Chika Nishiyama, RN, DrPH; Giuseppe Ristagno, MD, PhD; Federico Semeraro, MD; Christopher M. Smith, MD, MSc; Michael A. Smyth, BSc(hons), MSc, PhD; Christian Vaillancourt, MD, MSc; Jerry P. Nolan, MBChB; Mary Fran Hazinski, RN, MSN; and Peter T. Morley, MBBS; on behalf of the Adult Basic Life Support Collaborators

### Acknowledgments

The authors thank the following individuals (the Adult Basic Life Support Collaborators) for their contributions: Hildigunnur Svavarsdóttir, Violetta Raffay, Artem Kuzovlev, Jan-Thorsten Grasner, Ryan Dee, Michael Smith, and Kausala Rajendran.

## Disclosures

### Appendix 1. Writing Group Disclosures

| Writing Group Member   | Employment  | Research Grant  | Other Research Support | Speakers' Bureau/ Honoraria | Expert Witness         | Ownership Interest | Consultant/ Advisory Board   | Other  |
|------------------------|---|---|------------------------|-----------------------------|------------------------|--------------------|--|--|
| Theresa M. Olasveengen | Oslo University Hospital (Norway)   | Zoll Foundation*; Laerdal Foundation*   | None                   | None                        | None                   | None               | None   | None   |
| Mary E. Mancini        | University of Texas at Arlington  | None  | None                   | Stryker*                    | None                   | None               | None   | None   |
| Gavin D. Perkins       | Warwick Medical School and University Hospitals NHS Foundation Trust (United Kingdom) | National Institute for Health Research (Institutional funding relating to cardiac arrest research)†; British Heart Foundation (Institutional funding relating to cardiac arrest research)†; Resuscitation Council UK (Institutional funding relating to cardiac arrest research)† | None                   | None                        | None                   | None               | None   | None   |
| Suzanne Avis           | Ms School of Medicine (Australia)   | None  | None                   | None                        | None                   | None               | None   | None   |
| Steven Brooks          | Queen's University, Kingston General Hospital   | Canadian Institutes of Health Research (Peer-reviewed funding to study the PulsePoint mobile device application)†   | None                   | None                        | Borden-Ladner-Gervais† | None               | Heart and Stroke Foundation of Canada (member of the Resuscitation Advisory Committee and receive travel reimbursement)* | None   |
| Maaret Castrén         | Helsinki University Hospital (Finland)  | None  | None                   | None                        | None                   | None               | None   | None   |
| Sung Phil Chung        | Yonsei University (Republic of Korea)   | None  | None                   | None                        | None                   | None               | None   | None   |
| Julie Considine        | Deakin University   | None  | None                   | None                        | None                   | None               | None   | Salary: Deakin University†; Eastern Health†; College of Emergency Nursing Australasia* |
| Keith Couper           | University of Warwick, Warwick Medical School (United Kingdom)                        | None  | None                   | None                        | None                   | None               | None   | None   |
| Raffo Escalante        | Inter-American Heart Foundation (Peru)  | None  | None                   | None                        | None                   | None               | None   | None   |
| Tetsuo Hatanaka        | Emergency Life Saving Technique Academy (Japan)                                       | None  | None                   | None                        | None                   | None               | None   | None   |
| Mary Fran Hazinski     | Vanderbilt University   | None  | None                   | None                        | None                   | None               | American Heart Association†  | None   |
| Kevin K.C. Hung        | Chinese University of Hong Kong (Hong Kong)   | None  | None                   | None                        | None                   | None               | None   | None   |

(Continued)

## Appendix 1. Continued

| Writing Group Member   | Employment  | Research Grant   | Other Research Support | Speakers' Bureau/Honoraria | Expert Witness | Ownership Interest | Consultant/Advisory Board | Other |
|------------------------|---|--|------------------------|----------------------------|----------------|--------------------|---------------------------|-------|
| Peter Kudenchuk        | University of Washington  | NIH/NINDS (Principal Investigator for SIREN Network at University of Washington)†  | None                   | None                       | None           | None               | None                      | None  |
| Swee Han Lim           | Singapore General Hospital (Singapore)                            | None   | None                   | None                       | None           | None               | None                      | None  |
| Peter T. Morley        | University of Melbourne, Royal Melbourne Hospital (Australia)     | None   | None                   | None                       | None           | None               | None                      | None  |
| Chika Nishiyama        | Kyoto University (Japan)  | ZOLL Medical Corporation†  | None                   | None                       | None           | None               | None                      | None  |
| Jerry P. Nolan         | Warwick Medical School (United Kingdom)                           | None   | None                   | None                       | None           | None               | None                      | None  |
| Giuseppe Ristagno      | Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico (Italy) | None   | None                   | None                       | None           | None               | ZOLL Med. Corp†           | None  |
| Federico Semeraro      | Maggiore Hospital (Italy)   | None   | None                   | None                       | None           | None               | None                      | None  |
| Christopher M. Smith   | University of Warwick, Warwick Medical School (United Kingdom)    | Resuscitation Council UK (Grant of £23890 to conduct a simulation study into delivery of AEDs by drone. Awarded January 2020)*   | None                   | None                       | None           | None               | None                      | None  |
| Michael A. Smyth       | University of Warwick (United Kingdom)                            | None   | None                   | None                       | None           | None               | None                      | None  |
| Christian Vaillancourt | University of Ottawa, The Ottawa Hospital (Canada)                | Cardiac Arrhythmia Network of Canada (CANet) (PI on study on telecommunication-assisted CPR)†; Canadian Institutes of Health Research (Co-PI for Canadian Resuscitation Outcomes Consortium (CanROC))†; Heart and Stroke Foundation of Canada (Co-PI for Canadian Resuscitation Outcomes Consortium (CanROC))† | None                   | None                       | None           | None               | None                      | None  |

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

\*Modest.

†Significant.

## Appendix 2. Reviewer Disclosures

| Reviewer          | Employment  | Research Grant      | Other Research Support | Speakers' Bureau/Honoraria | Expert Witness | Ownership Interest | Consultant/Advisory Board | Other |
|-------------------|---|---------------------|------------------------|----------------------------|----------------|--------------------|---------------------------|-------|
| Henry Halperin    | Johns Hopkins University  | Zoll Medical*; NIH* | None                   | None                       | None           | None               | None                      | None  |
| Jonathan Jui      | Oregon Health and Science University  | None                | None                   | None                       | None           | None               | None                      | None  |
| Fred Severyn      | Denver Health and Hospital Authority and University of Colorado; University of Arkansas | None                | None                   | None                       | None           | None               | None                      | None  |
| Robert A. Swor    | William Beaumont Hospital   | None                | None                   | None                       | None           | None               | None                      | None  |
| Andrew H. Travers | Emergency Health Services, Nova Scotia (Canada)   | None                | None                   | None                       | None           | None               | None                      | None  |

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10,000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10,000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

\*Significant.

## REFERENCES

- Sayre MR, Koster RW, Botha M, Cave DM, Cudnik MT, Handley AJ, Hatanaka T, Hazinski MF, Jacobs I, Monsieurs K, et al; on behalf of the Adult Basic Life Support Chapter Collaborators. Part 5: adult basic life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122(suppl 2):S298–S324. doi: 10.1161/CIRCULATIONAHA.110.970996
- Koster RW, Sayre MR, Botha M, Cave DM, Cudnik MT, Handley AJ, Hatanaka T, Hazinski MF, Jacobs I, Monsieurs K, et al. Part 5: adult basic life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2010;81(suppl 1):e48–e70. doi: 10.1016/j.resuscitation.2010.08.005
- Travers AH, Perkins GD, Berg RA, Castren M, Considine J, Escalante R, Gazmuri RJ, Koster RW, Lim SH, Nation KJ, et al; on behalf of the Basic Life Support Chapter Collaborators. Part 3: adult basic life support and automated external defibrillation: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S51–S83. doi: 10.1161/CIR.0000000000000272
- Perkins GD, Travers AH, Berg RA, Castren M, Considine J, Escalante R, Gazmuri RJ, Koster RW, Lim SH, Nation K, et al. Part 3: adult basic life support and automated external defibrillation: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e43–e69. doi: 10.1016/j.resuscitation.2015.07.041
- Olasveengen TM, de Caen AR, Mancini ME, Maconochie IK, Aickin R, Atkins DL, Berg RA, Bingham RM, Brooks SC, Castrén M, et al; on behalf of the ILCOR Collaborators. 2017 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations summary. *Circulation*. 2017;136:e424–e440. doi: 10.1161/CIR.0000000000000541
- Olasveengen TM, de Caen AR, Mancini ME, Maconochie IK, Aickin R, Atkins DL, Berg RA, Bingham RM, Brooks SC, Castrén M, et al; on behalf of the ILCOR Collaborators. 2017 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations summary. *Resuscitation*. 2017;121:201–214. doi: 10.1016/j.resuscitation.2017.10.021
- Soar J, Maconochie I, Wyckoff MH, Olasveengen TM, Singletary EM, Greif R, Aickin R, Bhanji F, Donnino MW, Mancini ME, Wyllie JP, Zideman D, Andersen LW, Atkins DL, Aziz K, Bendall J, Berg KM, Berry DC, Bigham BL, Bingham R, Couto TB, Böttiger BW, Borra V, Bray JE, Breckwoldt J, Brooks SC, Buick J, Callaway CW, Carlson JN, Cassan P, Castrén M, Chang WT, Charlton NP, Cheng A, Chung SP, Considine J, Couper K, Dainty KN, Dawson JA, de Almeida MF, de Caen AR, Deakin CD, Drennan IR, Duff JP, Epstein JL, Escalante R, Gazmuri RJ, Gilfoyle E, Granfeldt A, Guerguerian AM, Guinsburg R, Hatanaka T, Holmberg MJ, Hood N, Hosono S, Hsieh MJ, Isayama T, Iwami T, Jensen JL, Kapadia V, Kim HS, Kleinman ME, Kudenchuk PJ, Lang E, Lavonas E, Liley H, Lim SH, Lockey A, Lofgren B, Ma MH, Markenson D, Meaney PA, Meyran D, Mildenhall L, Monsieurs KG, Montgomery W, Morley PT, Morrison LJ, Nadkarni VM, Nation K, Neumar RW, Ng KC, Nicholson T, Nikolaou N, Nishiyama C, Nuthall G, Ohshimo S, Okamoto D, O'Neil B, Yong-Kwang Ong G, Paiva EF, Parr M, Pellegrino JL, Perkins GD, Perlman J, Rabi Y, Reis A, Reynolds JC, Ristagno G, Roehr CC, Sakamoto T, Sandroni C, Schexnayder SM, Schofield BR, Shimizu N, Skrifvars MB, Smyth MA, Stanton D, Swain J, Szyld E, Tijssen J, Travers A, Trevisanuto D, Vaillancourt C, Van de Voorde P, Velaphi S, Wang TL, Weiner G, Welsford M, Woodin JA, Yeung J, Nolan JP, Hazinski MF. 2019 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: summary from the basic life support; education, implementation, and teams; and first aid task forces. *Circulation*. 2019;140:e826–e880. doi: 10.1161/CIR.0000000000000734
- Soar J, Maconochie I, Wyckoff MH, Olasveengen TM, Singletary EM, Greif R, Aickin R, Bhanji F, Donnino MW, Mancini ME, Wyllie JP, Zideman D, Andersen LW, Atkins DL, Aziz K, Bendall J, Berg KM, Berry DC, Bigham BL, Bingham R, Couto TB, Böttiger BW, Borra V, Bray JE, Breckwoldt J, Brooks SC, Buick J, Callaway CW, Carlson JN, Cassan P, Castrén M, Chang WT, Charlton NP, Cheng A, Chung SP, Considine J, Couper K, Dainty KN, Dawson JA, de Almeida MF, de Caen AR, Deakin CD, Drennan IR, Duff JP, Epstein JL, Escalante R, Gazmuri RJ, Gilfoyle E, Granfeldt A, Guerguerian AM, Guinsburg R, Hatanaka T, Holmberg MJ, Hood N, Hosono S, Hsieh MJ, Isayama T, Iwami T, Jensen JL, Kapadia V, Kim HS, Kleinman ME, Kudenchuk PJ, Lang E, Lavonas E, Liley H, Lim SH, Lockey A, Lofgren B, Ma MH, Markenson D, Meaney PA, Meyran D, Mildenhall L, Monsieurs KG, Montgomery W, Morley PT, Morrison LJ, Nadkarni VM, Nation K, Neumar RW, Ng KC, Nicholson T, Nikolaou N, Nishiyama C, Nuthall G, Ohshimo S, Okamoto D, O'Neil B, Ong GY, Paiva EF, Parr M, Pellegrino JL, Perkins GD, Perlman J, Rabi Y, Reis A, Reynolds JC, Ristagno G, Roehr CC, Sakamoto T, Sandroni C, Schexnayder SM, Schofield BR, Shimizu N, Skrifvars MB, Smyth MA, Stanton D, Swain J, Szyld E, Tijssen J, Travers A, Trevisanuto D, Vaillancourt C, Van de Voorde P, Velaphi S, Wang TL, Weiner G, Welsford M, Woodin JA, Yeung J, Nolan JP, Hazinski MF. 2019 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2019;145:95–150. doi: 10.1016/j.resuscitation.2019.10.016
- PRISMA. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) website. <http://www.prisma-statement.org/>. Accessed December 31, 2019.
- 5.2.1: study limitations. In: Schünemann H, Brozek J, Guyatt G, Oxman A, eds. *GRADE Handbook*; 2013. <https://gdt.gradepro.org/app/handbook/handbook.html#h.m9385o5z3li7>. Accessed December 31, 2019.
- Morley PT, Atkins DL, Finn JC, Maconochie I, Nolan JP, Rabi Y, Singletary EM, Wang TL, Welsford M, Olasveengen TM, et al. Evidence evaluation process and management of potential conflicts of interest: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(suppl 1):S28–S40 doi: 10.1161/CIR.0000000000000891



12. Morley PT, Zaritsky A. The evidence evaluation process for the 2005 International Consensus Conference on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation*. 2005;67:167–170. doi: 10.1016/j.resuscitation.2005.09.007
13. Morley PT, Atkins DL, Billi JE, Bossaert L, Callaway CW, de Caen AR, Deakin CD, Eigel B, Hazinski MF, Hickey RW, et al. Part 3: evidence evaluation process: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122(suppl 2):S283–S290. doi: 10.1161/CIRCULATIONAHA.110.970947
14. International Liaison Committee on Resuscitation website. <https://www.licor.org>. Accessed March 3, 2020.
15. Perkins GD, Morley PT, Nolan JP, Soar J, Berg K, Olasveengen T, Wyckoff M, Greif R, Singletary N, Castren M, de Caen A, Wang T, Escalante R, Merchant RM, Hazinski M, Kloeck D, Heriot G, Couper K, Neumar R. International Liaison Committee on Resuscitation: COVID-19 consensus on science, treatment recommendations and task force insights. *Resuscitation*. 2020;151:145–147. doi: 10.1016/j.resuscitation.2020.04.035
16. Clark JJ, Culley L, Eisenberg M, Henwood DK. Accuracy of determining cardiac arrest by emergency medical dispatchers. *Ann Emerg Med*. 1994;23:1022–1026. doi: 10.1016/s0196-0644(94)70097-4
17. Castrén M, Kuisma M, Serlachius J, Skrifvars M. Do health care professionals report sudden cardiac arrest better than laymen? *Resuscitation*. 2001;51:265–268. doi: 10.1016/s0300-9572(01)00422-1
18. Garza AG, Gratton MC, Chen JJ, Carlson B. The accuracy of predicting cardiac arrest by emergency medical services dispatchers: the calling party effect. *Acad Emerg Med*. 2003;10:955–960. doi: 10.1111/j.1553-2712.2003.tb00651.x
19. Hauff SR, Rea TD, Culley LL, Kerry F, Becker L, Eisenberg MS. Factors impeding dispatcher-assisted telephone cardiopulmonary resuscitation. *Ann Emerg Med*. 2003;42:731–737. doi: 10.1016/s0196-0644(03)00423-2
20. Kuisma M, Boyd J, Väyrynen T, Repo J, Nousila-Wiik M, Holmström P. Emergency call processing and survival from out-of-hospital ventricular fibrillation. *Resuscitation*. 2005;67:89–93. doi: 10.1016/j.resuscitation.2005.04.008
21. Flynn J, Archer F, Morgans A. Sensitivity and specificity of the medical priority dispatch system in detecting cardiac arrest emergency calls in Melbourne. *Prehosp Disaster Med*. 2006;21:72–76. doi: 10.1017/s1049023x00003381
22. Nurmi J, Pettilä V, Biber B, Kuisma M, Komulainen R, Castrén M. Effect of protocol compliance to cardiac arrest identification by emergency medical dispatchers. *Resuscitation*. 2006;70:463–469. doi: 10.1016/j.resuscitation.2006.01.016
23. Böhm K, Rosenqvist M, Hollenberg J, Biber B, Engerström L, Svensson L. Dispatcher-assisted telephone-guided cardiopulmonary resuscitation: an underused lifesaving system. *Eur J Emerg Med*. 2007;14:256–259. doi: 10.1097/MEJ.0b013e32823a3cd1
24. Ma MH, Lu TC, Ng JC, Lin CH, Chiang WC, Ko PC, Shih FY, Huang CH, Hsiung KH, Chen SC, Chen WJ. Evaluation of emergency medical dispatch in out-of-hospital cardiac arrest in Taipei. *Resuscitation*. 2007;73:236–245. doi: 10.1016/j.resuscitation.2006.09.005
25. Vaillancourt C, Verma A, Trickett J, Crete D, Beaudoin T, Nesbitt L, Wells GA, Stiell IG. Evaluating the effectiveness of dispatch-assisted cardiopulmonary resuscitation instructions. *Acad Emerg Med*. 2007;14:877–883. doi: 10.1197/j.aem.2007.06.021
26. Cairns KJ, Hamilton AJ, Marshall AH, Moore MJ, Adgey AA, Kee F. The obstacles to maximising the impact of public access defibrillation: an assessment of the dispatch mechanism for out-of-hospital cardiac arrest. *Heart*. 2008;94:349–353. doi: 10.1136/hrt.2006.109785
27. Berdowski J, Beekhuis F, Zwinderman AH, Tijssen JG, Koster RW. Importance of the first link: description and recognition of an out-of-hospital cardiac arrest in an emergency call. *Circulation*. 2009;119:2096–2102. doi: 10.1161/CIRCULATIONAHA.108.768325
28. Böhm K, Stålhandske B, Rosenqvist M, Ulfvarson J, Hollenberg J, Svensson L. Tuition of emergency medical dispatchers in the recognition of agonal respiration increases the use of telephone assisted CPR. *Resuscitation*. 2009;80:1025–1028. doi: 10.1016/j.resuscitation.2009.06.004
29. Roppolo LP, Westfall A, Pepe PE, Nobel LL, Cowan J, Kay JJ, Idris AH. Dispatcher assessments for agonal breathing improve detection of cardiac arrest. *Resuscitation*. 2009;80:769–772. doi: 10.1016/j.resuscitation.2009.04.013
30. Dami F, Fuchs V, Praz L, Vader JP. Introducing systematic dispatcher-assisted cardiopulmonary resuscitation (telephone-CPR) in a non-Advanced Medical Priority Dispatch System (AMPDS): implementation process and costs. *Resuscitation*. 2010;81:848–852. doi: 10.1016/j.resuscitation.2010.03.025
31. Lewis M, Stubbs BA, Eisenberg MS. Dispatcher-assisted cardiopulmonary resuscitation: time to identify cardiac arrest and deliver chest compression instructions. *Circulation*. 2013;128:1522–1530. doi: 10.1161/CIRCULATIONAHA.113.002627
32. Hardeland C, Olasveengen TM, Lawrence R, Garrison D, Lore M, Farstad G, Wik L. Comparison of Medical Priority Dispatch (MPD) and Criteria Based Dispatch (CBD) relating to cardiac arrest calls. *Resuscitation*. 2014;85:612–616. doi: 10.1016/j.resuscitation.2014.01.029
33. Stipulante S, Tubes R, El Fassi M, Donneau AF, Van Troyen B, Hartstein G, D'Orio V, Ghuysen A. Implementation of the ALERT algorithm, a new dispatcher-assisted telephone cardiopulmonary resuscitation protocol, in non-Advanced Medical Priority Dispatch System (AMPDS) Emergency Medical Services centres. *Resuscitation*. 2014;85:177–181. doi: 10.1016/j.resuscitation.2013.10.005
34. Tanaka Y, Nishi T, Takase K, Yoshita Y, Wato Y, Taniguchi J, Hamada Y, Inaba H. Survey of a protocol to increase appropriate implementation of dispatcher-assisted cardiopulmonary resuscitation for out-of-hospital cardiac arrest. *Circulation*. 2014;129:1751–1760. doi: 10.1161/CIRCULATIONAHA.113.004409
35. Travers S, Jost D, Gillard Y, Lanoe V, Bignand M, Domanski L, Tourtier JP; Paris Fire Brigade Cardiac Arrest Work Group. Out-of-hospital cardiac arrest phone detection: those who most need chest compressions are the most difficult to recognize. *Resuscitation*. 2014;85:1720–1725. doi: 10.1016/j.resuscitation.2014.09.020
36. Besnier E, Damm C, Jardel B, Veber B, Compere V, Dureuil B. Dispatcher-assisted cardiopulmonary resuscitation protocol improves diagnosis and resuscitation recommendations for out-of-hospital cardiac arrest. *Emerg Med Australas*. 2015;27:590–596. doi: 10.1111/1742-6723.12493
37. Dami F, Heymann E, Pasquier M, Fuchs V, Carron PN, Hugli O. Time to identify cardiac arrest and provide dispatch-assisted cardio-pulmonary resuscitation in a criteria-based dispatch system. *Resuscitation*. 2015;97:27–33. doi: 10.1016/j.resuscitation.2015.09.390
38. Fukushima H, Imanishi M, Iwami T, Kitaoka H, Asai H, Seki T, Kawai Y, Norimoto K, Urisono Y, Nishio K, Okuchi K. Implementation of a dispatch-instruction protocol for cardiopulmonary resuscitation according to various abnormal breathing patterns: a population-based study. *Scand J Trauma Resusc Emerg Med*. 2015;23:64. doi: 10.1186/s13049-015-0145-8
39. Fukushima H, Imanishi M, Iwami T, Seki T, Kawai Y, Norimoto K, Urisono Y, Hata M, Nishio K, Saeki K, Kurumatani N, Okuchi K. Abnormal breathing of sudden cardiac arrest victims described by laypersons and its association with emergency medical service dispatcher-assisted cardiopulmonary resuscitation instruction. *Emerg Med J*. 2015;32:314–317. doi: 10.1136/emered-2013-203112
40. Linderöth G, Hallas P, Lippert FK, Wibrandt I, Loumann S, Möller TP, Østergaard D. Challenges in out-of-hospital cardiac arrest - A study combining closed-circuit television (CCTV) and medical emergency calls. *Resuscitation*. 2015;96:317–322. doi: 10.1016/j.resuscitation.2015.06.003
41. Orpet R, Riesenberger R, Shin J, Subido C, Markl E, Rea T. Increasing bystander CPR: potential of a one question telecommunicator identification algorithm. *Scand J Trauma Resusc Emerg Med*. 2015;23:39. doi: 10.1186/s13049-015-0115-1
42. Vaillancourt C, Charette M, Kasaboski A, Hoad M, Larocque V, Crête D, Logan S, Lamoureux P, McBride J, Cheskes S, Wells GA, Stiell IG. Cardiac arrest diagnostic accuracy of 9-1-1 dispatchers: a prospective multi-center study. *Resuscitation*. 2015;90:116–120. doi: 10.1016/j.resuscitation.2015.02.027
43. Fukushima H, Panczyk M, Spaite DW, Chikani V, Dameff C, Hu C, Birkenes TS, Myklebust H, Sutter J, Langlais B, Wu Z, Bobrow BJ. Barriers to telephone cardiopulmonary resuscitation in public and residential locations. *Resuscitation*. 2016;109:116–120. doi: 10.1016/j.resuscitation.2016.07.241
44. Hardeland C, Sunde K, Ramsdal H, Hebert SR, Soilammi L, Westmark F, Nordum F, Hansen AE, Steen-Hansen JE, Olasveengen TM. Factors impacting upon timely and adequate allocation of prehospital medical assistance and resources to cardiac arrest patients. *Resuscitation*. 2016;109:56–63. doi: 10.1016/j.resuscitation.2016.09.027
45. Ho AF, Sim ZJ, Shahidah N, Hao Y, Ng YY, Leong BS, Zarinah S, Teo WK, Goh GS, Jaafar H, Ong ME. Barriers to dispatcher-assisted cardiopulmonary resuscitation in Singapore. *Resuscitation*. 2016;105:149–155. doi: 10.1016/j.resuscitation.2016.05.006
46. Möller TP, André C, Viereck S, Todorova L, Friberg H, Lippert FK. Recognition of out-of-hospital cardiac arrest by medical dispatchers in emergency medical dispatch centres in two countries. *Resuscitation*. 2016;109:1–8. doi: 10.1016/j.resuscitation.2016.09.012
47. Plodr M, Truhlar A, Krencikova J, Praunova M, Svaba V, Masek J, Bejrova D, Paral J. Effect of introduction of a standardized protocol

- in dispatcher-assisted cardiopulmonary resuscitation. *Resuscitation*. 2016;106:18–23. doi: 10.1016/j.resuscitation.2016.05.031
48. Deakin CD, England S, Diffey D. Ambulance telephone triage using 'NHS Pathways' to identify adult cardiac arrest. *Heart*. 2017;103:738–744. doi: 10.1136/heartjnl-2016-310651
  49. Fukushima H, Panczyk M, Hu C, Dameff C, Chikani V, Vadeboncoeur T, Spaite DW, Bobrow BJ. Description of abnormal breathing is associated with improved outcomes and delayed telephone cardiopulmonary resuscitation instructions. *J Am Heart Assoc*. 2017;6:e005058. doi: 10.1161/JAHA.116.005058
  50. Hardeland C, Skåre C, Kramer-Johansen J, Birkenes TS, Myklebust H, Hansen AE, Sunde K, Olasveengen TM. Targeted simulation and education to improve cardiac arrest recognition and telephone assisted CPR in an emergency medical communication centre. *Resuscitation*. 2017;114:21–26. doi: 10.1016/j.resuscitation.2017.02.013
  51. Huang CH, Fan HJ, Chien CY, Seak CJ, Kuo CW, Ng CJ, Li WC, Weng YM. Validation of a Dispatch Protocol with Continuous Quality Control for Cardiac Arrest: A before-and-after study at a city fire department-based dispatch center. *J Emerg Med*. 2017;53:697–707. doi: 10.1016/j.jemermed.2017.06.028
  52. Nuño T, Bobrow BJ, Rogge-Miller KA, Panczyk M, Mullins T, Tormala W, Estrada A, Keim SM, Spaite DW. Disparities in telephone CPR access and timing during out-of-hospital cardiac arrest. *Resuscitation*. 2017;115:11–16. doi: 10.1016/j.resuscitation.2017.03.028
  53. Viereck S, Möller TP, Ersbøll AK, Bækgaard JS, Claesson A, Hollenberg J, Folke F, Lippert FK. Recognising out-of-hospital cardiac arrest during emergency calls increases bystander cardiopulmonary resuscitation and survival. *Resuscitation*. 2017;115:141–147. doi: 10.1016/j.resuscitation.2017.04.006
  54. Lee SY, Ro YS, Shin SD, Song KJ, Hong KJ, Park JH, Kong SY. Recognition of out-of-hospital cardiac arrest during emergency calls and public awareness of cardiopulmonary resuscitation in communities: A multilevel analysis. *Resuscitation*. 2018;128:106–111. doi: 10.1016/j.resuscitation.2018.05.008
  55. Shah M, Bartram C, Irwin K, Vellano K, McNally B, Gallagher T, Swor R. Evaluating Dispatch-Assisted CPR Using the CARES Registry. *Prehosp Emerg Care*. 2018;22:222–228. doi: 10.1080/10903127.2017.1376133
  56. Syväoja S, Salo A, Uusaro A, Jäntti H, Kuisma M. Witnessed out-of-hospital cardiac arrest- effects of emergency dispatch recognition. *Acta Anaesthesiol Scand*. 2018;62:558–567. doi: 10.1111/aas.13051
  57. Blomberg SN, Folke F, Ersbøll AK, Christensen HC, Torp-Pedersen C, Sayre MR, Counts CR, Lippert FK. Machine learning as a supportive tool to recognize cardiac arrest in emergency calls. *Resuscitation*. 2019;138:322–329. doi: 10.1016/j.resuscitation.2019.01.015
  58. Chien CY, Chien WC, Tsai LH, Tsai SL, Chen CB, Seak CJ, Chou YS, Ma M, Weng YM, Ng CJ, Lin CY, Tzeng IS, Lin CC, Huang CH. Impact of the caller's emotional state and cooperation on out-of-hospital cardiac arrest recognition and dispatcher-assisted cardiopulmonary resuscitation. *Emerg Med J*. 2019;36:595–600. doi: 10.1136/emmermed-2018-208353
  59. Derkenne C, Jost D, Thabouillot O, Briche F, Travers S, Frattini B, Lesaffre X, Kedzierewicz R, Roquet F, de Charry F, Prunet B; Paris Fire Brigade Cardiac Arrest Task Force. Improving emergency call detection of Out-of-Hospital Cardiac Arrests in the Greater Paris area: Efficiency of a global system with a new method of detection. *Resuscitation*. 2020;146:34–42. doi: 10.1016/j.resuscitation.2019.10.038
  60. Green JD, Ewings S, Wortham R, Walsh B. Accuracy of nature of call screening tool in identifying patients requiring treatment for out of hospital cardiac arrest. *Emerg Med J*. 2019;36:203–207. doi: 10.1136/emmermed-2017-207354
  61. Saberian P, Sadeghi M, Hasani-Sharamin P, Modabber M, Baratloo A. Diagnosis of out-of-hospital cardiac arrest by emergency medical dispatchers: a diagnostic accuracy study. *Aust J Paramedicine*. 2019;16.
  62. Nikolaou N, Dainty KN, Couper K, Morley P, Tijssen J, Vaillancourt C; International Liaison Committee on Resuscitation's (ILCOR) Basic Life Support and Pediatric Task Forces. A systematic review and meta-analysis of the effect of dispatcher-assisted CPR on outcomes from sudden cardiac arrest in adults and children. *Resuscitation*. 2019;138:82–105. doi: 10.1016/j.resuscitation.2019.02.035
  63. Ashoor HM, Lillie E, Zarin W, Pham B, Khan PA, Nincic V, Yazdi F, Ghassemi M, Ivory J, Cardoso R, Perkins GD, de Caen AR, Tricco AC; ILCOR Basic Life Support Task Force. Effectiveness of different compression-to-ventilation methods for cardiopulmonary resuscitation: A systematic review. *Resuscitation*. 2017;118:112–125. doi: 10.1016/j.resuscitation.2017.05.032
  64. Kudenchuk PJ, Redshaw JD, Stubbs BA, Fahrenbruch CE, Dumas F, Phelps R, Blackwood J, Rea TD, Eisenberg MS. Impact of changes in resuscitation practice on survival and neurological outcome after out-of-hospital cardiac arrest resulting from nonshockable arrhythmias. *Circulation*. 2012;125:1787–1794. doi: 10.1161/CIRCULATIONAHA.111.064873
  65. Paradis NA, Martin GB, Rivers EP, Goetting MG, Appleton TJ, Feingold M, Nowak RM. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. *JAMA*. 1990;263:1106–1113.
  66. Christenson J, Andrusiek D, Everson-Stewart S, Kudenchuk P, Hostler D, Powell J, Callaway CW, Bishop D, Vaillancourt C, Davis D, Aufderheide TP, Idris A, Stouffer JA, Stiell I, Berg R; Resuscitation Outcomes Consortium Investigators. Chest compression fraction determines survival in patients with out-of-hospital ventricular fibrillation. *Circulation*. 2009;120:1241–1247. doi: 10.1161/CIRCULATIONAHA.109.852202
  67. Stiell IG, Brown SP, Nichol G, Cheskes S, Vaillancourt C, Callaway CW, Morrison LJ, Christenson J, Aufderheide TP, Davis DP, Free C, Hostler D, Stouffer JA, Idris AH; Resuscitation Outcomes Consortium Investigators. What is the optimal chest compression depth during out-of-hospital cardiac arrest resuscitation of adult patients? *Circulation*. 2014;130:1962–1970. doi: 10.1161/CIRCULATIONAHA.114.008671
  68. Hallstrom A, Cobb L, Johnson E, Copass M. Cardiopulmonary resuscitation by chest compression alone or with mouth-to-mouth ventilation. *N Engl J Med*. 2000;342:1546–1553. doi: 10.1056/NEJM200005253422101
  69. Cheskes S, Schmicker RH, Verbeek PR, Salcido DD, Brown SP, Brooks S, Menegazzi JJ, Vaillancourt C, Powell J, May S, et al; and the Resuscitation Outcomes Consortium (ROC) investigators. The impact of per-shock pause on survival from out-of-hospital shockable cardiac arrest during the Resuscitation Outcomes Consortium PRIMED trial. *Resuscitation*. 2014;85:336–342. doi: 10.1016/j.resuscitation.2013.10.014
  70. Nichol G, Leroux B, Wang H, Callaway CW, Sopko G, Weisfeldt M, Stiell I, Morrison LJ, Aufderheide TP, Cheskes S, Christenson J, Kudenchuk P, Vaillancourt C, Rea TD, Idris AH, Colella R, Isaacs M, Straight R, Stephens S, Richardson J, Conde J, Schmicker RH, Egan D, May S, Ornato JP; ROC Investigators. Trial of Continuous or Interrupted Chest Compressions during CPR. *N Engl J Med*. 2015;373:2203–2214. doi: 10.1056/NEJMoa1509139
  71. Iwami T, Kitamura T, Kiyohara K, Kawamura T. Dissemination of Chest Compression-Only Cardiopulmonary Resuscitation and Survival After Out-of-Hospital Cardiac Arrest. *Circulation*. 2015;132:415–422. doi: 10.1161/CIRCULATIONAHA.114.014905
  72. Trowbridge C, Parekh JN, Ricard MD, Potts J, Patrickson WC, Cason CL. A randomized cross-over study of the quality of cardiopulmonary resuscitation among females performing 30:2 and hands-only cardiopulmonary resuscitation. *BMC Nurs*. 2009;8:6. doi: 10.1186/1472-6955-8-6
  73. Heidenreich JW, Berg RA, Higdon TA, Ewy GA, Kern KB, Sanders AB. Rescuer fatigue: standard versus continuous chest-compression cardiopulmonary resuscitation. *Acad Emerg Med*. 2006;13:1020–1026. doi: 10.1197/j.aem.2006.06.049
  74. Odegaard S, Saether E, Steen PA, Wik L. Quality of lay person CPR performance with compression: ventilation ratios 15:2, 30:2 or continuous chest compressions without ventilations on manikins. *Resuscitation*. 2006;71:335–340. doi: 10.1016/j.resuscitation.2006.05.012
  75. Yuksen C, Prachanukool T, Aramvanitch K, Thongwicht N, Sawanyawisuth K, Sittichanbunha Y. Is a mechanical-assist device better than manual chest compression? A randomized controlled trial. *Open Access Emerg Med*. 2017;9:63–67. doi: 10.2147/OAEM.S133074
  76. Manders S, Geijssels FE. Alternating providers during continuous chest compressions for cardiac arrest: every minute or every two minutes? *Resuscitation*. 2009;80:1015–1018. doi: 10.1016/j.resuscitation.2009.05.014
  77. Shin J, Hwang SY, Lee HJ, Park CJ, Kim YJ, Son YJ, Seo JS, Kim JJ, Lee JE, Lee IM, et al. Comparison of CPR quality and rescuer fatigue between standard 30:2 CPR and chest compression-only CPR: a randomized cross-over manikin trial. *Scand J Trauma Resusc Emerg Med*. 2014;22:59. doi: 10.1186/s13049-014-0059-x
  78. Heidenreich JW, Bonner A, Sanders AB. Rescuer fatigue in the elderly: standard vs. hands-only CPR. *J Emerg Med*. 2012;42:88–92. doi: 10.1016/j.jemermed.2010.05.019
  79. Min MK, Yeom SR, Ryu JH, Kim YI, Park MR, Han SK, Lee SH, Cho SJ. A 10-s rest improves chest compression quality during hands-only cardiopulmonary resuscitation: a prospective, randomized crossover study using a manikin model. *Resuscitation*. 2013;84:1279–1284. doi: 10.1016/j.resuscitation.2013.01.035
  80. Ashton A, McCluskey A, Gwinnutt CL, Keenan AM. Effect of rescuer fatigue on performance of continuous external chest compressions over 3 min. *Resuscitation*. 2002;55:151–155. doi: 10.1016/s0300-9572(02)00168-5
  81. Huseyin TS, Matthews AJ, Wills P, O'Neill VM. Improving the effectiveness of continuous closed chest compressions: an exploratory study. *Resuscitation*. 2002;54:57–62. doi: 10.1016/s0300-9572(02)00040-0

82. Hightower D, Thomas SH, Stone CK, Dunn K, March JA. Decay in quality of closed-chest compressions over time. *Ann Emerg Med.* 1995;26:300–303. doi: 10.1016/s0196-0644(95)70076-5
83. Lucía A, de las Heras JF, Pérez M, Elvira JC, Carvajal A, Alvarez AJ, Chicharro JL. The importance of physical fitness in the performance of adequate cardiopulmonary resuscitation. *Chest.* 1999;115:158–164. doi: 10.1378/chest.115.1.158
84. Ochoa FJ, Ramalle-Gómara E, Lisa V, Saralegui I. The effect of rescuer fatigue on the quality of chest compressions. *Resuscitation.* 1998;37:149–152. doi: 10.1016/s0300-9572(98)00057-4
85. Riera SQ, González BS, Alvarez JT, Fernández Mdel M, Saura JM. The physiological effect on rescuers of doing 2min of uninterrupted chest compressions. *Resuscitation.* 2007;74:108–112. doi: 10.1016/j.resuscitation.2006.10.031
86. Skulec R, Truhlar A, Vondruska V, Parizkova R, Dudakova J, Astapenko D, Suchy T, Lehmann C, Cerny V. Rescuer fatigue does not correlate to energy expenditure during simulated basic life support. *Signa Vitae.* 2016;12:58–62. doi: 10.22514/SV121.102016.10
87. Oh J, Chee Y, Song Y, Lim T, Kang H, Cho Y. A novel method to decrease mattress compression during CPR using a mattress compression cover and a vacuum pump. *Resuscitation.* 2013;84:987–991. doi: 10.1016/j.resuscitation.2012.12.027
88. Perkins GD, Benny R, Giles S, Gao F, Tweed MJ. Do different mattresses affect the quality of cardiopulmonary resuscitation? *Intensive Care Med.* 2003;29:2330–2335. doi: 10.1007/s00134-003-2014-6
89. Tweed M, Tweed C, Perkins GD. The effect of differing support surfaces on the efficacy of chest compressions using a resuscitation manikin model. *Resuscitation.* 2001;51:179–183. doi: 10.1016/s0300-9572(01)00404-x
90. Song Y, Oh J, Lim T, Chee Y. A new method to increase the quality of cardiopulmonary resuscitation in hospital. *Conf Proc IEEE Eng Med Biol Soc.* 2013;2013:469–472. doi: 10.1109/EMBC.2013.6609538
91. Jäntti H, Silfvast T, Turpeinen A, Kiviniemi V, Uusaro A. Quality of cardiopulmonary resuscitation on manikins: on the floor and in the bed. *Acta Anaesthesiol Scand.* 2009;53:1131–1137. doi: 10.1111/j.1399-6576.2009.01966.x
92. Ahn HJ, Cho Y, You YH, Min JH, Jeong WJ, Ryu S, Lee JW, Cho SU, Oh SK, Park JS, et al. Effect of using a home-bed mattress on bystander chest compression during out-of-hospital cardiac arrest. *Hong Kong J Emerg Med.* 2019. <https://doi.org/10.1177/1024907919856485> Accessed March 20, 2020.
93. Andersen LØ, Isbye DL, Rasmussen LS. Increasing compression depth during manikin CPR using a simple backboard. *Acta Anaesthesiol Scand.* 2007;51:747–750. doi: 10.1111/j.1399-6576.2007.01304.x
94. Fischer EJ, Mayrand K, Ten Eyck RP. Effect of a backboard on compression depth during cardiac arrest in the ED: a simulation study. *Am J Emerg Med.* 2016;34:274–277. doi: 10.1016/j.ajem.2015.10.035
95. Perkins GD, Smith CM, Augre C, Allan M, Rogers H, Stephenson B, Thickett DR. Effects of a backboard, bed height, and operator position on compression depth during simulated resuscitation. *Intensive Care Med.* 2006;32:1632–1635. doi: 10.1007/s00134-006-0273-8
96. Sanri E, Karacabey S. The Impact of Backboard Placement on Chest Compression Quality: A Mannequin Study. *Prehosp Disaster Med.* 2019;34:182–187. doi: 10.1017/S1049023X19000153
97. Sato H, Komazawa N, Ueki R, Yamamoto N, Fujii A, Nishi S, Kaminoh Y. Backboard insertion in the operating table increases chest compression depth: a manikin study. *J Anesth.* 2011;25:770–772. doi: 10.1007/s00540-011-1196-2
98. Putzer G, Fiala A, Braun P, Neururer S, Biechl K, Keilig B, Ploner W, Fop E, Paal P. Manual versus mechanical chest compressions on surfaces of varying softness with or without backboards: a randomized, crossover manikin study. *J Emerg Med.* 2016;50:594–600.e1. doi: 10.1016/j.jemermed.2015.10.002
99. Lin Y, Wan B, Belanger C, Hecker K, Gilfoyle E, Davidson J, Cheng A. Reducing the impact of intensive care unit mattress compressibility during CPR: a simulation-based study. *Adv Simul (Lond).* 2017;2:22. doi: 10.1186/s41077-017-0057-y
100. Noordergraaf GJ, Paulussen IW, Venema A, van Berkomp PF, Woerlee PH, Scheffer GJ, Noordergraaf A. The impact of compliant surfaces on in-hospital chest compressions: effects of common mattresses and a backboard. *Resuscitation.* 2009;80:546–552. doi: 10.1016/j.resuscitation.2009.03.023
101. Beesems SG, Koster RW. Accurate feedback of chest compression depth on a manikin on a soft surface with correction for total body displacement. *Resuscitation.* 2014;85:1439–1443. doi: 10.1016/j.resuscitation.2014.08.005
102. Nishisaki A, Maltese MR, Niles DE, Sutton RM, Urbano J, Berg RA, Nadkarni VM. Backboards are important when chest compressions are provided on a soft mattress. *Resuscitation.* 2012;83:1013–1020. doi: 10.1016/j.resuscitation.2012.01.016
103. Lee S, Oh J, Kang H, Lim T, Kim W, Chee Y, Song Y, Ahn C, Cho JH. Proper target depth of an accelerometer-based feedback device during CPR performed on a hospital bed: a randomized simulation study. *Am J Emerg Med.* 2015;33:1425–1429. doi: 10.1016/j.ajem.2015.07.010
104. Oh J, Song Y, Kang B, Kang H, Lim T, Suh Y, Chee Y. The use of dual accelerometers improves measurement of chest compression depth. *Resuscitation.* 2012;83:500–504. doi: 10.1016/j.resuscitation.2011.09.028
105. Ruiz de Gauna S, González-Otero DM, Ruiz J, Gutiérrez JJ, Russell JK. A feasibility study for measuring accurate chest compression depth and rate on soft surfaces using two accelerometers and spectral analysis. *Biomed Res Int.* 2016;2016:6596040. doi: 10.1155/2016/6596040
106. Hellevuo H, Sainio M, Huhtala H, Olkkola KT, Tenhunen J, Hoppu S. The quality of manual chest compressions during transport—effect of the mattress assessed by dual accelerometers. *Acta Anaesthesiol Scand.* 2014;58:323–328. doi: 10.1111/aas.12245
107. Cloete G, Dellimore KH, Scheffer C, Smuts MS, Wallis LA. The impact of backboard size and orientation on sternum-to-spine compression depth and compression stiffness in a manikin study of CPR using two mattress types. *Resuscitation.* 2011;82:1064–1070. doi: 10.1016/j.resuscitation.2011.04.003
108. Cloete G, Dellimore KH, Scheffer C. Comparison of experimental chest compression data to a theoretical model for the mechanics of constant peak displacement cardiopulmonary resuscitation. *Acad Emerg Med.* 2011;18:1167–1176. doi: 10.1111/j.1553-2712.2011.01213.x
109. Cloete G, Dellimore K, Scheffer C. The impact of various backboard configurations on compression stiffness in a manikin study of CPR. *Conf Proc IEEE Eng Med Biol Soc.* 2011;2011:2484–2487. doi: 10.1109/EMBS.2011.6090689
110. Cheng A, Belanger C, Wan B, Davidson J, Lin Y. Effect of emergency department mattress compressibility on chest compression depth using a standardized cardiopulmonary resuscitation board, a slider transfer board, and a flat spine board: a simulation-based study. *Simul Healthc.* 2017;12:364–369. doi: 10.1097/SIH.0000000000000245
111. Perkins GD, Kocierz L, Smith SC, McCulloch RA, Davies RP. Compression feedback devices overestimate chest compression depth when performed on a bed. *Resuscitation.* 2009;80:79–82. doi: 10.1016/j.resuscitation.2008.08.011
112. Marsch S, Tschann F, Semmer NK, Zobrist R, Hunziker PR, Hunziker S. ABC versus CAB for cardiopulmonary resuscitation: a prospective, randomized simulator-based trial. *Swiss Med Wkly.* 2013;143:w13856. doi: 10.4414/smw.2013.13856
113. Lubrano R, Cecchetti C, Bellelli E, Gentile I, Loayza Levano H, Orsini F, Bertazzoni G, Messi G, Rugolotto S, Pirozzi N, Elli M. Comparison of times of intervention during pediatric CPR maneuvers using ABC and CAB sequences: a randomized trial. *Resuscitation.* 2012;83:1473–1477. doi: 10.1016/j.resuscitation.2012.04.011
114. Kobayashi M, Fujiwara A, Morita H, Nishimoto Y, Mishima T, Nitta M, Hayashi T, Hotta T, Hayashi Y, Hachisuka E, Sato K. A manikin-based observational study on cardiopulmonary resuscitation skills at the Osaka Senri medical rally. *Resuscitation.* 2008;78:333–339. doi: 10.1016/j.resuscitation.2008.03.230
115. Sekiguchi H, Kondo Y, Kukita I. Verification of changes in the time taken to initiate chest compressions according to modified basic life support guidelines. *Am J Emerg Med.* 2013;31:1248–1250. doi: 10.1016/j.ajem.2013.02.047
116. Kamikura T, Iwasaki H, Myojo Y, Sakagami S, Takei Y, Inaba H. Advantage of CPR-first over call-first actions for out-of-hospital cardiac arrests in nonelderly patients and of noncardiac aetiology. *Resuscitation.* 2015;96:37–45. doi: 10.1016/j.resuscitation.2015.06.027
117. Wik L, Hansen TB, Fylling F, Steen T, Vaagenes P, Auestad BH, Steen PA. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with out-of-hospital ventricular fibrillation: a randomized trial. *JAMA.* 2003;289:1389–1395. doi: 10.1001/jama.289.11.1389
118. Baker PW, Conway J, Cotton C, Ashby DT, Smyth J, Woodman RJ, Grantham H; Clinical Investigators. Defibrillation or cardiopulmonary resuscitation first for patients with out-of-hospital cardiac arrests found by paramedics to be in ventricular fibrillation? A randomised control trial. *Resuscitation.* 2008;79:424–431. doi: 10.1016/j.resuscitation.2008.07.017
119. Orlowski JP. Optimum position for external cardiac compression in infants and young children. *Ann Emerg Med.* 1986;15:667–673. doi: 10.1016/s0196-0644(86)80423-1



120. Cha KC, Kim HJ, Shin HJ, Kim H, Lee KH, Hwang SO. Hemodynamic effect of external chest compressions at the lower end of the sternum in cardiac arrest patients. *J Emerg Med*. 2013;44:691–697. doi: 10.1016/j.jemermed.2012.09.026
121. Qvigstad E, Kramer-Johansen J, Tømte Ø, Skålhegg T, Sørensen Ø, Sunde K, Olasveengen TM. Clinical pilot study of different hand positions during manual chest compressions monitored with capnography. *Resuscitation*. 2013;84:1203–1207. doi: 10.1016/j.resuscitation.2013.03.010
122. Considine J, Gazmuri RJ, Perkins GD, Kudenchuk PJ, Olasveengen TM, Vaillancourt C, Nishiyama C, Hatanaka T, Mancini ME, Chung SP, Escalante-Kanashiro R, Morley P. Chest compression components (rate, depth, chest wall recoil and leaning): A scoping review. *Resuscitation*. 2020;146:188–202. doi: 10.1016/j.resuscitation.2019.08.042
123. Cheskes S, Common MR, Byers AP, Zhan C, Silver A, Morrison LJ. The association between chest compression release velocity and outcomes from out-of-hospital cardiac arrest. *Resuscitation*. 2015;86:38–43. doi: 10.1016/j.resuscitation.2014.10.020
124. Hwang SO, Cha KC, Kim K, Jo YH, Chung SP, You JS, Shin J, Lee HJ, Park YS, Kim S, et al. A randomized controlled trial of compression rates during cardiopulmonary resuscitation. *J Korean Med Sci*. 2016;31:1491–1498. doi: 10.3346/jkms.2016.31.9.1491
125. Kilgannon JH, Kirchhoff M, Pierce L, Aunchman N, Trzeciak S, Roberts BW. Association between chest compression rates and clinical outcomes following in-hospital cardiac arrest at an academic tertiary hospital. *Resuscitation*. 2017;110:154–161. doi: 10.1016/j.resuscitation.2016.09.015
126. Kovacs A, Vadeboncoeur TF, Stolz U, Spaite DW, Irisawa T, Silver A, Bobrow BJ. Chest compression release velocity: Association with survival and favorable neurologic outcome after out-of-hospital cardiac arrest. *Resuscitation*. 2015;92:107–114. doi: 10.1016/j.resuscitation.2015.04.026
127. Sainio M, Hopppu S, Huhtala H, Eilevstjønn J, Olkkola KT, Tenhunen J. Simultaneous beat-to-beat assessment of arterial blood pressure and quality of cardiopulmonary resuscitation in out-of-hospital and in-hospital settings. *Resuscitation*. 2015;96:163–169. doi: 10.1016/j.resuscitation.2015.08.004
128. Sutton RM, Case E, Brown SP, Atkins DL, Nadkarni VM, Kaltman J, Callaway C, Idris A, Nichol G, Hutchison J, Drennan IR, Austin M, Daya M, Cheskes S, Nuttall J, Herren H, Christenson J, Andrusiek D, Vaillancourt C, Menegazzi JJ, Rea TD, Berg RA; ROC Investigators. A quantitative analysis of out-of-hospital pediatric and adolescent resuscitation quality—A report from the ROC epistry-cardiac arrest. *Resuscitation*. 2015;93:150–157. doi: 10.1016/j.resuscitation.2015.04.010
129. Sutton RM, Reeder RW, Landis W, Meert KL, Yates AR, Berger JT, Newth CJ, Carcillo JA, McQuillen PS, Harrison RE, Moler FW, Pollack MM, Carpenter TC, Notterman DA, Holubkov R, Dean JM, Nadkarni VM, Berg RA; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network (CPCCRN) Investigators. Chest compression rates and pediatric in-hospital cardiac arrest survival outcomes. *Resuscitation*. 2018;130:159–166. doi: 10.1016/j.resuscitation.2018.07.015
- 129a. Riyapan S, Naulnark T, Ruangsomboon O, Chaisirin W, Limsuwat C, Prapruetkit N, Chakorn T, Monsomboon A. Improving quality of chest compression in Thai emergency department by using real-time audiovisual feedback cardio-pulmonary resuscitation monitoring. *J Med Assoc Thai*. 2019;102:245–251.
130. Edelson DP, Abella BS, Kramer-Johansen J, Wik L, Myklebust H, Barry AM, Merchant RM, Hoek TL, Steen PA, Becker LB. Effects of compression depth and pre-shock pauses predict defibrillation failure during cardiac arrest. *Resuscitation*. 2006;71:137–145. doi: 10.1016/j.resuscitation.2006.04.008
131. Kramer-Johansen J, Myklebust H, Wik L, Fellows B, Svensson L, Sørebo H, Steen PA. Quality of out-of-hospital cardiopulmonary resuscitation with real time automated feedback: a prospective interventional study. *Resuscitation*. 2006;71:283–292. doi: 10.1016/j.resuscitation.2006.05.011
132. Kern KB, Sanders AB, Raife J, Milander MM, Otto CW, Ewy GA. A study of chest compression rates during cardiopulmonary resuscitation in humans. The importance of rate-directed chest compressions. *Arch Intern Med*. 1992;152:145–149.
133. Idris AH, Guffey D, Pepe PE, Brown SP, Brooks SC, Callaway CW, Christenson J, Davis DP, Daya MR, Gray R, Kudenchuk PJ, Larsen J, Lin S, Menegazzi JJ, Sheehan K, Sopko G, Stiell I, Nichol G, Aufderheide TP; Resuscitation Outcomes Consortium Investigators. Chest compression rates and survival following out-of-hospital cardiac arrest. *Crit Care Med*. 2015;43:840–848. doi: 10.1097/CCM.0000000000000824
134. Idris AH, Guffey D, Aufderheide TP, Brown S, Morrison LJ, Nichols P, Powell J, Daya M, Bigham BL, Atkins DL, Berg R, Davis D, Stiell I, Sopko G, Nichol G; Resuscitation Outcomes Consortium (ROC) Investigators. Relationship between chest compression rates and outcomes from cardiac arrest. *Circulation*. 2012;125:3004–3012. doi: 10.1161/CIRCULATIONAHA.111.059535
135. Abella BS, Sandbo N, Vassilatos P, Alvarado JP, O'Hearn N, Wigder HN, Hoffman P, Tynus K, Vanden Hoek TL, Becker LB. Chest compression rates during cardiopulmonary resuscitation are suboptimal: a prospective study during in-hospital cardiac arrest. *Circulation*. 2005;111:428–434. doi: 10.1161/01.CIR.0000153811.84257.59
136. Ornato JP, Gonzalez ER, Garnett AR, Levine RL, McClung BK. Effect of cardiopulmonary resuscitation compression rate on end-tidal carbon dioxide concentration and arterial pressure in man. *Crit Care Med*. 1988;16:241–245. doi: 10.1097/00003246-198803000-00007
137. Bohn A, Weber TP, Wecker S, Harding U, Osada N, Van Aken H, Lukas RP. The addition of voice prompts to audiovisual feedback and debriefing does not modify CPR quality or outcomes in out of hospital cardiac arrest—a prospective, randomized trial. *Resuscitation*. 2011;82:257–262. doi: 10.1016/j.resuscitation.2010.11.006
138. Vadeboncoeur T, Stolz U, Panchal A, Silver A, Venuti M, Tobin J, Smith G, Nunez M, Karammooz M, Spaite D, Bobrow B. Chest compression depth and survival in out-of-hospital cardiac arrest. *Resuscitation*. 2014;85:182–188. doi: 10.1016/j.resuscitation.2013.10.002
139. Hellevuo H, Sainio M, Nevalainen R, Huhtala H, Olkkola KT, Tenhunen J, Hopppu S. Deeper chest compression - more complications for cardiac arrest patients? *Resuscitation*. 2013;84:760–765. doi: 10.1016/j.resuscitation.2013.02.015
140. Stiell IG, Brown SP, Christenson J, Cheskes S, Nichol G, Powell J, Bigham B, Morrison LJ, Larsen J, Hess E, Vaillancourt C, Davis DP, Callaway CW; Resuscitation Outcomes Consortium (ROC) Investigators. What is the role of chest compression depth during out-of-hospital cardiac arrest resuscitation? *Crit Care Med*. 2012;40:1192–1198. doi: 10.1097/CCM.0b013e31823bc8bb
141. Babbis CF, Kemeny AE, Quan W, Freeman G. A new paradigm for human resuscitation research using intelligent devices. *Resuscitation*. 2008;77:306–315. doi: 10.1016/j.resuscitation.2007.12.018
142. Sutton RM, French B, Niles DE, Donoghue A, Topjian AA, Nishisaki A, Effelman J, Wolfe H, Berg RA, Nadkarni VM, et al. 2010 American Heart Association recommended compression depths during pediatric in-hospital resuscitations are associated with survival. *Resuscitation*. 2014;85:1179–1184. doi: 10.1016/j.resuscitation.2014.05.007
143. Jost D, Degrange H, Verret C, Hersan O, Banville IL, Chapman FW, Lank P, Petit JL, Fuilla C, Migliani R, et al; and the DEFI 2005 Work Group. DEFI 2005: a randomized controlled trial of the effect of automated external defibrillator cardiopulmonary resuscitation protocol on outcome from out-of-hospital cardiac arrest. *Circulation*. 2010;121:1614–1622. doi: 10.1161/CIRCULATIONAHA.109.878389
144. Berdowski J, Tijssen JG, Koster RW. Chest compressions cause recurrence of ventricular fibrillation after the first successful conversion by defibrillation in out-of-hospital cardiac arrest. *Circ Arrhythm Electrophysiol*. 2010;3:72–78. doi: 10.1161/CIRCEP.109.902114
145. Beesems SG, Berdowski J, Hulleman M, Blom MT, Tijssen JG, Koster RW. Minimizing pre- and post-shock pauses during the use of an automatic external defibrillator by two different voice prompt protocols. A randomized controlled trial of a bundle of measures. *Resuscitation*. 2016;106:1–6. doi: 10.1016/j.resuscitation.2016.06.009
146. Kellum MJ, Kennedy KW, Ewy GA. Cardiocerebral resuscitation improves survival of patients with out-of-hospital cardiac arrest. *Am J Med*. 2006;119:335–340. doi: 10.1016/j.amjmed.2005.11.014
147. Rea TD, Helbock M, Perry S, Garcia M, Cloyd D, Becker L, Eisenberg M. Increasing use of cardiopulmonary resuscitation during out-of-hospital ventricular fibrillation arrest: survival implications of guideline changes. *Circulation*. 2006;114:2760–2765. doi: 10.1161/CIRCULATIONAHA.106.654715
148. Bobrow BJ, Clark LL, Ewy GA, Chikani V, Sanders AB, Berg RA, Richman PB, Kern KB. Minimally interrupted cardiac resuscitation by emergency medical services for out-of-hospital cardiac arrest. *JAMA*. 2008;299:1158–1165. doi: 10.1001/jama.299.10.1158
149. Conover Z, Kern KB, Silver AE, Bobrow BJ, Spaite DW, Indik JH. Resumption of chest compressions after successful defibrillation and risk for recurrence of ventricular fibrillation in out-of-hospital cardiac arrest. *Circ Arrhythm Electrophysiol*. 2014;7:633–639. doi: 10.1161/CIRCEP.114.001506
150. Hostler D, Everson-Stewart S, Rea TD, Stiell IG, Callaway CW, Kudenchuk PJ, Sears GK, Emerson SS, Nichol G; Resuscitation Outcomes Consortium (ROC) Investigators. Relationship between chest compression rates and outcomes from cardiac arrest. *Circulation*. 2012;125:3004–3012. doi: 10.1161/CIRCULATIONAHA.111.059535



- Consortium Investigators. Effect of real-time feedback during cardiopulmonary resuscitation outside hospital: prospective, cluster-randomised trial. *BMJ*. 2011;342:d512. doi: 10.1136/bmj.d512
151. Couper K, Kimani PK, Abella BS, Chilwan M, Cooke MW, Davies RP, Field RA, Gao F, Quinton S, Stallard N, Woolley S, Perkins GD; Cardiopulmonary Resuscitation Quality Improvement Initiative Collaborators. The system-wide effect of real-time audiovisual feedback and postevent debriefing for in-hospital cardiac arrest: the cardiopulmonary resuscitation quality improvement initiative. *Crit Care Med*. 2015;43:2321–2331. doi: 10.1097/CCM.0000000000001202
  152. Sainio M, Kämäräinen A, Huhtala H, Aaltonen P, Tenhunen J, Olkkola KT, Hoppu S. Real-time audiovisual feedback system in a physician-staffed helicopter emergency medical service in Finland: the quality results and barriers to implementation. *Scand J Trauma Resusc Emerg Med*. 2013;21:50. doi: 10.1186/1757-7241-21-50
  153. Bobrow BJ, Vadeboncoeur TF, Stolz U, Silver AE, Tobin JM, Crawford SA, Mason TK, Schirmer J, Smith GA, Spaite DW. The influence of scenario-based training and real-time audiovisual feedback on out-of-hospital cardiopulmonary resuscitation quality and survival from out-of-hospital cardiac arrest. *Ann Emerg Med*. 2013;62:47–56.e1. doi: 10.1016/j.annemergmed.2012.12.020
  154. Riyapan S, Naulnark T, Ruangsomboon O, W. C, Limsuwat C, Prapruetkit N, Chakorn T, Monsomboon A. Improving quality of chest compression in Thai emergency department by using real-time audiovisual feedback cardio-pulmonary resuscitation monitoring. *J Med Assoc Thai*. 2019;245–251.
  155. Abella BS, Edelson DP, Kim S, Retzer E, Myklebust H, Barry AM, O'Hearn N, Hoek TL, Becker LB. CPR quality improvement during in-hospital cardiac arrest using a real-time audiovisual feedback system. *Resuscitation*. 2007;73:54–61. doi: 10.1016/j.resuscitation.2006.10.027
  156. Sutton RM, Niles D, French B, Maltese MR, Leffelman J, Eilevstjonn J, Wolfe H, Nishisaki A, Meaney PA, Berg RA, et al. First quantitative analysis of cardiopulmonary resuscitation quality during in-hospital cardiac arrests of young children. *Resuscitation*. 2014;85:70–74. doi: 10.1016/j.resuscitation.2013.08.014
  157. Agerskov M, Hansen MB, Nielsen AM, Møller TP, Wissenberg M, Rasmussen LS. Return of spontaneous circulation and long-term survival according to feedback provided by automated external defibrillators. *Acta Anaesthesiol Scand*. 2017;61:1345–1353. doi: 10.1111/aas.12992
  158. Lukas RP, Gräsner JT, Seewald S, Lefering R, Weber TP, Van Aken H, Fischer M, Bohn A. Chest compression quality management and return of spontaneous circulation: a matched-pair registry study. *Resuscitation*. 2012;83:1212–1218. doi: 10.1016/j.resuscitation.2012.03.027
  159. Goharani R, Vahedian-Azimi A, Farzanegan B, Bashar FR, Hajiesmaeili M, Shojaei S, Madani SJ, Gohari-Moghaddam K, Hatamian S, Mosavinasab SMM, Khoshfetrat M, Khabiri Khatir MA, Miller AC; MORZAK Collaborative. Real-time compression feedback for patients with in-hospital cardiac arrest: a multi-center randomized controlled clinical trial. *J Intensive Care*. 2019;7:5. doi: 10.1186/s40560-019-0357-5
  160. Vahedian-Azimi A, Hajiesmaeili M, Amirsavadkouhi A, Jamaati H, Izadi M, Madani SJ, Hashemian SM, Miller AC. Effect of the Cardio First Angel™ device on CPR indices: a randomized controlled clinical trial. *Crit Care*. 2016;20:147. doi: 10.1186/s13054-016-1296-3
  161. Chiang WC, Chen WJ, Chen SY, Ko PC, Lin CH, Tsai MS, Chang WT, Chen SC, Tsan CY, Ma MH. Better adherence to the guidelines during cardiopulmonary resuscitation through the provision of audio-prompts. *Resuscitation*. 2005;64:297–301. doi: 10.1016/j.resuscitation.2004.09.010
  162. Caldwell G, Millar G, Quinn E, Vincent R, Chamberlain DA. Simple mechanical methods for cardioversion: defence of the precordial thump and cough version. *Br Med J (Clin Res Ed)*. 1985;291:627–630. doi: 10.1136/bmj.291.6496.627
  163. Petelenz T, Iwiński J, Chlebowczyk J, Czyz Z, Flak Z, Fiutowski L, Zaorski K, Petelenz T, Zeman S. Self-administered cough cardiopulmonary resuscitation (c-CPR) in patients threatened by MAS events of cardiovascular origin. *Wiad Lek*. 1998;51:326–336.
  164. Marozsán I, Albared JL, Szatmáry LJ. Life-threatening arrhythmias stopped by cough. *Cor Vasa*. 1990;32:401–408.
  165. Niemann JT, Rosborough J, Hausknecht M, Brown D, Criley JM. Cough-CPR: documentation of systemic perfusion in man and in an experimental model: a “window” to the mechanism of blood flow in external CPR. *Crit Care Med*. 1980;8:141–146. doi: 10.1097/00003246-198003000-00011
  166. Nehme Z, Andrew E, Bernard SA, Smith K. Treatment of monitored out-of-hospital ventricular fibrillation and pulseless ventricular tachycardia utilising the precordial thump. *Resuscitation*. 2013;84:1691–1696. doi: 10.1016/j.resuscitation.2013.08.011
  167. Pellis T, Kette F, Lovisa D, Franceschino E, Magagnin L, Mercante WP, Kohl P. Utility of pre-cordial thump for treatment of out of hospital cardiac arrest: a prospective study. *Resuscitation*. 2009;80:17–23. doi: 10.1016/j.resuscitation.2008.10.018
  168. Gertsch M, Hottinger S, Hess T. Serial chest thumps for the treatment of ventricular tachycardia in patients with coronary artery disease. *Clin Cardiol*. 1992;15:181–188. doi: 10.1002/clc.4960150309
  169. Rajagopalan RS, Appu KS, Sultan SK, Jagannadham TG, Nityanandan K, Sethuraman S. Precordial thump in ventricular tachycardia. *J Assoc Physicians India*. 1971;19:725–729.
  170. Miller J, Tresch D, Horwitz L, Thompson BM, Aprahamian C, Darin JC. The precordial thump. *Ann Emerg Med*. 1984;13(9Pt2):791–794. doi: 10.1016/s0196-0644(84)80439-4
  171. Rahner E, Zeh E. [Regulation of ventricular tachycardia with precordial fist blow]. *Med Welt*. 1978;29:1659–1663.
  172. Cotoi S, Ștefănescu M, Moldovan D. An attempt to localize the accessory pathway in WPW syndrome using electrocardiographic methods. *Med Interne*. 1980;18:265–277.
  173. Pennington JE, Taylor J, Lown B. Chest thump for reverting ventricular tachycardia. *N Engl J Med*. 1970;283:1192–1195. doi: 10.1056/NEJM197011262832204
  174. Morgera T, Baldi N, Chersevani D, Medugno G, Camerini F. Chest thump and ventricular tachycardia. *Pacing Clin Electrophysiol*. 1979;2:69–75. doi: 10.1111/j.1540-8159.1979.tb05178.x
  175. Haman L, Parizek P, Vojacek J. Precordial thump efficacy in termination of induced ventricular arrhythmias. *Resuscitation*. 2009;80:14–16. doi: 10.1016/j.resuscitation.2008.07.022
  176. Amir O, Schliamser JE, Nemer S, Arie M. Ineffectiveness of precordial thump for cardioversion of malignant ventricular tachyarrhythmias. *Pacing Clin Electrophysiol*. 2007;30:153–156. doi: 10.1111/j.1540-8159.2007.00643.x
  177. Befeler B. Mechanical stimulation of the heart: its therapeutic value in tachyarrhythmias. *Chest*. 1978;73:832–838. doi: 10.1378/chest.73.6.832
  178. Miller J, Addas A, Akhtar M. Electrophysiology studies: precordial thumping patients paced into ventricular tachycardia. *J Emerg Med*. 1985;3:175–179. doi: 10.1016/0736-4679(85)90069-1
  179. Nejima J. [Clinical features and treatment of ventricular tachycardia associated with acute myocardial infarction]. *Nihon Ika Daigaku Zasshi*. 1991;58:40–49.
  180. Volkmann H, Klumbies A, Kühnert H, Paliege R, Dannberg G, Siegert K. [Terminating ventricular tachycardias by mechanical heart stimulation with precordial thumps]. *Z Kardiol*. 1990;79:717–724.
  181. Klumbies A, Paliege R, Volkmann H. [Mechanical emergency stimulation in asystole and extreme bradycardia]. *Z Gesamte Inn Med*. 1988;43:348–352.
  182. Scherf D, Bornemann C. Thumping of the precordium in ventricular standstill. *Am J Cardiol*. 1960;5:30–40. doi: 10.1016/0002-9149(60)90006-0
  183. Iseri LT, Allen BJ, Baron K, Brodsky MA. Fist pacing, a forgotten procedure in bradyasystolic cardiac arrest. *Am Heart J*. 1987;113:1545–1550. doi: 10.1016/0002-8703(87)90697-1
  184. Paliege R, Volkmann H, Klumbies A. The fist as a pacemaker for the heart—investigations about the mechanical stimulation of the heart in case of emergency. *Deutsche Gesundheitswesen Zeitschrift für Klinische Medizin*. 1982;37:1094–1100.
  185. Holmberg MJ, Vognsen M, Andersen MS, Donnino MW, Andersen LW. Bystander automated external defibrillator use and clinical outcomes after out-of-hospital cardiac arrest: A systematic review and meta-analysis. *Resuscitation*. 2017;120:77–87. doi: 10.1016/j.resuscitation.2017.09.003
  186. Bækgaard JS, Viereck S, Møller TP, Ersbøll AK, Lippert F, Folke F. The Effects of Public Access Defibrillation on Survival After Out-of-Hospital Cardiac Arrest: A Systematic Review of Observational Studies. *Circulation*. 2017;136:954–965. doi: 10.1161/CIRCULATIONAHA.117.029067
  187. Gianotto-Oliveira R, Gonzalez MM, Vianna CB, Monteiro Alves M, Timerman S, Kalil Filho R, Kern KB. Survival after ventricular fibrillation cardiac arrest in the Sao Paulo metropolitan subway system: first successful targeted automated external defibrillator (AED) program in Latin America. *J Am Heart Assoc*. 2015;4:e002185. doi: 10.1161/JAHA.115.002185
  188. Nakahara S, Tomio J, Ichikawa M, Nakamura F, Nishida M, Takahashi H, Morimura N, Sakamoto T. Association of bystander interventions with neurologically intact survival among patients with bystander-witnessed out-of-hospital cardiac arrest in Japan. *JAMA*. 2015;314:247–254. doi: 10.1001/jama.2015.8068

189. Kitamura T, Kiyohara K, Sakai T, Matsuyama T, Hatakeyama T, Shimamoto T, Izawa J, Fujii T, Nishiyama C, Kawamura T, Iwami T. Public-access defibrillation and out-of-hospital cardiac arrest in Japan. *N Engl J Med*. 2016;375:1649–1659. doi: 10.1056/NEJMsa1600011
190. Fukuda T, Ohashi-Fukuda N, Kobayashi H, Gunshin M, Sera T, Kondo Y, Yahagi N. Public access defibrillation and outcomes after pediatric out-of-hospital cardiac arrest. *Resuscitation*. 2017;111:1–7. doi: 10.1016/j.resuscitation.2016.11.010
191. Takeuchi I, Nagasawa H, Jitsuiki K, Kondo A, Ohsaka H, Yanagawa Y. Impact of automated external defibrillator as a recent innovation for the resuscitation of cardiac arrest patients in an urban city of Japan. *J Emerg Trauma Shock*. 2018;11:217–220. doi: 10.4103/JETS.JETS\_79\_17
192. Kiguchi T, Kiyohara K, Kitamura T, Nishiyama C, Kobayashi D, Okabayashi S, Shimamoto T, Matsuyama T, Kawamura T, Iwami T. Public-access defibrillation and survival of out-of-hospital cardiac arrest in public vs. residential locations in Japan. *Circ J*. 2019;83:1682–1688. doi: 10.1253/circj.CJ-19-0065
193. Tay PJM, Pek PP, Fan Q, Ng YY, Leong BS, Gan HN, Mao DR, Chia MYC, Cheah SO, Doctor N, Tham LP, Ong MEH. Effectiveness of a community based out-of-hospital cardiac arrest (OHCA) interventional bundle: Results of a pilot study. *Resuscitation*. 2020;146:220–228. doi: 10.1016/j.resuscitation.2019.10.015
194. Matsui S, Kitamura T, Sado J, Kiyohara K, Kobayashi D, Kiguchi T, Nishiyama C, Okabayashi S, Shimamoto T, Matsuyama T, Kawamura T, Iwami T, Tanaka R, Kurosawa H, Nitta M, Sobue T. Location of arrest and survival from out-of-hospital cardiac arrest among children in the public-access defibrillation era in Japan. *Resuscitation*. 2019;140:150–158. doi: 10.1016/j.resuscitation.2019.04.045
195. Capucci A, Aschieri D, Piepoli MF, Bardy GH, Iconomu E, Arvedi M. Tripling survival from sudden cardiac arrest via early defibrillation without traditional education in cardiopulmonary resuscitation. *Circulation*. 2002;106:1065–1070. doi: 10.1161/01.cir.0000028148.62305.69
196. Kuisma M, Castrén M, Nurminen K. Public access defibrillation in Helsinki—costs and potential benefits from a community-based pilot study. *Resuscitation*. 2003;56:149–152. doi: 10.1016/s0300-9572(02)00344-1
197. Berdowski J, Blom MT, Bardai A, Tan HL, Tijssen JG, Koster RW. Impact of onsite or dispatched automated external defibrillator use on survival after out-of-hospital cardiac arrest. *Circulation*. 2011;124:2225–2232. doi: 10.1161/CIRCULATIONAHA.110.015545
198. Fordyce CB, Hansen CM, Kragholm K, Dupre ME, Jollis JG, Roettig ML, Becker LB, Hansen SM, Hinohara TT, Corbett CC, Monk L, Nelson RD, Pearson DA, Tyson C, van Diepen S, Anderson ML, McNally B, Granger CB. Association of public health initiatives with outcomes for out-of-hospital cardiac arrest at home and in public locations. *JAMA Cardiol*. 2017;2:1226–1235. doi: 10.1001/jamacardio.2017.3471
199. Andersen LW, Holmberg MJ, Granfeldt A, Løfgren B, Vellano K, McNally BF, Siegerink B, Kurth T, Donnino MW; CARES Surveillance Group. Neighborhood characteristics, bystander automated external defibrillator use, and patient outcomes in public out-of-hospital cardiac arrest. *Resuscitation*. 2018;126:72–79. doi: 10.1016/j.resuscitation.2018.02.021
200. Aschieri D, Penela D, Pelizzoni V, Guerra F, Vermi AC, Rossi L, Torretta L, Losi G, Villani GQ, Capucci A. Outcomes after sudden cardiac arrest in sports centres with and without on-site external defibrillators. *Heart*. 2018;104:1344–1349. doi: 10.1136/heartjnl-2017-312441
201. Pollack RA, Brown SP, Rea T, Aufderheide T, Barbic D, Buick JE, Christenson J, Idris AH, Jasti J, Kamp M, Kudenchuk P, May S, Muhr M, Nichol G, Ornato JP, Sopko G, Vaillancourt C, Morrison L, Weisfeldt M; ROC Investigators. Impact of bystander automated external defibrillator use on survival and functional outcomes in shockable observed public cardiac arrests. *Circulation*. 2018;137:2104–2113. doi: 10.1161/CIRCULATIONAHA.117.030700
202. Ringh M, Jonsson M, Nordberg P, Fredman D, Hasselqvist-Ax I, Håkansson F, Claesson A, Riva G, Hollenberg J. Survival after public access defibrillation in Stockholm, Sweden—a striking success. *Resuscitation*. 2015;91:1–7. doi: 10.1016/j.resuscitation.2015.02.032
203. Hansen SM, Hansen CM, Folke F, Rajan S, Kragholm K, Ejlskov L, Gislason G, Køber L, Gerdts TA, Hjortshøj S, Lippert F, Torp-Pedersen C, Wissenberg M. Bystander defibrillation for out-of-hospital cardiac arrest in public vs residential locations. *JAMA Cardiol*. 2017;2:507–514. doi: 10.1001/jamacardio.2017.0008
204. Claesson A, Herlitz J, Svensson L, Ottosson L, Bergfeldt L, Engdahl J, Ericson C, Sandén P, Axelsson C, Bremer A. Defibrillation before EMS arrival in western Sweden. *Am J Emerg Med*. 2017;35:1043–1048. doi: 10.1016/j.ajem.2017.02.030
205. Dicker B, Davey P, Smith T, Beck B. Incidence and outcomes of out-of-hospital cardiac arrest: a New Zealand perspective. *Emerg Med Australas*. 2018;30:662–671. doi: 10.1111/1742-6723.12966
206. Hallstrom AP, Ornato JP, Weisfeldt M, Travers A, Christenson J, McBurnie MA, Zalenski R, Becker LB, Schron EB, Proschan M; Public Access Defibrillation Trial Investigators. Public-access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med*. 2004;351:637–646. doi: 10.1056/NEJMoa040566
207. Culley LL, Rea TD, Murray JA, Welles B, Fahrenbruch CE, Olsufka M, Eisenberg MS, Copass MK. Public access defibrillation in out-of-hospital cardiac arrest: a community-based study. *Circulation*. 2004;109:1859–1863. doi: 10.1161/01.CIR.0000124721.83385.B2
208. Colquhoun M. Public access defibrillation. *Curr Opin Crit Care*. 2008;14:275–278. doi: 10.1097/MCC.0b013e3282faa748
209. Weisfeldt ML, Sitlani CM, Ornato JP, Rea T, Aufderheide TP, Davis D, Dreyer J, Hess EP, Jui J, Maloney J, Sopko G, Powell J, Nichol G, Morrison LJ; ROC Investigators. Survival after application of automatic external defibrillators before arrival of the emergency medical system: evaluation in the resuscitation outcomes consortium population of 21 million. *J Am Coll Cardiol*. 2010;55:1713–1720. doi: 10.1016/j.jacc.2009.11.077
210. Edwards MJ, Fothergill RT. Exercise-related sudden cardiac arrest in London: incidence, survival and bystander response. *Open Heart*. 2015;2:e000281. doi: 10.1136/openhrt-2015-000281
211. Capucci A, Aschieri D, Guerra F, Pelizzoni V, Nani S, Villani GQ, Bardy GH. Community-based automated external defibrillator only resuscitation for out-of-hospital cardiac arrest patients. *Am Heart J*. 2016;172:192–200. doi: 10.1016/j.ahj.2015.10.018
212. Garcia EL, Caffrey-Villari S, Ramirez D, Caron JL, Mannhart P, Reuter PG, Lapostolle F, Adnet F. [Impact of onsite or dispatched automated external defibrillator use on early survival after sudden cardiac arrest occurring in international airports]. *Presse Med*. 2017;46:e63–e68. doi: 10.1016/j.lpm.2016.09.027
213. Karam N, Marijon E, Dumas F, Offredo L, Beganton F, Bougouin W, Jost D, Lamhaut L, Empana JP, Cariou A, Spaulding C, Jouven X; Paris Sudden Death Expertise Center. Characteristics and outcomes of out-of-hospital sudden cardiac arrest according to the time of occurrence. *Resuscitation*. 2017;116:16–21. doi: 10.1016/j.resuscitation.2017.04.024
214. Deleted in proof.
215. Nas J, Thannhauser J, Herrmann JJ, van der Wulp K, van Grunsven PM, van Royen N, de Boer MJ, Bonnes JL, Brouwer MA. Changes in automated external defibrillator use and survival after out-of-hospital cardiac arrest in the Nijmegen area. *Neth Heart J*. 2018;26:600–605. doi: 10.1007/s12471-018-1162-9
216. Nehme Z, Andrew E, Bernard S, Haskins B, Smith K. Trends in survival from out-of-hospital cardiac arrests defibrillated by paramedics, first responders and bystanders. *Resuscitation*. 2019;143:85–91. doi: 10.1016/j.resuscitation.2019.08.018
217. Fleischhackl R, Roessler B, Domanovits H, Singer F, Fleischhackl S, Foitik G, Czech G, Mittlboeck M, Malzer R, Eisenburger P, Hoerauf K. Results from Austria's nationwide public access defibrillation (ANPAD) programme collected over 2 years. *Resuscitation*. 2008;77:195–200. doi: 10.1016/j.resuscitation.2007.11.019
218. Andersen LW, Holmberg MJ, Granfeldt A, James LP, Cautley L. Cost-effectiveness of public automated external defibrillators. *Resuscitation*. 2019;138:250–258. doi: 10.1016/j.resuscitation.2019.03.029
219. Gong Y, Yu T, Chen B, He M, Li Y. Removal of cardiopulmonary resuscitation artifacts with an enhanced adaptive filtering method: an experimental trial. *Biomed Res Int*. 2014;2014:140438. doi: 10.1155/2014/140438
220. Li Y, Bisera J, Tang W, Weil MH. Automated detection of ventricular fibrillation to guide cardiopulmonary resuscitation. *Crit Pathw Cardiol*. 2007;6:131–134. doi: 10.1097/HPC.0b013e31813429b0
221. Tan Q, Freeman GA, Geheb F, Bisera J. Electrocardiographic analysis during uninterrupted cardiopulmonary resuscitation. *Crit Care Med*. 2008;36(11 Suppl):S409–S412. doi: 10.1097/ccm.0b013e31818a7fbf
222. Werther T, Klotz A, Granegger M, Baubin M, Feichtinger HG, Amann A, Gilly H. Strong corruption of electrocardiograms caused by cardiopulmonary resuscitation reduces efficiency of two-channel methods for removing motion artefacts in non-shockable rhythms. *Resuscitation*. 2009;80:1301–1307. doi: 10.1016/j.resuscitation.2009.07.020
223. Li Y, Bisera J, Weil MH, Tang W. An algorithm used for ventricular fibrillation detection without interrupting chest compression. *IEEE Trans Biomed Eng*. 2012;59:78–86. doi: 10.1109/TBME.2011.2118755
224. Aramendi E, Ayala U, Irueta U, Alonso E, Eftestøl T, Kramer-Johansen J. Suppression of the cardiopulmonary resuscitation artefacts using the instantaneous chest compression rate extracted from the

- thoracic impedance. *Resuscitation*. 2012;83:692–698. doi: 10.1016/j.resuscitation.2011.11.029
225. Babaeizadeh S, Firoozabadi R, Han C, Helfenbein ED. Analyzing cardiac rhythm in the presence of chest compression artifact for automated shock advisory. *J Electrocardiol*. 2014;47:798–803. doi: 10.1016/j.jelectrocard.2014.07.021
  226. Gong Y, Gao P, Wei L, Dai C, Zhang L, Li Y. An Enhanced Adaptive Filtering Method for Suppressing Cardiopulmonary Resuscitation Artifact. *IEEE Trans Biomed Eng*. 2017;64:471–478. doi: 10.1109/TBME.2016.2564642
  227. Fumagalli F, Silver AE, Tan Q, Zaidi N, Ristagno G. Cardiac rhythm analysis during ongoing cardiopulmonary resuscitation using the Analysis During Compressions with Fast Reconfirmation technology. *Heart Rhythm*. 2018;15:248–255. doi: 10.1016/j.hrthm.2017.09.003
  228. Hu Y, Tang H, Liu C, Jing D, Zhu H, Zhang Y, Yu X, Zhang G, Xu J. The performance of a new shock advisory algorithm to reduce interruptions during CPR. *Resuscitation*. 2019;143:1–9. doi: 10.1016/j.resuscitation.2019.07.026
  229. Zhang G, Wu T, Wan Z, Song Z, Yu M, Wang D, Li L, Chen F. A new method to detect ventricular fibrillation from CPR artifact-corrupted ECG based on the ECG alone. *Biomedical Signal Processing and Control* 2016;29:67–75.
  230. Rad AB, Engan K, Katsaggelos AK, Kvaløy JT, Wik L, Kramer-Johansen J, Irueta U, Eftestøl T. Automatic cardiac rhythm interpretation during resuscitation. *Resuscitation*. 2016;102:44–50. doi: 10.1016/j.resuscitation.2016.01.015
  231. Zhang G, Wu T, Wan Z, Song Z, Yu M, Wang D, Li L, Chen F, Xu X. A method to differentiate between ventricular fibrillation and asystole during chest compressions using artifact-corrupted ECG alone. *Comput Methods Programs Biomed*. 2017;141:111–117. doi: 10.1016/j.cmpb.2017.01.015
  232. Partridge R, Tan Q, Silver A, Riley M, Geheb F, Raymond R. Rhythm analysis and charging during chest compressions reduces compression pause time. *Resuscitation*. 2015;90:133–137. doi: 10.1016/j.resuscitation.2015.02.025
  233. Jacobs IG, Finn JC, Oxer HF, Jelinek GA. CPR before defibrillation in out-of-hospital cardiac arrest: a randomized trial. *Emerg Med Australas*. 2005;17:39–45. doi: 10.1111/j.1742-6723.2005.00694.x
  234. Ma MH, Chiang WC, Ko PC, Yang CW, Wang HC, Chen SY, Chang WT, Huang CH, Chou HC, Lai MS, Chien KL, Lee BC, Hwang CH, Wang YC, Hsiung GH, Hsiao YW, Chang AM, Chen WJ, Chen SC. A randomized trial of compression first or analyze first strategies in patients with out-of-hospital cardiac arrest: results from an Asian community. *Resuscitation*. 2012;83:806–812. doi: 10.1016/j.resuscitation.2012.01.009
  235. Stiell IG, Nichol G, Leroux BG, Rea TD, Ornato JP, Powell J, Christenson J, Callaway CW, Kudenchuk PJ, Aufderheide TP, Idris AH, Daya MR, Wang HE, Morrison LJ, Davis D, Andrusiek D, Stephens S, Cheskes S, Schmicker RH, Fowler R, Vaillancourt C, Hostler D, Zive D, Pirrallo RG, Vilke GM, Sopko G, Weisfeldt M; ROC Investigators. Early versus later rhythm analysis in patients with out-of-hospital cardiac arrest. *N Engl J Med*. 2011;365:787–797. doi: 10.1056/NEJMoa1010076
  236. Rea T, Prince D, Morrison L, Callaway C, Aufderheide T, Daya M, Stiell I, Christenson J, Powell J, Warden C, van Ottingham L, Kudenchuk P, Weisfeldt M. Association between survival and early versus later rhythm analysis in out-of-hospital cardiac arrest: do agency-level factors influence outcomes? *Ann Emerg Med*. 2014;64:1–8. doi: 10.1016/j.annemergmed.2014.01.014
  237. Sunde K, Jacobs I, Deakin CD, Hazinski MF, Kerber RE, Koster RW, Morrison LJ, Nolan JP, Sayre MR; on behalf of the Defibrillation Chapter Collaborators. Part 6: defibrillation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2010;81(suppl 1):e71–85. doi: 10.1016/j.resuscitation.2010.08.025
  238. Jacobs I, Sunde K, Deakin CD, Hazinski MF, Kerber RE, Koster RW, Morrison LJ, Nolan JP, Sayre MR; on behalf of the Defibrillation Chapter Collaborators. Part 6: Defibrillation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122 (Suppl 2):S325–337. doi: 10.1161/CIRCULATIONAHA.110.971010
  239. Drennan IR, Lin S, Sidalak DE, Morrison LJ. Survival rates in out-of-hospital cardiac arrest patients transported without prehospital return of spontaneous circulation: an observational cohort study. *Resuscitation*. 2014;85:1488–1493. doi: 10.1016/j.resuscitation.2014.07.011
  240. Gray WA, Capone RJ, Most AS. Unsuccessful emergency medical resuscitation—are continued efforts in the emergency department justified? *N Engl J Med*. 1991;325:1393–1398. doi: 10.1056/NEJM199111143252001
  241. Lewis LM, Ruoff B, Rush C, Stothert JC Jr. Is emergency department resuscitation of out-of-hospital cardiac arrest victims who arrive pulseless worthwhile? *Am J Emerg Med*. 1990;8:118–120. doi: 10.1016/0735-6757(90)90196-7
  242. Lim GH, Seow E. Resuscitation for patients with out-of-hospital cardiac arrest: Singapore. *Prehosp Disaster Med*. 2002;17:96–101. doi: 10.1017/s1049023x00000248
  243. Schoenenberger RA, von Planta M, von Planta I. Survival after failed out-of-hospital resuscitation. Are further therapeutic efforts in the emergency department futile? *Arch Intern Med*. 1994;154:2433–2437.
  244. Yates EJ, Smyth MA, Perkins GD. Reply to: Letter to the Editor re out-of-hospital cardiac arrest termination of resuscitation with ongoing CPR: An observational study. *Resuscitation*. 2018;133:e10. doi: 10.1016/j.resuscitation.2018.10.013
  245. Kellermann AL, Hackman BB, Somes G. Predicting the outcome of unsuccessful prehospital advanced cardiac life support. *JAMA*. 1993;270:1433–1436.
  246. Zive D, Koprowicz K, Schmidt T, Stiell I, Sears G, Van Ottingham L, Idris A, Stephens S, Daya M; Resuscitation Outcomes Consortium Investigators. Variation in out-of-hospital cardiac arrest resuscitation and transport practices in the Resuscitation Outcomes Consortium: ROC Epistry-Cardiac Arrest. *Resuscitation*. 2011;82:277–284. doi: 10.1016/j.resuscitation.2010.10.022
  247. Cheskes S, Byers A, Zhan C, Verbeek PR, Ko D, Drennan IR, Buick JE, Brooks SC, Lin S, Taher A, Morrison LJ; Rescu Epistry Investigators. CPR quality during out-of-hospital cardiac arrest transport. *Resuscitation*. 2017;114:34–39. doi: 10.1016/j.resuscitation.2017.02.016
  248. Ødegaard S, Olasveengen T, Steen PA, Kramer-Johansen J. The effect of transport on quality of cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Resuscitation*. 2009;80:843–848. doi: 10.1016/j.resuscitation.2009.03.032
  249. Olasveengen TM, Wik L, Steen PA. Quality of cardiopulmonary resuscitation before and during transport in out-of-hospital cardiac arrest. *Resuscitation*. 2008;76:185–190. doi: 10.1016/j.resuscitation.2007.07.001
  250. Roosa JR, Vadeboncoeur TF, Dommer PB, Panchal AR, Venuti M, Smith G, Silver A, Mullins M, Spate D, Bobrow BJ. CPR variability during ground ambulance transport of patients in cardiac arrest. *Resuscitation*. 2013;84:592–595. doi: 10.1016/j.resuscitation.2012.07.042
  251. Russi CS, Myers LA, Kolb LJ, Lohse CM, Hess EP, White RD. A Comparison of Chest Compression Quality Delivered During On-Scene and Ground Transport Cardiopulmonary Resuscitation. *West J Emerg Med*. 2016;17:634–639. doi: 10.5811/westjem.2016.6.29949
  252. Braunfels S, Meinhard K, Zieher B, Koetter KP, Maleck WH, Petroianu GA. A randomized, controlled trial of the efficacy of closed chest compressions in ambulances. *Prehosp Emerg Care*. 1997;1:128–131. doi: 10.1080/10903129708958804
  253. Havel C, Schreiber W, Trimmel H, Malzer R, Haugk M, Richling N, Riedmüller E, Sterz F, Herkner H. Quality of closed chest compression on a manikin in ambulance vehicles and flying helicopters with a real time automated feedback. *Resuscitation*. 2010;81:59–64. doi: 10.1016/j.resuscitation.2009.10.007
  254. Lipman SS, Wong JY, Arafeh J, Cohen SE, Carvalho B. Transport decreases the quality of cardiopulmonary resuscitation during simulated maternal cardiac arrest. *Anesth Analg*. 2013;116:162–167. doi: 10.1213/ANE.0b013e31826dd889
  255. Sunde K, Wik L, Steen PA. Quality of mechanical, manual standard and active compression-decompression CPR on the arrest site and during transport in a manikin model. *Resuscitation*. 1997;34:235–242. doi: 10.1016/s0300-9572(96)01087-8
  256. Chung TN, Kim SW, Cho YS, Chung SP, Park I, Kim SH. Effect of vehicle speed on the quality of closed-chest compression during ambulance transport. *Resuscitation*. 2010;81:841–847. doi: 10.1016/j.resuscitation.2010.02.024
  257. Roberts BG. Machine vs. manual cardiopulmonary resuscitation in moving vehicles. *EMT J*. 1979;3:30–34.
  258. Stone CK, Thomas SH. Can correct closed-chest compressions be performed during prehospital transport? *Prehosp Disaster Med*. 1995;10:121–123. doi: 10.1017/s1049023x00041856
  259. Stapleton ER. Comparing CPR during ambulance transport. Manual vs. mechanical methods. *JEMS*. 1991;16:63–4, 66, 68 passim.
  260. Axelsson C, Karlsson T, Axelsson AB, Herlitz J. Mechanical active compression-decompression cardiopulmonary resuscitation (ACD-CPR) versus manual CPR according to pressure of end tidal carbon dioxide (PetCO<sub>2</sub>) during CPR in out-of-hospital cardiac arrest (OHCA). *Resuscitation*. 2009;80:1099–1103. doi: 10.1016/j.resuscitation.2009.08.006
  261. Dickinson ET, Verdile VP, Schneider RM, Salluzzo RF. Effectiveness of mechanical versus manual chest compressions in out-of-hospital cardiac



- arrest resuscitation: a pilot study. *Am J Emerg Med*. 1998;16:289–292. doi: 10.1016/s0735-6757(98)90105-x
262. Hallstrom A, Rea TD, Sayre MR, Christenson J, Anton AR, Mosesso VN Jr, Van Ottingham L, Olsufka M, Pennington S, White LJ, Yahn S, Husar J, Morris MF, Cobb LA. Manual chest compression vs use of an automated chest compression device during resuscitation following out-of-hospital cardiac arrest: a randomized trial. *JAMA*. 2006;295:2620–2628. doi: 10.1001/jama.295.22.2620
  263. Axelsson C, Nestin J, Svensson L, Axelsson AB, Herlitz J. Clinical consequences of the introduction of mechanical chest compression in the EMS system for treatment of out-of-hospital cardiac arrest—a pilot study. *Resuscitation*. 2006;71:47–55. doi: 10.1016/j.resuscitation.2006.02.011
  264. Casner M, Andersen D, Isaacs SM. The impact of a new CPR assist device on rate of return of spontaneous circulation in out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2005;9:61–67. doi: 10.1080/109031205090891714
  265. Ong ME, Ornato JP, Edwards DP, Dhindsa HS, Best AM, Ines CS, Hickey S, Clark B, Williams DC, Powell RG, Overton JL, Peberdy MA. Use of an automated, load-distributing band chest compression device for out-of-hospital cardiac arrest resuscitation. *JAMA*. 2006;295:2629–2637. doi: 10.1001/jama.295.22.2629
  266. Kim TH, Shin SD, Song KJ, Hong KJ, Ro YS, Song SW, Kim CH. Chest compression fraction between mechanical compressions on a reducible stretcher and manual compressions on a standard stretcher during transport in out-of-hospital cardiac arrests: the ambulance stretcher innovation of asian cardiopulmonary resuscitation (ASIA-CPR) Pilot Trial. *Prehosp Emerg Care*. 2017;21:636–644. doi: 10.1080/10903127.2017.1317892
  267. Lyon RM, Crawford A, Crookston C, Short S, Clegg GR. The combined use of mechanical CPR and a carry sheet to maintain quality resuscitation in out-of-hospital cardiac arrest patients during extrication and transport. *Resuscitation*. 2015;93:102–106. doi: 10.1016/j.resuscitation.2015.05.030
  268. Wang HC, Chiang WC, Chen SY, Ke YL, Chi CL, Yang CW, Lin PC, Ko PC, Wang YC, Tsai TC, Huang CH, Hsiung KH, Ma MH, Chen SC, Chen WJ, Lin FY. Video-recording and time-motion analyses of manual versus mechanical cardiopulmonary resuscitation during ambulance transport. *Resuscitation*. 2007;74:453–460. doi: 10.1016/j.resuscitation.2007.01.018
  269. Kim TH, Hong KJ, Sang Do S, Kim CH, Song SW, Song KJ, Ro YS, Ahn KO, Jang DB. Quality between mechanical compression on reducible stretcher versus manual compression on standard stretcher in small elevator. *Am J Emerg Med*. 2016;34:1604–1609. doi: 10.1016/j.ajem.2016.05.072
  270. Putzer G, Braun P, Zimmermann A, Pedross F, Strapazon G, Brugger H, Paal P. LUCAS compared to manual cardiopulmonary resuscitation is more effective during helicopter rescue—a prospective, randomized, cross-over manikin study. *Am J Emerg Med*. 2013;31:384–389. doi: 10.1016/j.ajem.2012.07.018
  271. Fox J, Fiechter R, Gerstl P, Url A, Wagner H, Lüscher TF, Eriksson U, Wyss CA. Mechanical versus manual chest compression CPR under ground ambulance transport conditions. *Acute Card Care*. 2013;15:1–6. doi: 10.3109/17482941.2012.735675
  272. Gässler H, Ventzke MM, Lampl L, Helm M. Transport with ongoing resuscitation: a comparison between manual and mechanical compression. *Emerg Med J*. 2013;30:589–592. doi: 10.1136/emermed-2012-201142
  273. Cudnik MT, Schmicker RH, Vaillancourt C, Newgard CD, Christensen JM, Davis DP, Lowe RA; ROC Investigators. A geospatial assessment of transport distance and survival to discharge in out of hospital cardiac arrest patients: Implications for resuscitation centers. *Resuscitation*. 2010;81:518–523. doi: 10.1016/j.resuscitation.2009.12.030
  274. Geri G, Gilgan J, Wu W, Vijendira S, Ziegler C, Drennan IR, Morrison L, Lin S. Does transport time of out-of-hospital cardiac arrest patients matter? A systematic review and meta-analysis. *Resuscitation*. 2017;115:96–101. doi: 10.1016/j.resuscitation.2017.04.003
  275. Spaite DW, Bobrow BJ, Vadeboncoeur TF, Chikani V, Clark L, Mullins T, Sanders AB. The impact of prehospital transport interval on survival in out-of-hospital cardiac arrest: implications for regionalization of post-resuscitation care. *Resuscitation*. 2008;79:61–66. doi: 10.1016/j.resuscitation.2008.05.006
  276. Spaite DW, Stiell IG, Bobrow BJ, de Boer M, Maloney J, Denninghoff K, Vadeboncoeur TF, Dreyer J, Wells GA. Effect of transport interval on out-of-hospital cardiac arrest survival in the OPALS study: implications for triaging patients to specialized cardiac arrest centers. *Ann Emerg Med*. 2009;54:248–255. doi: 10.1016/j.annemergmed.2008.11.020
  277. Vilke GM, Smith AM, Ray LU, Steen PJ, Murrin PA, Chan TC. Airway obstruction in children aged less than 5 years: the prehospital experience. *Prehosp Emerg Care*. 2004;8:196–199. doi: 10.1016/j.prehos.2003.12.014
  278. Igarashi Y, Yokobori S, Yoshino Y, Masuno T, Miyauchi M, Yokota H. Pre-hospital removal improves neurological outcomes in elderly patient with foreign body airway obstruction. *Am J Emerg Med*. 2017;35:1396–1399. doi: 10.1016/j.ajem.2017.04.016
  279. Redding JS. The choking controversy: critique of evidence on the Heimlich maneuver. *Crit Care Med*. 1979;7:475–479.
  280. Guinane J, Lee SM. Fatal acute aortic dissection after back blows and chest thrusts delivered for choking episode. *Intern Med J*. 2018;48:1272–1273. doi: 10.1111/imj.14062
  281. Nowitz A, Lewer BM, Galletly DC. An interesting complication of the Heimlich manoeuvre. *Resuscitation*. 1998;39:129–131. doi: 10.1016/s0300-9572(98)00109-9
  282. Patterson DL, Brennan S, Cartwright T, Jolly W, Adlam JH, Waller BF. Traumatic rupture of an aortic ulcerative atherosclerotic plaque producing aortic dissection: a complication of interscapular back blows used to dislodge objects from the esophagus. *Clin Cardiol*. 1993;16:741–744. doi: 10.1002/clc.4960161011
  - 282a. Rakotoharinandrasana H, Petit E, Dumas P, Vandermarcq P, Gil R, Neau JP. Internal carotid artery dissection after Heimlich maneuver. *Ann Fr Anesth Reanim*. 2003;22:43–45. doi: 10.1016/s0750-7658(02)00007-2
  283. Boussuges S, Maitre Robert P, Bost M. [Use of the Heimlich Maneuver on children in the Rhône-Alpes area]. *Arch Fr Pediatr*. 1985;42:733–736.
  284. Heimlich HJ. A life-saving maneuver to prevent food-choking. *JAMA*. 1975;234:398–401.
  285. Soroudi A, Shipp HE, Stepanski BM, Ray LU, Murrin PA, Chan TC, Davis DP, Vilke GM. Adult foreign body airway obstruction in the prehospital setting. *Prehosp Emerg Care*. 2007;11:25–29. doi: 10.1080/10903120601023263
  286. Agia GA, Hurst DJ. Pneumomediastinum following the Heimlich maneuver. *JACEP*. 1979;8:473–475. doi: 10.1016/s0361-1124(79)80064-7
  287. Ayerdi J, Gupta SK, Sampson LN, Deshmukh N. Acute abdominal aortic thrombosis following the Heimlich maneuver. *Cardiovasc Surg*. 2002;10:154–156. doi: 10.1016/s0967-2109(01)00017-5
  288. Bintz M, Cogbill TH. Gastric rupture after the Heimlich maneuver. *J Trauma*. 1996;40:159–160. doi: 10.1097/00005373-199601000-00036
  289. Bouayed S, Sandu K, Teiga PS, Hallak B. Thoracocervicofacial emphysema after Heimlich's maneuver. *Case Rep Otolaryngol*. 2015;2015:427320. doi: 10.1155/2015/427320
  290. Caro Aguilera P, Peiró Aranda R, Pérez Ruiz E, Rodríguez Amuedo F, Pérez Frías J. [Haemoptysis after Heimlich manoeuvre]. *An Pediatr (Barc)*. 2008;68:533–534. doi: 10.1157/13120211
  291. Cecchetto G, Viel G, Cecchetto A, Kusstatscher S, Montisci M. Fatal splenic rupture following Heimlich maneuver: case report and literature review. *Am J Forensic Med Pathol*. 2011;32:169–171. doi: 10.1097/PAF.0b013e318219c878
  292. Chao CM, Lai CC, Tan CK. Gastric perforation after Heimlich maneuver. *Am J Med*. 2012;125:e7–e8. doi: 10.1016/j.amjmed.2012.02.008
  293. Chapman JH, Menapace FJ, Howell RR. Ruptured aortic valve cusp: a complication of the Heimlich maneuver. *Ann Emerg Med*. 1983;12:446–448. doi: 10.1016/s0196-0644(83)80344-8
  294. Chillag S, Krieg J, Bhargava R. The Heimlich maneuver: breaking down the complications. *South Med J*. 2010;103:147–150. doi: 10.1097/SMJ.0b013e3181c99140
  295. Cowan M, Bardole J, Dlesk A. Perforated stomach following the Heimlich maneuver. *Am J Emerg Med*. 1987;5:121–122. doi: 10.1016/0735-6757(87)90088-x
  296. Croom DW. Rupture of stomach after attempted Heimlich maneuver. *JAMA*. 1983;250:2602–2603. doi: 10.1001/jama.1983.03340190016011
  297. Desai SC, Chute DJ, Desai BC, Koloski ER. Traumatic dissection and rupture of the abdominal aorta as a complication of the Heimlich maneuver. *J Vasc Surg*. 2008;48:1325–1327. doi: 10.1016/j.jvs.2008.05.061
  298. Dupre MW, Silva E, Brotman S. Traumatic rupture of the stomach secondary to Heimlich maneuver. *Am J Emerg Med*. 1993;11:611–612. doi: 10.1016/0735-6757(93)90014-3
  299. Entel RJ, Hakki AH. Bone scan after the Heimlich maneuver. *Clin Nucl Med*. 1996;21:251. doi: 10.1097/00003072-199603000-00016
  300. Fearing NM, Harrison PB. Complications of the Heimlich maneuver: case report and literature review. *J Trauma*. 2002;53:978–979. doi: 10.1097/00005373-200211000-00026
  301. Feeney SN, Pegoli W, Gestring ML. Pancreatic transection as a complication of the Heimlich maneuver: case report and literature review. *J Trauma*. 2007;62:252–254. doi: 10.1097/TA.0b013e31802e2d3b
  302. Fink JA, Klein RL. Complications of the Heimlich maneuver. *J Pediatr Surg*. 1989;24:486–487. doi: 10.1016/s0022-3468(89)80408-7



303. Haynes DE, Haynes BE, Yong YV. Esophageal rupture complicating Heimlich maneuver. *Am J Emerg Med.* 1984;2:507–509. doi: 10.1016/0735-6757(84)90075-5
304. Herman A, Maiti A, Cherian SV, Estrada-Y-Martin RM. Heimlich Maneuver-Induced Diaphragmatic Rupture and Hiatal Hernia. *Am J Med Sci.* 2018;355:e13. doi: 10.1016/j.amjms.2017.11.009
305. Kirshner RL, Green RM. Acute thrombosis of abdominal aortic aneurysm subsequent to Heimlich maneuver: a case report. *J Vasc Surg.* 1985;2:594–596. doi: 10.1067/mva.1985.avs0020594
306. Koss SL, Karle WE, Dibelius G, Kamat A, Berzofsky C. Esophageal perforation as a complication of the Heimlich maneuver in a pediatric patient: A case report. *Ear Nose Throat J.* 2018;97:E1–E3. doi: 10.1177/014556131809700701
307. Kösser A, Lehmkuhl L, Gutberlet M. [Aortocaval fistula after administering the Heimlich maneuver—diagnosis with multislice CT]. *Rofo.* 2009;181:1089–1090. doi: 10.1055/s-0028-1109780
308. Lee KY, Wu YL, Ho SW. Silent Aortic Dissection after the Heimlich Maneuver: A Case Report. *J Emerg Med.* 2019;56:210–212. doi: 10.1016/j.jemermed.2018.09.053
309. Lee SL, Kim SS, Shekherdian S, Ledbetter DJ. Complications as a result of the Heimlich maneuver. *J Trauma.* 2009;66:E34–E35. doi: 10.1097/01.ta.0000219291.27245.90
310. Lette J, Levasseur A, Labonté C, Eybalin MC, Cerino M. Thoracic bone imaging after the Heimlich maneuver. *Clin Nucl Med.* 1990;15:512. doi: 10.1097/00003072-199007000-00014
311. Lin PH, Bush RL, Lumsden AB. Proximal aortic stent-graft displacement with type I endoleak due to Heimlich maneuver. *J Vasc Surg.* 2003;38:380–382. doi: 10.1016/s0741-5214(03)00294-5
312. Mack L, Forbes TL, Harris KA. Acute aortic thrombosis following incorrect application of the Heimlich maneuver. *Ann Vasc Surg.* 2002;16:130–133. doi: 10.1007/s10016-001-0147-z
313. Majumdar A, Sedman PC. Gastric rupture secondary to successful Heimlich manoeuvre. *Postgrad Med J.* 1998;74:609–610. doi: 10.1136/pgmj.74.876.609
314. Martin TJ, Bobba RK, Metzger R, Madalina M, Bollu M, Patel BG, Kazemi MM. Acute abdominal aortic thrombosis as a complication of the Heimlich maneuver. *J Am Geriatr Soc.* 2007;55:1146–1147. doi: 10.1111/j.1532-5415.2007.01221.x
315. Matharoo G, Kalra A, Phatak T, Bhattacharyya N. Diaphragmatic rupture with gastric volvulus after Heimlich maneuver. *Eur J Pediatr Surg.* 2013;23:502–504. doi: 10.1055/s-0032-1330850
316. Meredith MJ, Liebowitz R. Rupture of the esophagus caused by the Heimlich maneuver. *Ann Emerg Med.* 1986;15:106–107. doi: 10.1016/s0196-0644(86)80537-6
317. Olenchok SA Jr, Rowlands DM, Reed JF III, Garzia FM, Zasik JM. Dysphagia after Heimlich maneuver. *Chest.* 2004;125:302–304. doi: 10.1378/chest.125.1.302
318. Otero Palleiro MM, Barbagelata López C, Fernández Pretel MC, Salgado Fernández J. Hepatic rupture after Heimlich maneuver. *Ann Emerg Med.* 2007;49:825–826. doi: 10.1016/j.annemergmed.2006.12.019
319. Passik CS, Ackermann DM, Piehler JM, Edwards WD. Traumatic rupture of Ionescu-Shiley aortic valve after the Heimlich maneuver. *Arch Pathol Lab Med.* 1987;111:469–470.
320. Razaboni RM, Brathwaite CE, Dwyer WA Jr. Ruptured jejunum following Heimlich maneuver. *J Emerg Med.* 1986;4:95–98. doi: 10.1016/0736-4679(86)90070-3
321. Roehm EF, Twiest MW, Williams RC Jr. Abdominal aortic thrombosis in association with an attempted Heimlich maneuver. *JAMA.* 1983;249:1186–1187.
322. Sams JS. Dangers of the Heimlich maneuver for esophageal obstruction. *N Engl J Med.* 1989;321:980–981.
323. Skulberg A. Chest compression—an alternative to the Heimlich manoeuvre? *Resuscitation.* 1992;24:91. doi: 10.1016/0300-9572(92)90177-e
324. Tashtoush B, Schroeder J, Memarpour R, Oliveira E, Medina M, Hadeh A, Ramirez J, Smolley L. Food Particle Aspiration Associated with Hemorrhagic Shock: A Diagnostic Dilemma. *Case Rep Emerg Med.* 2015;2:75497. doi: 10.1155/2015/275497
325. Truong T, Salire K, De Cicco I, Cherian S, Aisenberg G. Incarcerated diaphragmatic hernia following Heimlich maneuver. *Proc (Bayl Univ Med Cent).* 2018;31:48–50. doi: 10.1080/08998280.2017.1391034
326. Tung PH, Law S, Chu KM, Law WL, Wong J. Gastric rupture after Heimlich maneuver and cardiopulmonary resuscitation. *Hepatogastroenterology.* 2001;48:109–111.
327. Ujjin V, Ratanasit S, Nagendran T. Diaphragmatic hernia as a complication of the Heimlich maneuver. *Int Surg.* 1984;69:175–176.
328. Valero V. Mesenteric laceration complicating a Heimlich maneuver. *Ann Emerg Med.* 1986;15:105–106. doi: 10.1016/s0196-0644(86)80536-4
329. van der Ham AC, Lange JF. Traumatic rupture of the stomach after Heimlich maneuver. *J Emerg Med.* 1990;8:713–715. doi: 10.1016/0736-4679(90)90284-3
330. Visintine RE, Baick CH. Ruptured stomach after Heimlich maneuver. *JAMA.* 1975;234:415.
331. Wolf DA. Heimlich trauma: a violent maneuver. *Am J Forensic Med Pathol.* 2001;22:65–67. doi: 10.1097/0000433-200103000-00013
332. Rakotoharinandrasana H, Petit E, Dumas P, Vandermarcq P, Gil R, Neau JP. [Internal carotid artery dissection after Heimlich maneuver]. *Ann Fr Anesth Reanim.* 2003;22:43–45. doi: 10.1016/s0750-7658(02)00007-2
333. Ülger H. Complications of the Heimlich maneuver: isolated sternum fracture. *Akademik Acil Tip Olgu Sunumlari Dergisi.* 2016;7:15–16.
334. Kinoshita K, Azuhata T, Kawano D, Kawahara Y. Relationships between pre-hospital characteristics and outcome in victims of foreign body airway obstruction during meals. *Resuscitation.* 2015;88:63–67. doi: 10.1016/j.resuscitation.2014.12.018
335. Abder-Rahman HA. Infants choking following blind finger sweep. *J Pediatr (Rio J).* 2009;85:273–275. doi: 10.2223/JPED.1892
336. Gjoni D, Mbamalu D, Banerjee A, James K. An unusual complication of an attempt to open the airway in a choking child. *Br J Hosp Med (Lond).* 2009;70:595. doi: 10.12968/hmed.2009.70.10.44630
337. Hartrey R, Bingham RM. Pharyngeal trauma as a result of blind finger sweeps in the choking child. *J Accid Emerg Med.* 1995;12:52–54. doi: 10.1136/emj.12.1.52
338. Heimlich HJ. Update on the Heimlich Maneuver. *Emerg Med Serv.* 1977;6:11–3, 16.
339. Kabbani M, Goodwin SR. Traumatic epiglottitis following blind finger sweep to remove a pharyngeal foreign body. *Clin Pediatr (Phila).* 1995;34:495–497. doi: 10.1177/000992289503400908
340. Mori T, Inoue N. Nasopharyngeal foreign body triggered by a blind finger sweep. *BMJ Case Rep.* 2016;2016:bcr2016216536. doi: 10.1136/bcr-2016-216536
341. Sridharan S, Amin MR, Branski RC. Vocal fold immobility after finger sweep self-extrusion of impacted food in a choking victim with resolution following laryngeal mask ventilation. *Ear Nose Throat J.* 2016;95:33–35.
342. Vunda A, Vandertuin L. Nasopharyngeal foreign body following a blind finger sweep. *J Pediatr.* 2012;160:353. doi: 10.1016/j.jpeds.2011.08.061
343. Sakai T, Kitamura T, Iwami T, Nishiyama C, Tanigawa-Sugihara K, Hayashida S, Nishiuchi T, Kajino K, Irisawa T, Shiozaki T, Ogura H, Tasaki O, Kuwagata Y, Hiraide A, Shimazu T. Effectiveness of prehospital Magill forceps use for out-of-hospital cardiac arrest due to foreign body airway obstruction in Osaka City. *Scand J Trauma Resusc Emerg Med.* 2014;22:53. doi: 10.1186/s13049-014-0053-3
344. Rouillon I, Charrier JB, Devictor D, Portier F, Lebret IK, Attal P, Le Pajolec C, Bobin S. Lower respiratory tract foreign bodies: a retrospective review of morbidity, mortality and first aid management. *Int J Pediatr Otorhinolaryngol.* 2006;70:1949–1955. doi: 10.1016/j.ijporl.2006.07.022
345. Saperstein DM, Pugliese PR, Ulteig C, Schreiber N. Successful use of a novel device called the Lifevac ton resuscitate choking victims world-wide results. 2018;12 doi: DOI: 10.4172/Clinical-Skills.1000136
- 345a. Centers for Disease Control and Prevention. CDC's response to the opioid overdose epidemic. <http://www.cdc.gov/opioids>. Accessed August 12, 2020.
346. Giglio RE, Li G, DiMaggio CJ. Effectiveness of bystander naloxone administration and overdose education programs: a meta-analysis. *Inj Epidemiol.* 2015;2:10. doi: 10.1186/s40621-015-0041-8
347. McDonald R, Strang J. Are take-home naloxone programmes effective? Systematic review utilizing application of the Bradford Hill criteria. *Addiction.* 2016;111:1177–1187. doi: 10.1111/add.13326
348. Wermeling DP. Review of naloxone safety for opioid overdose: practical considerations for new technology and expanded public access. *Ther Adv Drug Saf.* 2015;6:20–31. doi: 10.1177/2042098614564776
349. World Health Organization. Drowning. 2020. <https://www.who.int/news-room/fact-sheets/detail/drowning>. Accessed March 15, 2020.
350. Frates RC Jr. Analysis of predictive factors in the assessment of warm-water near-drowning in children. *Am J Dis Child.* 1981;135:1006–1008. doi: 10.1001/archpedi.1981.02130350010004
351. Nagel FO, Kibel SM, Beatty DW. Childhood near-drowning—factors associated with poor outcome. *S Afr Med J.* 1990;78:422–425.
352. Quan L, Wentz KR, Gore EJ, Copass MK. Outcome and predictors of outcome in pediatric submersion victims receiving prehospital care in King County, Washington. *Pediatrics.* 1990;86:586–593.

353. Niu YW, Chong WS, Lin MT, Tsao LY. An analysis of prognostic factors for submersion accidents in children. *Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi*. 1992;33:81–88.
354. Mizuta R, Fujita H, Osamura T, Kidowaki T, Kiyosawa N. Childhood drownings and near-drownings in Japan. *Acta Paediatr Jpn*. 1993;35:186–192. doi: 10.1111/j.1442-200x.1993.tb03036.x
355. Kyriacou DN, Arciniegas EL, Peek C, Kraus JF. Effect of immediate resuscitation on children with submersion injury. *Pediatrics*. 1994;94(2 Pt 1):137–142.
356. Al-Mofadda SM, Nassar A, Al-Turki A, Al-Salloun AA. Pediatric near drowning: the experience of King Khalid University Hospital. *Ann Saudi Med*. 2001;21:300–303. doi: 10.5144/0256-4947.2001.300
357. Blasco Alonso J, Moreno Pérez D, Milano Manso G, Calvo Macías C, Jurado Ortiz A. [Drowning in pediatric patients]. *An Pediatr (Barc)*. 2005;62:20–24. doi: 10.1157/13070176
358. Nitta M, Kitamura T, Iwami T, Nadkarni VM, Berg RA, Topjian AA, Okamoto Y, Nishiyama C, Nishuchi T, Hayashi Y, Nishimoto Y, Takasu A. Out-of-hospital cardiac arrest due to drowning among children and adults from the Utstein Osaka Project. *Resuscitation*. 2013;84:1568–1573. doi: 10.1016/j.resuscitation.2013.06.017
359. Quan L, Mack CD, Schiff MA. Association of water temperature and submersion duration and drowning outcome. *Resuscitation*. 2014;85:790–794. doi: 10.1016/j.resuscitation.2014.02.024
- 359a. Al-Qurashi FO, Yousef AA, Aljoudi A, Alzahrani SM, Al-Jawder NY, Al-Ahmar AK, Al-Majed MS, Abouollo HM. A review of nonfatal drowning in the pediatric-age group. *Pediatr Emerg Care*. 2019;35:782–786. doi: 10.1097/PEC.0000000000001232
360. Orlowski JP. Prognostic factors in pediatric cases of drowning and near-drowning. *JACEP*. 1979;8:176–179. doi: 10.1016/s0361-1124(79)80121-5
361. Mosayebi Z, Movahedian AH, Mousavi GA. Drowning in children in Iran: outcomes and prognostic factors. *Med J Malaysia*. 2011;66:187–190.
362. Claesson A, Lindqvist J, Ortenwall P, Herlitz J. Characteristics of life-saving from drowning as reported by the Swedish Fire and Rescue Services 1996–2010. *Resuscitation*. 2012;83:1072–1077. doi: 10.1016/j.resuscitation.2012.05.025
363. Dyson K, Morgans A, Bray J, Matthews B, Smith K. Drowning related out-of-hospital cardiac arrests: characteristics and outcomes. *Resuscitation*. 2013;84:1114–1118. doi: 10.1016/j.resuscitation.2013.01.020
364. Claesson A, Lindqvist J, Herlitz J. Cardiac arrest due to drowning—changes over time and factors of importance for survival. *Resuscitation*. 2014;85:644–648. doi: 10.1016/j.resuscitation.2014.02.006
365. Vähätalo R, Lunetta P, Olkkola KT, Suominen PK. Drowning in children: Utstein style reporting and outcome. *Acta Anaesthesiol Scand*. 2014;58:604–610. doi: 10.1111/aas.12298
366. Claesson A, Svensson L, Silfverstolpe J, Herlitz J. Characteristics and outcome among patients suffering out-of-hospital cardiac arrest due to drowning. *Resuscitation*. 2008;76:381–387. doi: 10.1016/j.resuscitation.2007.09.003
367. Forler J, Carsin A, Arlaud K, Bosdure E, Viard L, Paut O, Camboulives J, Dubus JC. [Respiratory complications of accidental drownings in children]. *Arch Pediatr*. 2010;17:14–18. doi: 10.1016/j.arcped.2009.09.021
368. Jeong J, Hong KJ, Shin SD, Ro YS, Song KJ, Lee EJ, Lee YJ, Ahn KO. Relationship between drowning location and outcome after drowning-associated out-of-hospital cardiac arrest: nationwide study. *Am J Emerg Med*. 2016;34:1799–1803. doi: 10.1016/j.ajem.2016.06.008
369. Biersens JJ, van der Velde EA, van Berkel M, van Zanten JJ. Submersion in The Netherlands: prognostic indicators and results of resuscitation. *Ann Emerg Med*. 1990;19:1390–1395. doi: 10.1016/s0196-0644(05)82604-6
370. Omar HR, Mirsaeidi M, Bosco G, Morgan K, Dalvi P, Helal E, Mangar D, Camporesi EM. Cardiovascular complications and mortality determinants in near drowning victims: a 5-year retrospective analysis. *J Crit Care*. 2017;37:237–239. doi: 10.1016/j.jccr.2016.09.010
371. Kruus S, Bergström L, Suutarinen T, Hyvönen R. The prognosis of near-drowned children. *Acta Paediatr Scand*. 1979;68:315–322. doi: 10.1111/j.1651-2227.1979.tb05013.x
372. Graf WD, Cummings P, Quan L, Brutocao D. Predicting outcome in pediatric submersion victims. *Ann Emerg Med*. 1995;26:312–319. doi: 10.1016/s0196-0644(95)70079-x
373. Torres SF, Rodríguez M, Iolster T, Siaba Serrate A, Cruz Iturrieta C, Martínez del Valle E, Schnitzler E, Roca Rivarola M. Near drowning in a pediatric population: epidemiology and prognosis. *Arch Argent Pediatr*. 2009;107:234–240. doi: 10.1590/S0325-00752009000300011
374. Kieboom JK, Verkade HJ, Burgerhof JG, Biersens JJ, Rheeën PF, Kneyber MC, Albers MJ. Outcome after resuscitation beyond 30 minutes in drowned children with cardiac arrest and hypothermia: Dutch nationwide retrospective cohort study. *BMJ*. 2015;350:h418. doi: 10.1136/bmj.h418
375. Joanknecht L, Argent AC, van Dijk M, van As AB. Childhood drowning in South Africa: local data should inform prevention strategies. *Pediatr Surg Int*. 2015;31:123–130. doi: 10.1007/s00383-014-3637-0
376. Shenoi RP, Koerner CE, Cruz AT, Frost MH, Jones JL, Camp EA, Alam S, Fraser JJ Jr. Factors associated with poor outcome in childhood swimming pool submersions. *Pediatr Emerg Care*. 2016;32:669–674. doi: 10.1097/PEC.0000000000000678
377. Anderson KC, Roy T, Danzl DF. Submersion incidents: a review of 39 cases and development of the submersion outcome score. *J Wilderness Med Care*. 1991;2:27–36.
378. Kaukinen L. Clinical course and prognostic signs in near-drowned patients. *Ann Chir Gynaecol*. 1984;73:34–39.
379. Veenhuizen L, Haasnoot K, van Vught AJ, Biersens JJ, Thunnissen BT, Gemke RJ. [Submersion in children; the role of hypothermia and development of adult respiratory distress syndrome]. *Ned Tijdschr Geneesk*. 1994;138:906–910.
380. Quan L, Kinder D. Pediatric submersions: prehospital predictors of outcome. *Pediatrics*. 1992;90:909–913.
381. Suominen PK, Korpela RE, Silfvast TG, Olkkola KT. Does water temperature affect outcome of nearly drowned children. *Resuscitation*. 1997;35:111–115. doi: 10.1016/s0300-9572(97)00036-1
382. Panzino F, Quintillá JM, Luaces C, Pou J. [Unintentional drowning by immersion. Epidemiological profile of victims attended in 21 Spanish emergency departments]. *An Pediatr (Barc)*. 2013;78:178–184. doi: 10.1016/j.anpedi.2012.06.014
383. Tobin JM, Ramos WD, Pu Y, Wernicki PG, Quan L, Rossano JW. Bystander CPR is associated with improved neurologically favourable survival in cardiac arrest following drowning. *Resuscitation*. 2017;115:39–43. doi: 10.1016/j.resuscitation.2017.04.004
384. Fukuda T, Ohashi-Fukuda N, Hayashida K, Kukita I. Association of bystander cardiopulmonary resuscitation and neurological outcome after out-of-hospital cardiac arrest due to drowning in Japan, 2013–2016. *Resuscitation*. 2019;141:111–120. doi: 10.1016/j.resuscitation.2019.06.005
385. Fukuda T, Ohashi-Fukuda N, Hayashida K, Kondo Y, Kukita I. Bystander-initiated conventional vs compression-only cardiopulmonary resuscitation and outcomes after out-of-hospital cardiac arrest due to drowning. *Resuscitation*. 2019;145:166–174. doi: 10.1016/j.resuscitation.2019.08.026
386. Bauman BD, Louiselle A, Nygaard RM, Vakayil V, Acton R, Hess D, Saltzman D, Kreykes N, Fischer G, Louie J, et al. Treatment of hypothermic cardiac arrest in the pediatric drowning victim, a case report, and systematic review. *Pediatr Emerg Care*. 2019:Epub ahead of print. doi: 10.1097/PEC.0000000000001735
387. Romlin BS, Winberg H, Janson M, Nilsson B, Björk K, Jeppsson A, Drake G, Claesson A. Excellent Outcome With Extracorporeal Membrane Oxygenation After Accidental Profound Hypothermia (13.8°C) and Drowning. *Crit Care Med*. 2015;43:e521–e525. doi: 10.1097/CCM.0000000000001283
388. Hilmo J, Naesheim T, Gilbert M. “Nobody is dead until warm and dead”: prolonged resuscitation is warranted in arrested hypothermic victims also in remote areas—a retrospective study from northern Norway. *Resuscitation*. 2014;85:1204–1211. doi: 10.1016/j.resuscitation.2014.04.029
389. Burke CR, Chan T, Brogan TV, Lequier L, Thiagarajan RR, Rycus PT, McMullan DM. Extracorporeal life support for victims of drowning. *Resuscitation*. 2016;104:19–23. doi: 10.1016/j.resuscitation.2016.04.005
390. Weuster M, Haney A, Panholzer B, Klüter T, van der Brelie M, van Laak U, Cremer J, Haake N. The Use of Extracorporeal Membrane Oxygenation Systems in Severe Accidental Hypothermia After Drowning: A Centre Experience. *ASAIO J*. 2016;62:157–162. doi: 10.1097/MAT.0000000000000312
391. Biermann D, Köhne M, Stark V, Müller G, Kozlik-Feldmann R, Singer D, Riso A, Sachweh J. Extracorporeal life support in children: outcome of pediatric drowning victims after cardiopulmonary resuscitation: late-breaking science abstracts from the American Heart Association’s Scientific Sessions 2018 and late-breaking abstracts in resuscitation science from the Resuscitation Science Symposium 2018. *Circulation*. 2019;138:e751–e782. doi: 10.1161/CIR.0000000000000636
392. Dragann BN, Melnychuk EM, Wilson CJ, Lambert RL, Maffei FA. Resuscitation of a Pediatric Drowning in Hypothermic Cardiac Arrest. *Air Med J*. 2016;35:86–87. doi: 10.1016/j.amj.2015.12.006
393. Stachon P, Kalbhenn J, Waltersbacher S, Bode C, Staudacher D. [Near-Drowning with Good Outcome after ECMO-Therapy and Therapeutic Hypothermia Despite 20 Minutes of Anoxia and 16 Hours of Hypoxia]. *Dtsch Med Wochenschr*. 2017;142:596–600. doi: 10.1055/s-0043-101530

- 
394. Mann C, Baer W, Riedel T. Epinephrine at 25°C core body temperature and during rewarming: case report of successful infant resuscitation after cold water submersion. *Pediatr Emerg Care*. 2019:Epub ahead of print. doi: 10.1097/PEC.0000000000001766
395. White L, Rogers J, Bloomingdale M, Fahrenbruch C, Culley L, Subido C, Eisenberg M, Rea T. Dispatcher-assisted cardiopulmonary resuscitation: risks for patients not in cardiac arrest. *Circulation*. 2010;121:91–97. doi: 10.1161/CIRCULATIONAHA.109.872366
396. Haley KB, Lerner EB, Pirrallo RG, Croft H, Johnson A, Uihlein M. The frequency and consequences of cardiopulmonary resuscitation performed by bystanders on patients who are not in cardiac arrest. *Prehosp Emerg Care*. 2011;15:282–287. doi: 10.3109/10903127.2010.541981
397. Moriwaki Y, Sugiyama M, Tahara Y, Iwashita M, Kosuge T, Harunari N, Arata S, Suzuki N. Complications of bystander cardiopulmonary resuscitation for unconscious patients without cardiopulmonary arrest. *J Emerg Trauma Shock*. 2012;5:3–6. doi: 10.4103/0974-2700.93094