

Toward Metabolomic Signatures of Cardiovascular Disease

Gregory D. Lewis, MD; Robert E. Gerszten, MD

Small biochemicals are the end result of all the regulatory complexity present in a cell, tissue, or organism, including transcriptional regulation, translational regulation, and posttranslational modifications. Metabolic changes are thus the most proximal reporters of the body's response to a disease process or drug therapy. In 1971, Robinson and coworkers¹ conceived the core idea that information-rich data reflecting the functional status of a complex biological system resides in the quantitative and qualitative pattern of metabolites in body fluids. In the same year, Horning and Horning² first used the term metabolic profiling to describe the output of a gas chromatogram from a patient sample. This new approach to the quantitative metabolic profiling of large numbers of small molecules in biofluids was ultimately termed "metabonomics" by Nicholson et al³ and "metabolomics" by others.

Article see p 207

Two core technologies are used to perform metabolic profiling: nuclear magnetic resonance and tandem mass spectrometry (MS/MS), as previously reviewed in *Circulation: Cardiovascular Genetics*.⁴ Nuclear magnetic resonance requires relatively little sample preparation and is nondestructive, allowing for subsequent structural analyses. However, the method tends to have low sensitivity and can detect only highly abundant analytes. Tandem mass spectrometry (MS/MS), coupled with liquid chromatography, on the other hand, has much higher sensitivity for small molecules and is also applicable to a wide range of biological fluids (including serum, plasma, and urine). Recent advances in MS technology now enable researchers to determine analyte masses with such high precision and accuracy that metabolites can be identified unambiguously even in complex fluids.

These technologies can be used to characterize biological samples either in a targeted manner or in a pattern discovery manner. In the former, the investigator targets a predefined set of metabolites for analysis. In the latter, the investigator must analyze a complex pattern of peaks—the molecular identities of the species giving rise to the peaks are generally not known. Although the targeted approach is more limited,

the analysis is more straightforward because the biochemicals giving rise to the signals have already been identified. The pattern discovery or fingerprint approach is inherently less biased, but the need for subsequent unambiguous identification of the peaks can be difficult.

The vision for human metabolic profiling also extends from seminal studies of inborn errors of metabolism in infants. Millington and coworkers⁵ pioneered the use of MS-based methods for monitoring fatty acid oxidation, and organic and selected amino acids. Rapid identification of subjects with fatty acid oxidation disorders, organic acidemias, and aminoacidopathies through metabolic profiling has led to dietary modification and amelioration of symptoms or disease onset. It is anticipated that a global metabolomic analysis of more common diseases such as atherosclerosis might identify new biomarkers or spotlight pathways for dietary or drug modulation.

As might be expected, however, the application of metabolomics to complex cardiovascular diseases has been more difficult than for Mendelian disorders. Important progress, however, is reported in this edition of *Circulation: Cardiovascular Genetics*. Shah et al⁶ used a targeted MS/MS-based platform to profile ≈70 metabolites in subjects from the CATHGEN biorepository who underwent cardiac catheterization to evaluate suspected ischemic heart disease. Particular strengths of this study include the relatively large number of patients studied, appropriate corrections for both clinical conditions and medications that may modulate metabolic profiles, and the use of derivation and replication groups. They demonstrate that peripheral blood metabolite profiles, one enriched for branched-chain amino acid metabolites and another comprising urea cycle metabolites, add to the discriminative capability for coronary artery disease (CAD) compared with models containing only clinical variables. They specifically document improvement in the c-statistic with incorporation of metabolite factors, compared with clinical characteristics and established circulating biomarkers. Though the increment in the c-statistic is relatively modest, it underscores the capacity for metabolic profiling to provide new information on top of known risk factors. Furthermore, the group reports that a novel metabolite cluster composed of dicarboxylacylcarnitines predicts subsequent cardiovascular events (death and myocardial infarction) in individuals with existing CAD.

The study from Shah et al,⁶ which highlights branched-chain amino acid from ≈70 metabolites profiled (Factor 4 in the manuscript), is also noteworthy in the context of experimental and clinical data suggesting that certain amino acids may be markers of insulin resistance.^{7–11}

From the Cardiology Division, Massachusetts General Hospital; and Harvard Medical School, Boston, Mass.

Correspondence to Robert E. Gerszten, MD, Cardiovascular Research Center, Massachusetts General Hospital, E 8307, 149 13th St, Charlestown, MA 02129. E-mail rgerszten@partners.org

(*Circ Cardiovasc Genet*. 2010;3:119-121.)

© 2010 American Heart Association, Inc.

Circ Cardiovasc Genet is available at
<http://circgenetics.ahajournals.org>

DOI: 10.1161/CIRCGENETICS.110.954941

Furthermore, studies of branched-chain amino acid supplementation in both animals⁷ and humans,¹² including prior work by the authors of this manuscript, indicate that circulating amino acids may directly promote insulin resistance, possibly via disruption of insulin signaling in skeletal muscle.^{7,10} Shah et al⁶ appropriately adjusted their clinical analyses for diabetes, although their metabolomics platform may be identifying a more subtle metabolic syndrome. Their compelling new findings highlight potential cross talk between insulin signaling and atherosclerosis that merits future investigation both in mechanistic studies in animal models and in clinical populations.

The authors also found an association between a group of urea cycle metabolites (Factor 9) and prevalent CAD, and note that this finding may reflect increased amino acid and ammonia catabolism. Alternatively, 2 of the 4 constituents of this factor (arginine and citrulline) are key substrates in the nitric oxide synthesis pathway, perhaps suggesting altered arginine bioavailability in modulating CAD. Indeed, another recent mass spectrometry-based study found that a low arginine to citrulline plus ornithine ratio in peripheral blood predicts CAD and adverse outcomes.¹³

The study from Shah et al⁶ also highlights an unanticipated metabolite signature, enriched for small- and medium-chain dicarboxylacylcarnitines, that predicts future cardiovascular events. Carnitine and its acyl esters are essential compounds for the metabolism of fatty acids. The main function of carnitine is to assist in the transport and metabolism of fatty acids in the mitochondria, where they are oxidized as a major source of energy.¹⁴ The acyl-CoA dehydrogenases, in turn, are a family of enzymes involved in the mitochondrial β -oxidation of fatty acids. Medium-chain acyl-CoA dehydrogenase acts on fatty acyl-CoA molecules from 4 to 12 carbons in length. Deficiency in this enzyme is the most common defect observed in the process of mitochondrial β -oxidation of fatty acids and is one of the most common inherited disorders of metabolism. Affected individuals can experience severe hypoglycemia and catastrophic nervous system injury in the setting of fasting. How the metabolomic findings in adults with CAD might relate in any way to Mendelian disorders remains unknown. Future studies must also address the source of the dicarboxylacylcarnitines, the atherosclerosis-relevant cell types in which they are acting, and the functional effects of their buildup.

The metabolomic studies were performed using a targeted liquid chromatography–MS/MS-based platform. At first glance, the number of metabolites assayed may seem modest in the context of the estimated universe of total human metabolites (≈ 5000).¹⁵ However, the analytes chosen are key intermediaries in the metabolism of proteins, carbohydrates, and fats. Moreover, use of the targeted approach that incorporates isotope-labeled standards ensures the unambiguous identification of the analytes of interest. The use of standards also allows the investigators to report absolute quantities as opposed to relative quantities of these metabolites. Their rigorous methodology will thus facilitate both animal studies to manipulate levels of these metabolites in physiologically relevant contexts and future biomarker studies in other disease populations.

The relative contributions of diet and genetics to this newly described metabolic disarray also merits further investigation. Indeed, a previous study by this group has documented the heritability of metabolomic profiles in families with premature CAD. Ultimately, the integration of metabolomics findings with genetic data will provide an opportunity to study whether a biomarker also plays a role in disease pathogenesis. For a biomarker that has a causal role, the expected random distribution in a population of a polymorphism that determines high or low biomarker concentrations would be skewed in individuals depending on their disease status. Data from the so-called Mendelian randomization studies are accumulating for several biomarkers.

The study by Shah et al⁶ thus represents important progress in the integration of metabolomics toward our understanding of cardiovascular disease. Although metabolite profiling technologies are still under development, they complement other functional genomic approaches, such as high throughput genome sequencing, RNA expression analysis, and proteomics. Together, they hold great promise to transform our ability to profile samples with the goal of illuminating biology and discovering valuable clinical biomarkers.

Sources of Funding

This work was supported by the National Institutes of Health (K23HL091106 to G.D. Lewis and R01 HL072872 and R01DK081572 to R.E. Gerszten), the Donald W. Reynolds Foundation (to R.E. Gerszten), Fondation Leducq (to R.E. Gerszten), American Heart Association Fellow-to-Faculty Award (G.D. Lewis), and Established Investigator Award (R.E. Gerszten).

Disclosures

None.

References

- Pauling L, Robinson AB, Teranishi R, Cary P. Quantitative analysis of urine vapor and breath by gas-liquid partition chromatography. *Proc Natl Acad Sci U S A*. 1971;68:2374–2376.
- Horning EC, Horning MG. Metabolic profiles: gas-phase methods for analysis of metabolites. *Clin Chem*. 1971;17:802–809.
- Nicholson JK, Lindon JC, Holmes E. 'Metabonomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica*. 1999;29:1181–1189.
- Mayr M. Metabolomics: ready for the prime time? *Circ Cardiovasc Genet*. 2008;1:58–65.
- Roe CR, Millington DS, Maltby DA. Identification of 3-methylglutaryl carnitine. A new diagnostic metabolite of 3-hydroxy-3-methylglutaryl-coenzyme A lyase deficiency. *J Clin Invest*. 1986;77:1391–1394.
- Shah SH, Bain JR, Muehlbauer MJ, Stevens RD, Crosslin DR, Haynes C, Dungan J, Newby LK, Hauser ER, Ginsburg GS, Newgard CB, Kraus WE. Association of a peripheral blood metabolic profile with coronary artery disease and risk of subsequent cardiovascular events. *Circ Cardiovasc Genet*. 2010;3:207–214.
- Newgard CB, An J, Bain JR, Muehlbauer MJ, Stevens RD, Lien LF, Ha AM, Shah SH, Arlotto M, Slentz CA, Rochon J, Gallup D, Ilkayeva O, Wenner BR, Yancy WS Jr, Eisenson H, Musante G, Surwit RS, Millington DS, Butler MD, Svetkey LP. A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Cell Metab*. 2009;9:311–326.
- Huffman KM, Shah SH, Stevens RD, Bain JR, Muehlbauer M, Slentz CA, Tanner CJ, Kuchibhatla M, Houmard JA, Newgard CB, Kraus WE. Relationships between circulating metabolic intermediates and insulin action in overweight to obese, inactive men and women. *Diabetes Care*. 2009;32:1678–1683.

9. Shaham O, Wei R, Wang TJ, Ricciardi C, Lewis GD, Vasan RS, Carr SA, Thadhani R, Gerszten RE, Mootha VK. Metabolic profiling of the human response to a glucose challenge reveals distinct axes of insulin sensitivity. *Mol Syst Biol.* 2008;4:214.
10. Patti ME, Brambilla E, Luzi L, Landaker EJ, Kahn CR. Bidirectional modulation of insulin action by amino acids. *J Clin Invest.* 1998;101:1519–1529.
11. Wopereis S, Rubingh CM, van Erk MJ, Verheij ER, van Vliet T, Cnubben NH, Smilde AK, van der Greef J, van Ommen B, Hendriks HF. Metabolic profiling of the response to an oral glucose tolerance test detects subtle metabolic changes. *PLoS One.* 2009;4:e4525.
12. Krebs M, Krssak M, Bernroider E, Anderwald C, Brehm A, Meyerspeer M, Nowotny P, Roth E, Waldhausl W, Roden M. Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes.* 2002;51:599–605.
13. Tang WH, Wang Z, Cho L, Brennan DM, Hazen SL. Diminished global arginine bioavailability and increased arginine catabolism as metabolic profile of increased cardiovascular risk. *J Am Coll Cardiol.* 2009;53:2061–2067.
14. Ferdinandusse S, Denis S, Van Roermund CW, Wanders RJ, Dacremont G. Identification of the peroxisomal beta-oxidation enzymes involved in the degradation of long-chain dicarboxylic acids. *J Lipid Res.* 2004;45:1104–1111.
15. Gerszten RE, Wang TJ. The search for new cardiovascular biomarkers. *Nature.* 2008;451:949–952.

KEY WORDS: Editorials ■ cardiovascular diseases ■ genetics ■ metabolomics
■ functional genomics ■ atherosclerosis