

# The Stability of the Blood Oxygenation Level-Dependent Functional MRI Response to Motor Tasks Is Altered in Patients With Chronic Ischemic Stroke

Kelley C. Mazzetto-Betti, MSci; Renata F. Leoni, MSci; Octavio M. Pontes-Neto, MD, PhD; Antonio C. Santos, MD, PhD; Joao P. Leite, MD, PhD; Afonso C. Silva, PhD; Draulio B. de Araujo, PhD

**Background and Purpose**—Functional MRI is a powerful tool to investigate recovery of brain function in patients with stroke. An inherent assumption in functional MRI data analysis is that the blood oxygenation level-dependent (BOLD) signal is stable over the course of the examination. In this study, we evaluated the validity of such assumption in patients with chronic stroke.

**Methods**—Fifteen patients performed a simple motor task with repeated epochs using the paretic and the unaffected hand in separate runs. The corresponding BOLD signal time courses were extracted from the primary and supplementary motor areas of both hemispheres. Statistical maps were obtained by the conventional General Linear Model and by a parametric General Linear Model.

**Results**—Stable BOLD amplitude was observed when the task was executed with the unaffected hand. Conversely, the BOLD signal amplitude in both primary and supplementary motor areas was progressively attenuated in every patient when the task was executed with the paretic hand. The conventional General Linear Model analysis failed to detect brain activation during movement of the paretic hand. However, the proposed parametric General Linear Model corrected the misdetection problem and showed robust activation in both primary and supplementary motor areas.

**Conclusions**—The use of data analysis tools that are built on the premise of a stable BOLD signal may lead to misdetection of functional regions and underestimation of brain activity in patients with stroke. The present data urge the use of caution when relying on the BOLD response as a marker of brain reorganization in patients with stroke. (*Stroke*. 2010; 41:1921-1926.)

**Key Words:** BOLD signal instability ■ cortical plasticity ■ patients with stroke ■ primary motor cortex ■ supplementary motor cortex

In recent years, functional neuroimaging has made significant contributions to the understanding of neurophysiological and neuropathological processes that occur after an ischemic brain insult.<sup>1,2</sup> Among all neuroimaging techniques, functional MRI (fMRI) has clearly proven advantageous due to its noninvasiveness and good spatiotemporal resolution. Furthermore, the vast literature of fMRI studies in the healthy brain can provide the normal standard against which studies of neuropathological cases can be compared such as stroke. Most of such studies are based on the blood oxygenation level-dependent (BOLD) mechanism, which relies on net changes in the oxidative state of hemoglobin during neural activity. As regional cerebral blood flow increases more than the increase in the regional cerebral metabolic rate of oxygen, a net reduction of the local concentration of deoxyhemoglo-

bin results, leading to local MRI signal increases of T2\*-weighted images.

To properly perform fMRI studies using standard block design paradigms, one needs to assume that the BOLD signal is stable throughout data acquisition.<sup>3,4</sup> Among other factors, this stability is highly dependent on a normal coupling between neural activity and vascular response. However, it is well known that neurovascular coupling is disrupted in pathological diseases of the brain.<sup>5,6</sup> Therefore, the assumption of BOLD stability may not be valid in fMRI studies of patients with stroke, leading to misinterpretation of results obtained with conventional statistical analysis such as the General Linear Model (GLM).

The purpose of the present study was 3-fold: (1) to evaluate the dynamic stability of the BOLD signal amplitude in the

Received May 13, 2010; accepted June 2, 2010.

From the Department of Neuroscience and Behavioral Sciences (K.C.M.-B., O.M.P.-N., A.C.S., J.P.L., D.B.d.A.), FMRP, University of Sao Paulo, Ribeirao Preto, Brazil, the Department of Physics and Mathematics (R.F.L., D.B.d.A.), FFCLRP, University of Sao Paulo, Ribeirao Preto, Brazil, the Cerebral Microcirculation Unit (R.F.L., A.C.S.), Laboratory of Functional and Molecular Imaging, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Md; the Edmond and Lily Safra International Institute of Neuroscience of Natal (D.B.d.A.), Natal, Brazil; and Onofre Lopes University Hospital (D.B.d.A.), Federal University of Rio Grande do Norte, Natal, Brazil.

Correspondence to Draulio B. de Araujo, PhD, Onofre Lopes University Hospital (HUOL), Coordenacao do CDI, Av. Nilo Pecanha, 620, 59012-300, Natal, RN, Brazil. E-mail draulio@ufrnet.br, draulio@natalneuro.org.br

© 2010 American Heart Association, Inc.

*Stroke* is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.110.590471

**Table 1. Demographic and Clinical Details of the 15 Chronic Stroke Survivors Evaluated in This Study**

Case No.	Gender	Age, Years	Time From Stroke, Years	Anatomic Location	Paretic Side	Motor Impairment (Fugl-Meyer Score)
1	M	61	10	Corona radiata	Left	69
2	F	57	1.5	Corona radiata	Left	84
3	M	64	1	Inferior frontal gyrus	Right	59
4	F	38	1.5	Internal capsule	Left	89
5	M	67	1.3	Postcentral gyrus	Right	74
6	M	71	1	Corona radiata	Left	42
7	M	69	9	Superior temporal gyrus	Right	62
8	M	58	2	Angular gyrus	Left	68
9	F	57	1.5	Corona radiata	Left	82
10	M	55	4.5	Corona radiata	Right	73
11	F	48	2	Internal capsule	Left	75
12	M	55	5	Corona radiata	Right	69
13	F	45	2	Supramarginal gyrus	Right	57
14	F	64	1.5	Inferior parietal lobule	Left	63
15	F	43	3	Internal capsule	Left	51

M indicates male; F, female.

primary motor cortex (M1) and supplementary motor area (SMA) in response to a motor task performed by patients with chronic stroke in a single fMRI run; (2) to compare the BOLD response to movement of the paretic versus the unaffected hand in these patients; and (3) to evaluate the performance of the conventional GLM and to propose an alternative strategy to analyze fMRI of patients with stroke.

## Materials and Methods

### Subjects

Fifteen patients with chronic ischemic stroke in the unilateral middle cerebral artery territory (7 women; mean age, 57 years) were included in the study. Two were excluded from analyses due to uncorrectable amounts of head movement during the MRI acquisition, leaving 13 participants (5 women) in the final data set. The study was approved by the local Research Ethics Committee. All patients gave their written, informed consent.


For each patient, the affected territory of the middle cerebral artery stroke was evaluated by CT and MRI. The motor function was

evaluated by the Fugl-Meyer motor assessment using a validated scale that varies from 0 (most impairment) to 100 (no motor impairment).<sup>7</sup> Demographic and clinical details of relevance are presented in Table 1. None of the patients presented carotid or intracranial stenosis.

### MRI Acquisition

MR images were acquired in a 1.5-T scanner (Magnetom Vision; Siemens). The functional data set was acquired with an echoplanar sequence (TR/TE=4500/66.0 ms, flip angle=90°, voxel size=1.64 mm×1.64 mm×5.00 mm). Also, a high-resolution T1-weighted image (1 mm<sup>3</sup>) was acquired using a 3-dimensional fast spoiled gradient echo (TR/TE=9.7/4.0 ms, flip angle=12°).

A block paradigm was designed with 6 blocks of rest (27 seconds) intercalated by 5 blocks of activity (27 seconds) when the patient was cued to open and close her hand. It consisted of 2 runs, 1 with the paretic and the other with the unaffected hand. Before being placed in the MRI scanner, each patient performed a “dry run” to ensure task compliance and to guarantee a stable frequency of movement throughout the examination.



**Figure 1.** Two predictors created to extract the modulated fMRI response. Light gray columns correspond to periods when the motor task was being performed and the dark gray columns correspond to rest periods. (A) The top row predictor is commonly used in block paradigm fMRI statistical analysis and is designed to be stable throughout task performance; (B) the predictor at the bottom row is weighted to take into account progressive attenuation of the fMRI response in each epoch.

**Table 2. The Average BOLD Signal (Percent Change) in Response to the Motor Task Performed With the Paretic or Unaffected Hand\***

	Paretic Hand		Unaffected Hand		<i>P</i>
	Mean	SD	Mean	SD	
Mean of 5 blocks					
BOLD (percent signal change)					
M1	2.24	0.81	3.08	0.70	0.024
SMA	1.85	0.86	2.71	0.89	0.042
Mean of the last block					
BOLD (percent signal change)					
M1	1.58	0.76	2.94	0.91	0.002
SMA	1.43	0.63	2.68	0.80	0.027

\*Mean values were obtained from the average of all patients, for a given hemisphere, over all 5 blocks and only the last block. A paired Student *t* test was applied to inspect for statistical difference across hemispheres.

### Image Processing and Analysis

Regions of interest were chosen over M1 and SMA of each hemisphere of all patients. The average amplitude of the BOLD response was computed in every region of interest for each hand (unaffected and paretic). Absolute BOLD amplitude differences between the 2 hemispheres were evaluated using a Student *t* test. Statistical differences along task performance were evaluated using a parametric 1-way analysis of variance test with Bonferroni post hoc correction for multiple comparisons (5 time points). Unless otherwise specified, statistical significance was taken at  $P < 0.05$ .

fMRI was analyzed in BrainVoyager QX 1.9 (Brain Innovation, Maastricht, The Netherlands). Preprocessing included 3-dimensional motion correction, linear trend removal, temporal high-pass filtering at 0.01 Hz, and slice-scan-time correction with a sinc interpolation. To improve anatomic localization, statistical maps were coregistered onto high-resolution T1 images.

fMRI statistical maps were computed by 2 methods: a conventional GLM and a parametric modulation method capable of detecting data trending. Specifically, parametric GLM includes 2 predictors to explain the variance of the data. The first 1 is based on the conventional hemodynamic response function with no weighting. The second presents a hemodynamic model weighted according to a specific modulation (Figure 1). Both maps were corrected for multiple comparisons using a false discovery rate with a threshold set to  $q(\text{FDR}) < 0.05$ .


### Results

All patients presented stroke of the region supplied by the middle cerebral artery of either the right or left hemisphere and the time of stroke varied from 1 to 10 years (mean, 3.25 years). Fugl-Meyer evaluation showed a mean score of 69.5, varying from 42 to 89, which is considered severe motor impairment<sup>7</sup> (Table 1).


To investigate for asymmetries in signal amplitude, the average BOLD response (percent change) was calculated for both M1 and SMA of each hemisphere (affected and unaffected). Table 2 shows the mean BOLD signal values obtained from the average of all patients in each region of interest. To allow comparison across patients, signal amplitude was normalized to the first activation block. Signal asymmetry was statistically significant when all 5 blocks of task were taken into account ( $P < 0.024$  for M1 and  $P < 0.042$  for SMA), and it increases when the average was computed with respect only to the last block ( $P < 0.002$  for M1 and  $P < 0.027$  for SMA).

To further investigate this finding, and to demonstrate that this is representative, the amplitude and standard deviation of the normalized BOLD response was calculated for each of the 5 task epochs averaged across all patients (Figure 2) for each hemisphere and every region of interest. Figure 2 shows the results obtained in SMA and M1 for the unaffected (Figure 2A) and paretic hands (Figure 2B). There was no significant difference in the response of the unaffected hand showing that the BOLD amplitude is preserved across repeated epochs in either M1 (Figure 2A, left) or SMA (Figure 2A, right). Conversely, the BOLD response of the paretic hand progressively attenuates in M1 (Figure 2B, left) as well as in SMA (Figure 2B, right). The responses in M1 in the last 2 epochs are significantly smaller than the response in either the first or second epochs ( $P < 0.001$ ). The same trend was found in SMA, in which the amplitudes to the fourth and fifth epochs are significantly smaller than the initial 1 ( $P < 0.05$ ).

To evaluate the impact of such attenuation over the computation of fMRI statistical maps, the conventional GLM and parametric GLM methods were applied to the data of every patient. Results from a representative patient are shown in Figures 3 and 4.



**Figure 2.** Amplitude and standard deviation of the mean BOLD signal amplitude in SMA (left) and M1 (right) throughout a single run of 5 epochs (x axis) normalized to the first epoch and averaged across patients obtained in response to movement of the unaffected hand (A) and the paretic hand (B). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.0001$  compared with the first response. ††† $P < 0.001$  compared with the second response. ‡ $P < 0.05$  compared with the third response.




**Figure 3.** Functional activation maps obtained from a representative patient when moving the unaffected hand. Statistically significant results ( $q(\text{FDR}) < 0.05$ ) were found in M1 and SMA by both analysis methods: the conventional GLM (A) and the parametric modulated GLM (B). The BOLD signal time courses show a stable response throughout task execution both in M1 as well as in SMA.

Figure 3 presents statistical maps obtained by GLM (Figure 3A) and parametric GLM (Figure 3B) when the patient was moving the unaffected hand and the extracted BOLD time courses from M1 and SMA. The functional maps are quite similar, demonstrating that both analysis methods provide robust detection of activation in M1 and SMA in response to the task. Moreover, it is clear that the BOLD response is stable throughout the 5 epochs comprising movement of the unaffected hand.

Figure 4 shows the functional maps and corresponding BOLD time series obtained when the same patient moved the paretic hand. The conventional GLM failed to detect activation in M1 and SMA (Figure 4A). Conversely, when parametric GLM was applied, robust activation in both M1 and SMA was observed (Figure 4B). The fMRI time courses in both regions show a progressive and significant attenuation of the BOLD response to movement of the paretic hand.

### Discussion

In the present study, the stability of the BOLD response was quantitatively analyzed across repeated epochs when patients with chronic stroke performed a motor task. The key finding of this work is the remarkable progressive attenuation of the BOLD response when stroke survivors were moving their paretic hand.<sup>8</sup> Such short-term BOLD attenuation was consistently observed in every single patient in the present study and led to a significant underestimation of the region of activation when the conventional GLM analysis was applied to the data. Besides being an unforeseen phenomenon, this result puts into question the simple use of presence or absence of activation in ipsilateral and contralateral M1 and SMA as



**Figure 4.** Functional activation maps obtained from a representative patient when moving the paretic hand. Application of the conventional GLM analysis failed to identify regions of activation in all but an extremely limited area close to M1 and SMA (A). On the other hand, the parametric modulated GLM successfully detected activation in both M1 and SMA (B). In both regions, the BOLD signal time courses show a remarkable and progressive attenuation during the execution of the motor task.

detected with standard fMRI analysis techniques such as the GLM as a marker of functional status in patients with stroke.

Although a few previous studies have already reported a decrease in BOLD signal in patients with chronic stroke, such amplitude variations were observed in different fMRI sessions.<sup>6,9–12</sup> Herein, a remarkable and consistent rapid decrease of the BOLD signal amplitude was detected during a single fMRI session in all studied patients. A similar finding has been reported by our group in a single stroke patient study.<sup>8</sup>

As noted here, individual characteristics of the BOLD response may confound interpretation of brain activation. Besides signal analysis strategies such as the one implemented in this study, many attempts have been made to calibrate the BOLD signal and so to have a more reliable evaluation of fMRI signal. Most of them are based on hypercapnia challenges,<sup>13,14</sup> which attempts to model inherent individual brain perfusion characteristics to better estimate the observed BOLD signal.<sup>15</sup>

The present results can most probably be explained according to 3 hypotheses: muscular fatigue or weakness; habituation associated with brain plasticity; or altered neurovascular coupling.

The muscular fatigue hypothesis needs to be considered in light of the simplicity of the motor task executed, although none of the patients reported difficulty in executing the task. It is well known that damage of brain tissue after stroke affects corticospinal and other supraspinal motor pathways. The reduction in neural activity along these pathways promotes a disruption of primary force control mechanisms, affecting the fast recruitment of spinal motoneurons and resulting in slower movements and a decreased muscle strength.<sup>16</sup> Previously conducted studies of motor unit activity in hemiplegic patients have suggested that motor unit



firing rates tend overall to be decreased relative to ipsilesional limbs, which affects the capacity to produce fused contraction.<sup>17</sup> Moreover, previous fMRI studies have demonstrated that, although the response in M1 is dependent on the duration and motor strength of the task, the response in SMA is invariably constant.<sup>18</sup> In our study, we found that the BOLD response in SMA was also progressively attenuated. Therefore, taken together, the simplicity of the motor task and the decreased response in SMA, it is unlikely that muscular fatigue alone can explain the observed phenomena.

A second explanation relates to a possible plasticity-induced habituation. It is well known that the brain shows a remarkable capacity of reorganization as a consequence of acquiring new skills and behaviors or learning new tasks.<sup>19</sup> In our case, the repetitive nature of the task could have induced habituation, leading to progressive attenuation of the BOLD response. Previously, Karni and colleagues conducted longitudinal fMRI studies to evaluate the effect of task performance on M1 BOLD response.<sup>20,21</sup> They observed that the performance of a repeated motor task by healthy subjects in different sessions leads to a smaller area of activation than that obtained the first time subjects executed the task. Furthermore, Carel and colleagues studied 2 groups of healthy subjects, one with and the other without daily passive training.<sup>22</sup> Two fMRI sessions were accomplished separated by one month. The untrained group showed a decrease activity of M1 and SMA, consistent with a habituation effect.<sup>22</sup> Another study from the same group failed to observe any habituation effects within a single fMRI session and only comparisons of the fMRI response to different sessions separated by 5 hours or by 49 days showed significant changes in the extent of the area of activation.<sup>23</sup> We believe pure habituation is not the most probable explanation for the observed attenuation, mainly for 2 reasons. First, only the regions responding to movement of the paretic hand, but not the ones responding to movement of the unaffected hand, exhibited habituation. Second, should habituation be the main cause for the signal decrease, one should expect the regions associated with the unaffected hand to yield the strongest signal attenuation due to a greater ability of the patients in executing the task with this hand.

A third explanation for the present results focuses on an altered neurovascular coupling. Previous studies have documented impairment of the cerebrovascular reactivity in stroke.<sup>2,11,24</sup> For instance, analysis of the BOLD response to finger tapping showed that stroke survivors present a slower rate of rise and maximum signal intensity than healthy control subjects.<sup>11</sup> Interestingly, such changes were found in both hemispheres (affected and unaffected), suggesting that the differences may be due to a diffuse functional neurovascular pathology.<sup>11</sup> Moreover, stroke survivors showed decreased BOLD amplitudes both in M1 and in SMA in the ipsilesional hemisphere regardless of whether a unilateral or a bilateral motor task was performed,<sup>2</sup> much like in the present study (Table 2). Furthermore, a recent arterial spin labeling study demonstrated a general hypoperfusion and an increased transit time in sensorimotor and supplementary motor cortices of stroke survivors.<sup>25</sup> Therefore, we believe that the present data are consistent with this hypothesis: because

hypoperfusion is temporally preserved, the maintained local increase of metabolic demand by neuronal activity would result in a progressive local attenuation of the BOLD signal.

## Conclusion

The present study reports on the existence of a progressive short-term attenuation of the BOLD signal in stroke. Regardless of the cause of the observed phenomena, the present data urge the use of caution when relying on the magnitude and spatial extent of the BOLD response in M1 and SMA of stroke survivors as a marker of brain plasticity and reorganization. Therefore, the use of new strategies should be considered, as, for instance, the analysis method proposed in this study.

## Acknowledgments

We thank the Brazilian Financial Agencies FAPESP (05/03225-7), CNPq, CAPES, and FINEP for financial support.

## Source of Funding

This research was supported in part by the Brazilian financial agencies CAPES (PROCAD-NF: 23/2010); FAPESP (05/03225-7), CNPq and FINEP and by the Intramural Research Program of the National Institutes of Health, National Institute of Neurological Disorders and Stroke (Alan P. Koretsky, Scientific Director).

## Disclosures

None.

## References

1. Calautti C, Baron JC. Functional neuroimaging studies of motor recovery after stroke in adults—a review. *Stroke*. 2003;34:1553–1566.
2. Krainik A, Hund-Georgiadis M, Zysset S, von Cramon DY. Regional impairment of cerebrovascular reactivity and BOLD signal in adults after stroke. *Stroke*. 2005;36:1146–1152.
3. de Araujo DB, Tedeschi W, Santos AC, Elias J, Neves UPC, Baffa O. Shannon entropy applied to the analysis of event-related fMRI time series. *Neuroimage*. 2003;20:311–317.
4. Sturzbecher MJ, Tedeschi W, Cabella BCT, Baffa O, Neves UPC, De Araujo DB. Non-extensive entropy and the extraction of BOLD spatial information in event-related functional MRI. *Phys Med Biol*. 2009;54:161–174.
5. Girouard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease. *J Appl Physiol*. 2006;100:328–335.
6. Rossini PM, Altamura C, Ferretti A, Vernieri F, Zappasodi F, Caulo M, Pizzella V, Del Gratta C, Romani GL, Tecchio F. Does cerebrovascular disease affect the coupling between neuronal activity and local haemodynamics? *Brain*. 2004;127:99–110.
7. Sanford J, Moreland J, Swanson LR, Stratford PW, Gowland C. Reliability of the Fugl-Meyer assessment for testing motor-performance in patients following stroke. *Phys Ther*. 1993;73:447–454.
8. Carvalho SMF, Pontes-Neto OM, Fabio SRC, Leite JP, Santos AC, de Araujo DB. Rapid BOLD fMRI signal loss in the primary motor cortex of a stroke patient. *Arq Neuropsiquiatr*. 2008;66:885–887.
9. D'Esposito M, Deouell LY, Gazzaley A. Alterations in the BOLD fMRI signal with ageing and disease: a challenge for neuroimaging. *Nat Rev Neurosci*. 2003;4:863–872.
10. Hamzei F, Knab R, Weiller C, Rother J. The influence of extra- and intracranial artery disease on the BOLD signal in fMRI. *Neuroimage*. 2003;20:1393–1399.
11. Pineiro R, Pendlebury S, Johansen-Berg H, Matthews PM. Altered hemodynamic responses in patients after subcortical stroke measured by functional MRI. *Stroke*. 2002;33:103–109.
12. Rother J, Knab R, Hamzei F, Fiehler J, Reichenbach JR, Buchel C, Weiller C. Negative dip in BOLD fMRI is caused by blood flow–oxygen consumption uncoupling in humans. *Neuroimage*. 2002;15:98–102.

13. Andrade KC, Pontes-Neto OM, Leite JP, Santos AC, Baffa O, de Araujo DB. Quantitative aspects of brain perfusion dynamic induced by BOLD fMRI. *Arq Neuropsiquiatr*. 2006;64:895–898.
14. Leoni RF, Mazzeto-Betti KC, Andrade KC, de Araujo DB. Quantitative evaluation of hemodynamic response after hypercapnia among different brain territories by fMRI. *Neuroimage*. 2008;41:1192–1198.
15. Thomason ME, Foland LC, Glover GH. Calibration of BOLD fMRI using breath holding reduces group variance during a cognitive task. *Hum Brain Mapp*. 2007;28:59–68.
16. Schieber MH, Poliakov AV. Partial inactivation of the primary motor cortex hand area: effects on individuated finger movements. *J Neurosci*. 1998;18:9038–9054.
17. Gemperline JJ, Allen S, Walk D, Rymer WZ. Characteristics of motor unit discharge in subjects with hemiparesis. *Muscle Nerve*. 1995;18:1101–1114.
18. Peck KK, Sunderland A, Peters AM, Butterworth S, Clark P, Gowland PA. Cerebral activation during a simple force production task: changes in the time course of the haemodynamic response. *Neuroreport*. 2001;12:2813–2816.
19. Draganski B, May A. Training-induced structural changes in the adult human brain. *Behav Brain Res*. 2008;192:137–142.
20. Karni A, Meyer G, Jezzard P, Adams MM, Turner R, Ungerleider LG. Functional MRI evidence for adult motor cortex plasticity during motor skill learning. *Nature*. 1995;377:155–158.
21. Karni A, Meyer G, Rey-Hipolito C, Jezzard P, Adams MM, Turner R, Ungerleider LG. The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex. *Proc Natl Acad Sci U S A*. 1998;95:861–868.
22. Carel C, Loubinoux I, Boulanouar K, Manelfe C, Rascol O, Celsis P, Chollet F. Neural substrate for the effects of passive training on sensorimotor cortical representation: a study with functional magnetic resonance imaging in healthy subjects. *J Cerebr Blood Flow Metab*. 2000;20:478–484.
23. Loubinoux I, Carel C, Alary F, Boulanouar K, Viillard G, Manelfe C, Rascol O, Celsis P, Chollet F. Within-session and between-session reproducibility of cerebral sensorimotor activation: a test–retest effect evidenced with functional magnetic resonance imaging. *J Cerebr Blood Flow Metab*. 2001;21:592–607.
24. Murata Y, Sakatani K, Hoshino T, Fujiwara N, Kano T, Nakamura S, Katayama Y. Effects of cerebral ischemia on evoked cerebral blood oxygenation responses and BOLD contrast functional MRI in stroke patients. *Stroke*. 2006;37:2514–2520.
25. Brumm KP, Perthen JE, Liu TT, Haist F, Ayalon L, Love T. An arterial spin labeling investigation of cerebral blood flow deficits in chronic stroke survivors. *Neuroimage*. 2010;51:995–1005.